

Full title: Key genes in the phenylpropanoids biosynthesis pathway have different expression patterns under various abiotic stresses in the Iranian red and green cultivars of sweet basil (*Ocimum basilicum* L.)

Informative title: Expression patterns of key genes in the phenylpropanoids biosynthesis pathway of sweet basil.

Mojgan Shahivand¹, Reza Mirderikvand^{2, 3*}, Masoud Gomarian¹, Kamran Samiei³

1-Mojgan Shahivand, PhD Candidate.

Affiliation: Department of Agronomy and Plant Breeding, Arak Branch, Islamic Azad University, Arak, Iran.

Address: Department of Agronomy and Plant Breeding, Arak Branch, Islamic Azad University.

Email: mojganshahivand@gmail.com. mobile: 09163666616.

2-***Reza Mir Drikvand**. Assistant professor. Corresponding Author:

Affiliation: Department of Agronomy and Plant Breeding, Khorramabad Branch, Islamic Azad University, Khorramabad. Iran.

Address: Department of Agronomy and Plant Breeding, Khorramabad Branch, Islamic Azad University. Email: drikvand_r@yahoo.com. Tel: 066-33120025 – Fax: 066-33120026. Mobile: 09163672770

ORCID: 0000-0001-5702-6779

1-Masoud Gomarian, Assistant professor.

Affiliation: Department of Agronomy and plant Breeding, Arak Branch, Islamic Azad University, Arak. Iran. Email: msgomarian@yahoo.com

Address: Department of Plant Breeding, Arak Branch, Islamic Azad University.

Email: msgomarian@yahoo.com. Mobile: 09188113385.

3- Kamran Samiei. Assistant Professor.

Affiliation: Department of Agriculture, Kangavar Branch, Islamic Azad University, Kangavar, Iran.

Address: Department of Agriculture, Kangavar Branch, Islamic Azad University Email: Kamransamiei@yahoo.com. Mobile: 09184251225.

Contributions

All authors contributed to the study conception and design. The laboratory experiments, data collection, and analysis were performed by Reza Mir Drikvand, Kamran Samiei and Mojgan Shahivand. Masoud Gomarian helped in survey and editing of the manuscript. All authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Consent to participate

All authors and consent.

Consent for publication

All authors consent.

key message

In the present study we investigate the expression of three key biosynthetic genes of sweet basil under five environmental stresses., for the first time we evaluated the effect of cold, drought, heat, light and salt stresses on expression of *4CL*, *C4H* and *CVOMT* genes in the Iranian red and green cultivars of sweet basil

Abstract

Phenylpropanoids are a large and important family of plant secondary metabolites. The biosynthetic pathway of phenylpropanoids is induced in plants under environmental stresses to cope with these harmful conditions. In the present study, for the first time, we identified and characterized one reference gene (*ACTIN*) and three key biosynthetic genes (*4CL*, *C4H* and *CVOMT*) in the Iranian red and green cultivars of sweet basil. Also, the expression patterns of *4CL*, *C4H* and *CVOMT* biosynthetic genes were determined for the first time in the Iranian red and green cultivars of sweet basil under cold, drought, heat, light and salt stresses. The results showed that the *ACTIN*, *4CL*, *C4H* and *CVOMT* genes identified in the Iranian cultivars are identical to other cultivars in terms of all characteristics such as ORF length, protein sequence

length, molecular weight, functional domains, lack of signal peptide, subcellular localization site, and secondary structures. Our results also revealed that the *4CL*, *C4H* and *CVOMT* biosynthetic genes have different expression in the Iranian red and green cultivars of sweet basil under abiotic stresses and their expression patterns are cultivar-dependent. The findings of this study can advance our knowledge of phenylpropanoids biosynthesis in plants under environmental stresses. These findings also can be used in plant breeding programs for stress tolerance in sweet basil and other plants.

Keywords: Abiotic stress, Gene expression pattern, *Ocimum basilicum* L., Phenylpropanoids, Sweet basil

Introduction

The medicinal plants produce a wide range of valuable therapeutic and industrial secondary metabolites (Saxena et al. 2013; Van Wyk and Wink 2018). Due to advantages such as easy availability, safety and cost-effectiveness, the medicinal plants have been widely used in various industries (Lim 2012; Saxena et al. 2013; Van Wyk and Wink 2018). Among all medicinal plants, the members of genus *Ocimum* are very valuable for their therapeutic compounds and they contain various secondary metabolites with medicinal properties (Miraj and Kiani 2016; Purushothaman et al. 2018). On the other hand, among 150 species of the genus *Ocimum*, secondary metabolites of sweet basil (*Ocimum basilicum* L.) are the most widely used in the pharmaceutical and cosmetics industries and this plant is the major essential oil-producing crop which is cultivated commercially all over the world (Nerio et al. 2010; Bilal et al. 2012; Miraj and Kiani 2016; Purushothaman et al. 2018). Sweet basil is an annual, herbaceous, aromatic medicinal plant of the Lamiaceae family (Khair-ul-Bariyah et al. 2012; Miraj and Kiani 2016).

The sweet basil contains chemical compounds such as terpenes, tannins, flavonoids and phenylpropanoids (Grayer et al. 1996; Bilal et al. 2012). Two phenylpropanoids, methyl eugenol and methyl chavicol, are the most important ingredients in sweet basil essential oil (Simon et al. 1990; Joshi 2014; Tahsili et al. 2012). The phenylpropanoids are produced from phenylalanine and tyrosine through shikimate pathway. Several enzymes are involved in phenylpropanoids biosynthesis from phenylalanine in plants (Herrmann and Weaver 1999; Haslam 2014). The phenylpropanoids biosynthetic pathway in plants begins with phenylalanine, which is then deaminated to cinnamic acid by phenylalanine ammonia-lyase (*PAL*) enzyme. The *PAL* enzyme by converting phenylalanine to cinnamic acid plays a vital role in phenylpropanoids biosynthesis in plants (Achnine et al. 2004; Swedan 2013; Khakdan et al. 2018). The second key enzyme in phenylpropanoids biosynthesis, cinnamate 4-hydroxylase (*C4H*), converts cinnamic acid to p-coumaric acid (Li et al. 2019). The *C4H* acts via an enzyme complex and connects phenylpropanoid biosynthesis to other metabolic processes (Winkel-Shirley 1999; Achnine et al. 2004). A small family of *C4H* genes exists in some plant such as *Zea mays* and *Medicago sativa*, whereas, other plants such as *Brassica napus*, *Pisum sativum*, *Arabidopsis thaliana* and *Petroselinum crispum* contain only one *C4H* gene (Frank et al. 1996; Koopmann et al. 1999; Chen et al. 2007; Andersen et al. 2008; Xu et al. 2009; Douglas et al. 2011). Over-expression of *C4H* enhanced decursinol angelate level in hairy roots of *Angelica gigas* (Park et al. 2012). The antisense suppression of *C4H* gene reduced levels of chlorogenic acid, flavonoids and lignins in

M. sativa transgenic plants (Reddy et al. 2005). Members of 4-coumarate-CoA ligase (*4CL*) gene family catalyze the conversion of 4-coumaric acid to 4-coumaroyl-CoA and participate in phenylpropanoid biosynthesis in plants (Hu et al. 1998; Hamberger and Hahlbrock 2004; Wang et al. 2016). Members of *4CL* gene family are also necessary in production of 4-cinnamic acid CoA esters that are required for the formation of plant products such as lignin, stilbenes and flavonoids (Wang et al. 2016). Over-expression of *4CL* gene enhanced flavone level in hairy roots of *Scutellaria baicalensis* (Kim et al. 2014). On the other hand, down-regulation of *4CL* reduced lignin content in transgenic switchgrass (Xu et al. 2011). The chavicol O-methyltransferase (*CVOMT*) transfers a methyl group from S-adenosylmethionine to chavicol and produces methylchavicol (Gang et al. 2002). The *CVOMT* has higher expression level in young leaves of basil compared to older leaves (Gang et al. 2002). Different chemotypes the red and green sweet basil contain various phenylpropanoid content and some of them are rich in specific phenylpropanoids (Beatovic et al. 2015; Bilal et al. 2012; Simon et al. 1990). Previous studies have reported that phenylpropanoids biosynthesis in plants is affected by growth and development, biotic and abiotic stresses (Batard et al. 1997; Solecka 1997; Bellés et al. 2008; Silva et al. 2012; Niinemets et al. 2013; Meek 2015; Deng and Lu 2017; Rastogi et al. 2019). In a study, the insect and mechanical induced stresses enhanced production of phenylpropanoids such as eugenol, eugenol acetate, dillapiol and myristicin in the leaves of *Mangifera indica* (Silva et al. 2012). In the other study, the cold, drought, flood and salt stresses changed expression of phenylpropanoids biosynthetic genes and also altered accumulation level of these compounds in *Ocimum tenuiflorum* (Rastogi et al. 2019). In the similar study, water stress increased expression of genes involved in phenylpropanoids biosynthesis such as *PAL*, *C3H*, *4CL* in *Oryza sativa* (Yang et al. 2006; Moura et al. 2010). Similar results were also observed in *Citrullus lanatus* plant under drought stress (Yoshimura et al. 2008; Moura et al. 2010). Because of the effect of environmental stresses on expression of phenylpropanoids biosynthetic genes and phenylpropanoid content, the current study was conducted to investigate the expression of three key biosynthetic genes in the Iranian red and green cultivars of sweet basil under five environmental stresses. In the present study, for the first time we evaluated the effect of cold, drought, heat, light and salt stresses on expression of *4CL*, *C4H* and *CVOMT* genes in the Iranian red and green cultivars of sweet basil.

Material and Methods

Plant Materials, Stress Treatments and Sampling

Seeds of the Iranian red and green cultivars of sweet basil were provided from seed and plant improvement institute of Iran. Seeds were surface sterilized by 5% (v/v) sodium hypochlorite and sown plastic pots containing sterilized soil mixture (40% clay and 60% peats). Pots were fed with NPK fertilizer (20:20:20) and kept at 20/20°C (light/dark) with a photoperiod of 16/8 h. The regular irrigation of pots was performed to maintain normal soil moisture level (Chang et al. 2005). Five weeks plants (7-8 leaves plants) were used to stress treatment. For cold stress, plants were kept in 10°C for 24 h and then transferred to 8°C for 24 h. For heat stress, pots containing five weeks plants were kept in 30°C for 48 h. For drought stress, the field capacity (FC) of pots was determined using the gravimetric approach (Belhassen 1997) and pots containing five weeks plants were kept in 50% FC for one week. For light stress, pots were transferred to dark and kept them in 16°C for 48 h. Finally, for salt stress, pots were irrigated with 100mM NaCl for 48 h. The control plants were well irrigated and kept in normal condition (20/20°C light/dark and

1 photoperiod of 16/8 h) until subsequent analyses. Leaves of control and stressed plants were
2 harvested and immediately frozen in liquid N₂. Frozen leaf tissues were transferred to -80°C and
3 kept for subsequent analyses.

4 5 **RNA Isolation and cDNA Synthesis for real-time PCR**

6 Frozen leaf tissues of control and stressed plants were thoroughly ground in liquid N₂ using
7 mortar and pestle. Total RNA was isolated from ground tissues using Yekta Tajhiz Azma RNA
8 isolation kit (Tehran, Iran) according to the manufacturer's instructions. The integrity of isolated
9 total RNA was evaluated using spectrophotometer and agarose gel electrophoresis. Removing of
10 genomic DNA from isolated total RNA was performed using Yekta Tajhiz Azma DNase I kit
11 (Tehran, Iran) according to the manufacturer's instructions. Two micrograms of DNase I-treated
12 total RNA was used for first strand cDNA synthesis. The cDNA was synthesized using Yekta
13 Tajhiz Azma cDNA synthesis kit (Tehran, Iran) according to the manufacturer's instructions. The
14 complete synthesis of first strand of cDNA and DNA removal from total RNA and was
15 confirmed by PCR using specific *ACTIN* primers (Table 1).

16 17 **Molecular Cloning of Biosynthetic Genes and Design of Primers**

18 To identify coding sequences of *O. basilicum* biosynthetic genes, the protein sequences of
19 known *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes were retrieved from the NCBI protein database.
20 The protein sequences of each gene were separately aligned against the EST library of *O.*
21 *basilicum* using NCBI tBLASTn tool (Johnson et al. 2008). The resulting ESTs for each gene
22 were separately pooled and assembled in CLC Genomics Workbench 9.0 software (Qiagen,
23 Valencia, USA). The resulting contigs were compared with the GenBank nr/nt database using
24 BLASTn tool and the contigs with similarity to known *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes
25 were selected for further analyses. The selected contigs were evaluated for the presence of
26 complete open reading frames (ORFs) of *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes. The complete
27 ORFs of *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes were used to design primers by Allele ID 6.0
28 software (Table 1). The complete ORFs of *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes were
29 amplified by PCR. PCR was performed using gene specific primers and *Pfu* DNA polymerase
30 (Thermo Fisher Scientific, Lenexa, USA) on cDNA template. The PCR products were tailed
31 using *Taq* DNA polymerase (SinaClon BioScience, Tehran, Iran) and cloned into pTZ57R/T
32 plasmid (Thermo Fisher Scientific, Lenexa, USA). The plasmids containing *ACTIN*, *4CL*, *C4H*,
33 and *CVOMT* gene fragments were sequenced. The results of sequencing were analyzed in CLC
34 Genomics Workbench 9.0 software (Qiagen, Valencia, USA) and compared to GenBank using
35 BLASTn tool. The specific primers for real-time PCR were designed based on the sequence of
36 identified genes. The complete coding sequences of identified genes were inserted to Allele ID
37 7.0 software and specific primers were designed (Table 1). The specificity of the designed
38 primers was evaluated by PCR (sweet basil genomic DNA as template) and NCBI Primer
39 BLAST tool.

40 41 **In silico and phylogenetic analyses**

42 In silico analyses of identified genes were performed using ORF finder, CDD, Interproscan,
43 Pfam, ExPASy ProtParam, CELLO2GO, SignalP, Phyre2, Vector NTI 10.3, and CLC Genomics
44 Workbench 9.0 tools (Gasteiger et al. 2005; McGuffin et al. 2000; Yu et al. 2014; Marchler-
45 Bauer et al. 2014; Finn et al. 2016; Jones et al. 2014; Petersen et al. 2011; Kelley et al. 2015).

Multiple sequence alignment (MSA) of protein sequences of *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes was performed using Vector NTI 10.3.

Gene expression analysis by Real-time PCR

To minimize sampling and pipetting errors, the cDNAs were diluted in distilled water. Five microliters of the diluted cDNAs (1:10) was used as template in the real-time PCR. Real-time PCR was performed using Yekta Tajhiz Azma SYBR Premix (Tehran, Iran) on a Bio-rad thermocycler. The specificity of the amplifications was evaluated by melting curve analysis and agarose gel electrophoresis. All amplification reactions were performed in three biological and three technical replications. The efficiency, reproducibility and dynamic range of amplification reactions were evaluated using a standard curve. The identified *ACTIN* gene was used as reference gene. The resulted cycle thresholds (C_t) of all genes were normalized by the C_t values of the *ACTIN* gene. The relative expression of *4CL*, *C4H* and *CVOMT* genes were determined using the $2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen 2001). The Mean of C_t values were calculated across biological and technical replicates. Statistical analysis was performed using MINITAB 14 software. The Tukey's test was used for multiple comparisons of means at $P < 0.05$.

Results

Characterization of the *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes

The PCR amplified fragments of *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes and confirmed the results of EST assembly. The gel electrophoresis showed products with size of 1,334, 1,815, 1,214, and 1631 bp for *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes, respectively. The results of sequencing showed complete ORFs with size of 1,134, 1,704, 1,071, and 1,518 bp for *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes, respectively (Table 2) (Supplementary file). The complete ORFs of *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes began with ATG as start codon and ended with TAA and TGA as stop codons (Supplementary file). The *ACTIN*, *4CL*, *C4H*, and *CVOMT* genes were completely identical in red and green cultivars. For this reason, only one sequence was included in the analyses. The identified *ACTIN* gene showed 89.07% identity to *Agastache rugosa*, 88.27% identity to *Sesamum indicum*, 87.92% identity to *Erythranthe lewisii*, 88.22% identity to *Leucas cephalotes* and 86.64% identity *Erythranthe guttatus*. The nucleotide sequence of *4CL* gene identified from Iranian cultivars of sweet basil showed 100% identity to *O. basilicum*, 92.9% identity to *O. tenuiflorum*, 85.34% identity to *Agastache rugosa*, 83.9% identity to *Salvia miltiorrhiza* and 81.72% identity to *Scutellaria baicalensis*. The nucleotide sequence of *C4H* gene displayed 100% identity to *O. basilicum*, 94.66% identity to *O. tenuiflorum*, 87.94% identity to *S. miltiorrhiza*, 86.89% identity to *Mentha spicata* and 86.36% identity to *Mentha aquatica*. The identified *CVOMT* gene exhibited 100% identity to *O. basilicum*, 98.71% identity to *Ocimum americanum*, 96.22% identity to *Ocimum africanum*, 96.12% identity to *Ocimum kilimandscharicum* and 91.67% identity *O. tenuiflorum*. These results were similar in two Iranian red and green cultivars of sweet basil.

In frame translation of complete ORFs produced proteins with length of 377, 567, 356, 505 and amino acids for *ACTIN*, *4CL*, *CVOMT*, and *C4H* genes, respectively (Table 2). The molecular weight of *ACTIN*, *4CL*, *CVOMT*, and *C4H* proteins were computed 41.664, 61.12, 39.922, and 58.044 kDa (Table 2).

The signal peptide prediction showed no signal sequence in the structure of *ACTIN*, *4CL*, *C4H*, and *CVOMT* proteins (Table 3) (Supplementary file). The subcellular localization prediction exhibited that all identified proteins are localized in the cytoplasm (Table 3) (Supplementary

file). The secondary structure prediction revealed the presence of alpha helices and beta strands with variable ratios in all identified proteins (Table 3). The search for functional domain showed the presence of NBD_sugar-kinase_HSP70 domain in *ACTIN*, 4-coumarate-CoA ligase domain in *4CL*, tans-cinnamate 4-monooxygenase domain in *C4H*, and AdoMet-MTase Dimerization domains in *CVOMT* structure (Table 3) (Supplementary file).

The protein sequence of *O. basilicum ACTIN* exhibited 97-99% identity to its orthologs in other mono- and dicotyledon plant species (Figure 1). The *O. basilicum ACTIN* showed the highest and lowest identity with *Sesamum indicum* (99%) and *Z. mays* (67%), respectively (Figure 1).

The MSA results showed that the *O. basilicum 4CL* has 67-86% identity to its orthologs in other mono- and dicotyledon plant species (Figure 2). The *O. basilicum 4CL* gene showed the highest and lowest identity with *Sesamum indicum* (86%) and *Triticum aestivum* (67%), respectively (Figure 2).

The MSA results showed that the *O. basilicum C4H* has 85-92% identity to its orthologs in other mono- and dicotyledon plant species (Figure 3). The *O. basilicum C4H* gene showed the highest and lowest identity with *Striga asiatica* (92%) and *Sorghum bicolor* (85%), respectively (Figure 3).

The protein sequence of *O. basilicum CVOMT* exhibited 42-60% identity to its orthologs in other mono- and dicotyledon plant species (Figure 4). The *O. basilicum CVOMT* showed the highest and lowest identity with *Catalpa bungei* (60%) and *Asparagus officinalis* (42%), respectively (Figure 4).

Gene expression analysis

The results of relative expression of *4CL*, *C4H*, and *CVOMT* genes in red and green cultivars of basil under different environmental stresses treatments have been shown in the figures 1, 2 and 3. Analysis of variance revealed significant differences in the expression of *4CL* gene under different stresses in both red and green cultivars of basil. In the cold, heat and light stresses similar significant changes in expression of *4CL* gene were observed in both red and green cultivars (Figure 5). The expression of *4CL* gene was significantly altered in green cultivar under drought and salt stresses, while, in the red cultivar, no significant changes were observed in expression of this gene (Figure 5). The expression level of *4CL* gene was significantly decreased under cold, heat and light stresses in both red and green cultivars (Figure 5). In drought and salt stresses, the expression level of *4CL* gene was significantly decreased in green cultivar, while, it remained unchanged in the red cultivar (Figure 5). Comparison of means showed that the changes in the expression level of *4CL* gene under different stresses were similar in both red and green cultivars except in the red cultivar exposed to salt stress (Figure 5).

The results of analysis of variance showed significant changes in the expression of *C4H* gene under different stresses in both red and green cultivars of basil. In the salt and light stresses a similar significant changes in expression of *C4H* gene was observed in both red and green cultivars (Figure 6). On the other hand, in the drought and heat stresses a different significant change in expression of *C4H* gene was observed in both red and green cultivars (Figure 6). The expression of *C4H* gene was significantly altered in red cultivar under cold stress, while, in the green cultivar, no significant changes were observed in expression of this gene (Figure 6). The expression level of *C4H* gene was significantly decreased under salt and light stresses in both red and green cultivars (Figure 6).

Analysis of variance showed significant differences in the expression of *CVOMT* gene under different stresses in both red and green cultivars of basil. The expression of *CVOMT* gene

significantly increased under the drought stress in both red and green cultivars (Figure 7). In the salt stress, the expression of *CVOMT* gene was significantly decreased in the green cultivar, but it remained unchanged in the red cultivar (Figure 7). In the cold stress, the expression of *CVOMT* gene was significantly increased in the green cultivar and decreased in the red cultivar (Figure 7). In heat and light stresses, the expression level of *CVOMT* gene was significantly decreased in both red and green cultivars. This reduction was the same in both cultivars under the light stress, but in heat stress, the green cultivar showed a much greater reduction in the expression of *CVOMT* gene (Figure 7).

Discussion

Among all species of the genus *Ocimum*, secondary metabolites of *O. basilicum* (sweet basil) are the most widely used in the pharmaceutical and cosmetics industries (Nerio et al. 2010; Bilal et al. 2012; Miraj and Kiani 2016; Purushothaman et al. 2018). In the present study, we investigated the effect of abiotic stresses on the expression of *4CL*, *C4H*, and *CVOMT* biosynthetic genes in the Iranian red and green cultivars of sweet basil. For this purpose, we first identified and characterized *4CL*, *C4H*, and *CVOMT* biosynthetic genes and *ACTIN* reference gene in the tested cultivars. In the next step, we examined the expression of these genes under the cold, drought, heat, light and salt stresses in the leaves of these cultivars.

The *ACTIN* genes encode the essential multi-functional proteins that found in all eukaryotic organisms. The *ACTIN* plays important role in many important cellular processes, including cell division, signaling, cytokinesis, cell motility, and organelle movement (Dominguez and Holmes 2011; Gunning et al. 2015). The *ACTIN* gene has been identified from different plant species. The *ACTIN* genes identified from the Iranian red and green cultivars of basil did not show any difference with other basil cultivars. Our results showed that the *ACTIN* genes identified in Iranian cultivars were identical to other cultivars in terms of all characteristics such as ORF length, protein sequence length, molecular weight, functional domains, lack of signal peptide, subcellular localization site, and secondary structures. Due to the optimized structure, the protein sequence of *ACTIN* has been highly conserved during evolution (Dominguez and Holmes 2011; Gunning et al. 2015).

The *4CL* gene catalyzes the conversion of 4-coumaric acid to 4-coumaroyl-CoA and participates in phenylpropanoid biosynthesis in plants (Hu et al. 1998; Hamberger and Hahlbrock 2004; Wang et al. 2016). The *4CL* gene has been identified from different plant species. The *4CL* genes identified from the Iranian red and green cultivars of basil did not show any difference with other basil cultivars. Our results showed that the *4CL* genes identified in Iranian cultivars were identical to other cultivars in terms of all characteristics such as ORF length, protein sequence length, molecular weight, functional domains, lack of signal peptide, subcellular localization site, and secondary structures.

Numerous studies reported that the expression of *4CL* gene involved in biosynthesis of phenylpropanoid decreased under abiotic stresses in plants (Lavhale et al. 2018). The expression of *4CL* gene decreased under wounding stress in *A. thaliana* (Soltani et al. 2006). The expression of *4CL* gene decreased in *O. basilicum* in response to drought stress (Mandoulakani et al. 2017). Our results were consistent with the previous studies. The expression of *4CL* gene significantly decreased in green cultivar under cold, drought, heat, light, and salt stresses. The expression of this gene also significantly decreased in the red cultivar under cold, heat and light stresses and it remained unchanged under drought and salinity stresses. In the heat stress, the expression of *4CL* gene in green cultivar showed a greater decrease compared to red cultivar. These results indicate

1 that the expression of *4CL* gene is not similar in different cultivars of basil and it depends on the
2 cultivar.

3 The *C4H* gene is a key enzyme in phenylpropanoids biosynthesis that catalyzes conversion of
4 cinnamic acid to p-coumaric acid (Li et al. 2019). So far, the *C4H* gene have been identified in *Z.*
5 *mays*, *M. sativa*, *B. napus*, *P. sativum*, *A. thaliana* and *P. crispum* plants (Frank et al. 1996;
6 Koopmann et al. 1999; Chen et al. 2007; Andersen et al. 2008; Xu et al. 2009; Douglas et al.
7 2011). Our results demonstrated that the *C4H* genes identified from the Iranian red and green
8 cultivars of basil were identical to other cultivars in terms of all characteristics such as ORF
9 length, protein sequence length, molecular weight, functional domains, lack of signal peptide,
10 subcellular localization site, and secondary structures. These genes did not show any difference
11 with other basil cultivars.

12 Several studies have reported that the expression of *C4H* gene in plants changes under abiotic
13 stresses (Sharma et al. 2019). The wounding, cold and salt stresses increased the expression of
14 *C4H* gene in the stem tissues of *Hibiscus cannabinus* L. (Kim et al. 2013). The expression of
15 *C4H* gene remained unchanged in the cold-stressed leaf tissues of *O. basilicum* var. keshkeni
16 luvelou (Rezaie et al. 2020). The *C4H* gene highly expressed under the wounding and salt
17 stresses in the *Carthamus tinctorius* (Sadeghi et al. 2013). The copper stress increased the
18 expression of *C4H* gene in the *Vitis vinifera* (Leng et al. 2015). The expression of *C4H* gene
19 increased under the drought stress in the *Fragaria ananassa* and *Lotus japonicas* plants (Perin et
20 al. 2019; García-Calderón et al. 2015). The salt stress increased the expression of *C4H* gene in
21 the *F. ananassa* and *Olea europaea* plants (Perin et al. 2019; Rossi et al. 2016). The expression
22 of *C4H* gene increased under the chilling stress in the *Prunus persica* (Wang et al. 2019).

23 Our results indicated that the expression of *C4H* gene show different pattern under the drought,
24 cold, and heat stresses in the red and green cultivars. The expression of *C4H* gene significantly
25 increased in green cultivar under the drought stress, while it significantly decreased in red
26 cultivar. In the cold stress, the expression of *C4H* gene significantly increased in red cultivar and
27 it remained unchanged in green cultivar. The expression of *C4H* gene significantly increased in
28 red cultivar under the heat stress, while it significantly decreased in green cultivar. The salt and
29 light stresses significantly decreased the expression of *C4H* gene in red and green cultivars. In
30 the light stress, the expression of *C4H* gene in green cultivar showed a greater decrease
31 compared to red cultivar. These results indicate that the expression of *C4H* gene is not similar in
32 different cultivars of basil and it depends on the cultivar.

33 The *CVOMT* catalyzes the conversion of chavicol to methylchavicol (Gang et al. 2002). The
34 *CVOMT* genes identified from the Iranian red and green cultivars of basil did not show any
35 difference with other basil cultivars. Our results showed that the *CVOMT* genes identified in
36 Iranian cultivars were identical to other cultivars in terms of all characteristics such as ORF
37 length, protein sequence length, molecular weight, functional domains, lack of signal peptide,
38 subcellular localization site, and secondary structures.

39 The previous studies have shown that the expression of *CVOMT* gene in *O. basilicum* increases
40 under drought stresses (Mandoulakani et al. 2017; Lavhale et al. 2018; Khakdan et al. 2017).

41 The expression of *CVOMT* gene significantly decreased in the cold-stressed leaf tissues of *O.*
42 *basilicum* var. keshkeni luvelou (Rezaie et al. 2020). In case of drought stress, our results were
43 consistent with the previous studies. The expression of *CVOMT* gene significantly increased
44 under the drought stress in both green and red cultivars. In the cold stress, the expression of
45 *CVOMT* gene significantly increased in green cultivar, while it significantly decreased in red
46 cultivar. The expression of *CVOMT* gene significantly decreased under salt stress in green

cultivar and it remained unchanged in red cultivar. In the heat stress, the expression of *CVOMT* gene in green cultivar showed a greater decrease compared to red cultivar. The light stress significantly decreased the expression of *CVOMT* gene in both red and green cultivars. These results indicate that the expression of *CVOMT* gene is not similar in different cultivars of basil and it depends on the cultivar.

The abiotic environmental stresses cause damage to plant processes, including growth and development, metabolism, photosynthesis and nutrition by inducing the production of free radicals (Sharma et al. 2019; Huang et al. 2019). Thus, plants activate protective systems, including radical scavenging processes and antioxidant compounds (such as phenylpropanoids and other phenolic compounds) to cope with stress condition (Jenks and Hasegawa 2005; Pereira 2016; Korkina 2007). Phenolic compounds act as strong radical scavenging agents due to their ability to donate electrons and hydrogen atoms (Huang et al. 2019; Sharma et al. 2019).

Previous studies have reported that different cultivars of a plant species produce different ratios of the phenolic compounds in response to environmental stresses. The different cultivars of *V. vinifera* produced different composition of phenolic compounds under drought stress (Pinasseau et al. 2017). The results of other study indicated that the phenylpropanoids accumulate at different levels in the UV-sensitive and UV-resistant cultivars of *Vaccinium corymbosum* (Escobar et al. 2017). In the similar study, the water stress had different effects on phenylpropanoids content and on expression of their biosynthetic genes of *O. basilicum* cultivars (Khakdan et al. 2017). In the study of Rossi et al. (2016), the salt-tolerant and salt-sensitive cultivars of *O. europaea* accumulated different levels of phenylpropanoid under salt stress (Rossi et al. 2016).

Different expression pattern of phenylpropanoids biosynthetic genes under environmental stresses in different cultivars of basil can be caused by structural and morphological differences, differences in cell signaling processes, genetic and epigenetic differences. Cultivars with different morphology and structure show different levels of stress resistance and also display different responses to stress. The cell signaling processes are closely related to plant morphology and differences in morphology affect signaling processes. Differences in gene expression regulation, the presence or absence of specific regulatory cis-acting elements in the promoter of biosynthetic genes differences in the number of these elements are the most important genetic differences that may exist among cultivars. Distribution of particular cis-acting elements in the promoter of biosynthetic genes can change the gene expression pattern in different cultivars. Finally, epigenetic differences such as histone modifications and chromatin remodeling are other factors that can cause different responses in cultivars.

Conclusion

To our knowledge, the current study is the first report of the identification and characterization of three phenylpropanoids biosynthetic genes in the Iranian red and green cultivars of sweet basil. Also in our study, the expression patterns of *4CL*, *C4H* and *CVOMT* biosynthetic genes were determined for the first time in the Iranian red and green cultivars of sweet basil under cold, drought, heat, light and salt stresses. Our results showed that the *4CL*, *C4H* and *CVOMT* genes identified in Iranian cultivars are identical to other cultivars in terms of all characteristics such as ORF length, protein sequence length, molecular weight, functional domains, lack of signal peptide, subcellular localization site, and secondary structures. Our results also revealed that the expression pattern of *4CL*, *C4H* and *CVOMT* biosynthetic genes is cultivar-dependent. However,

to better understand the differences among different sweet basil cultivars in response to environmental stresses, the transcriptome of cultivars should be evaluated under stress condition. The findings of the current study can advance our knowledge of phenylpropanoids biosynthesis in plants under environmental stresses. These findings also can be used in plant breeding programs for stress tolerance in sweet basil and other plants.

Compliance with ethical standards

Conflict of interest: The authors have no conflicts of interest to declare.

References

- Achnine L, Blancaflor EB, Rasmussen S, Dixon RA (2004) Colocalization of L-phenylalanine ammonia-lyase and cinnamate 4-hydroxylase for metabolic channeling in phenylpropanoid biosynthesis. *The Plant Cell* 16 (11):3098-3109
- Andersen JR, Zein I, Wenzel G, Darnhofer B, Eder J, Ouzunova M, Lübberstedt T (2008) Characterization of phenylpropanoid pathway genes within European maize (*Zea mays* L.) inbreds. *BMC Plant Biology* 8 (1):2
- Batard Y, Schalk M, Pierrel M-A, Zimmerlin A, Durst F, Werck-Reichhart D (1997) Regulation of the cinnamate 4-hydroxylase (CYP73A1) in Jerusalem artichoke tubers in response to wounding and chemical treatments. *Plant physiology* 113 (3):951-959
- Beatovic D, Krstic-Milosevic D, Trifunovic S, Siljegovic J, Glamoclija J, Ristic M, Jelacic S (2015) Chemical composition, antioxidant and antimicrobial activities of the essential oils of twelve *Ocimum basilicum* L. cultivars grown in Serbia. *Records of Natural Products* 9 (1):62
- Belhassen E (1997) Drought tolerance in higher plants: genetical, physiological and molecular biological analysis. Springer,
- Bellés JM, López-Gresa MP, Fayos J, Pallás V, Rodrigo I, Conejero V (2008) Induction of cinnamate 4-hydroxylase and phenylpropanoids in virus-infected cucumber and melon plants. *Plant Science* 174 (5):524-533
- Bilal A, Jahan N, Ahmed A, Bilal SN, Habib S, Hajra S (2012) Phytochemical and pharmacological studies on *Ocimum basilicum* Linn-A review. *International Journal of Current Research and Review* 4 (23)
- Chang X, Alderson P, Wright C (2005) Effect of temperature integration on the growth and volatile oil content of basil (*Ocimum basilicum* L.). *The Journal of Horticultural Science and Biotechnology* 80 (5):593-598
- Chen A-H, Chai Y-R, Li J-N, Chen L (2007) Molecular cloning of two genes encoding cinnamate 4-hydroxylase (C4H) from oilseed rape (*Brassica napus*). *BMB Reports* 40 (2):247-260
- Deng Y, Lu S (2017) Biosynthesis and regulation of phenylpropanoids in plants. *Critical reviews in plant sciences* 36 (4):257-290
- Dominguez R, Holmes KC (2011) Actin structure and function.

- Douglas CJ, Ehrling J, Harding SA, Joshi C, DiFazio S, Kole C (2011) Phenylpropanoid and phenolic metabolism in *Populus*: gene family structure and comparative and functional genomics. *Genetics, genomics and breeding of poplars* Enfield, NH: Science Publishers:304-326
- Escobar AL, de Oliveira Silva FM, Acevedo P, Nunes-Nesi A, Alberdi M, Reyes-Díaz M (2017) Different levels of UV-B resistance in *Vaccinium corymbosum* cultivars reveal distinct backgrounds of phenylpropanoid metabolites. *Plant Physiology and Biochemistry* 118:541-550
- Finn RD, Coghill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi M, Sangrador-Vegas A (2016) The Pfam protein families database: towards a more sustainable future. *Nucleic acids research* 44 (D1):D279-D285
- Frank MR, Deyneka JM, Schuler MA (1996) Cloning of wound-induced cytochrome P450 monooxygenases expressed in pea. *Plant physiology* 110 (3):1035-1046
- Gang DR, Lavid N, Zubieta C, Chen F, Beuerle T, Lewinsohn E, Noel JP, Pichersky E (2002) Characterization of phenylpropene O-methyltransferases from sweet basil: facile change of substrate specificity and convergent evolution within a plant O-methyltransferase family. *The Plant Cell* 14 (2):505-519
- García-Calderón M, Pons-Ferrer T, Mrazova A, Pal'ove-Balang P, Vilkova M, Pérez-Delgado CM, Vega JM, Eliašová A, Repčák M, Márquez AJ (2015) Modulation of phenolic metabolism under stress conditions in a *Lotus japonicus* mutant lacking plastidic glutamine synthetase. *Frontiers in plant science* 6:760
- Gasteiger E, Hoogland C, Gattiker A, Duvaud Se, Wilkins MR, Appel RD, Bairoch A (2005) Protein identification and analysis tools on the ExPASy server. Springer,
- Grayer RJ, Kite GC, Goldstone FJ, Bryan SE, Paton A, Putievsky E (1996) Intraspecific taxonomy and essential oil chemotypes in sweet basil, *Ocimum basilicum*. *Phytochemistry* 43 (5):1033-1039
- Gunning PW, Ghoshdastider U, Whitaker S, Popp D, Robinson RC (2015) The evolution of compositionally and functionally distinct actin filaments. *Journal of cell science* 128 (11):2009-2019
- Hamberger B, Hahlbrock K (2004) The 4-coumarate: CoA ligase gene family in *Arabidopsis thaliana* comprises one rare, sinapate-activating and three commonly occurring isoenzymes. *Proceedings of the National Academy of Sciences* 101 (7):2209-2214
- Haslam E (2014) The shikimate pathway: biosynthesis of natural products series. Elsevier,
- Herrmann KM, Weaver LM (1999) The shikimate pathway. *Annual review of plant biology* 50 (1):473-503
- Hu W-J, Kawaoka A, Tsai C-J, Lung J, Osakabe K, Ebinuma H, Chiang VL (1998) Compartmentalized expression of two structurally and functionally distinct 4-coumarate: CoA ligase genes in aspen (*Populus tremuloides*). *Proceedings of the National Academy of Sciences* 95 (9):5407-5412
- Huang H, Ullah F, Zhou D-X, Yi M, Zhao Y (2019) Mechanisms of ROS regulation of plant development and stress responses. *Frontiers in Plant Science* 10
- Jenks MA, Hasegawa PM (2005) Plant abiotic stress. Wiley Online Library,
- Johnson M, Zaretskaya I, Raytselis Y, Merezuk Y, McGinnis S, Madden TL (2008) NCBI BLAST: a better web interface. *Nucleic acids research* 36 (suppl_2):W5-W9

- 1 Jones P, Binns D, Chang H-Y, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell
- 2 A, Nuka G (2014) InterProScan 5: genome-scale protein function classification.
- 3 Bioinformatics 30 (9):1236-1240
- 4 Joshi RK (2014) Chemical composition and antimicrobial activity of the essential oil of *Ocimum*
- 5 *basilicum* L.(sweet basil) from Western Ghats of North West Karnataka, India. *Ancient*
- 6 *science of life* 33 (3):151
- 7 Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ (2015) The Phyre2 web portal for
- 8 protein modeling, prediction and analysis. *Nature protocols* 10 (6):845
- 9 Khair-ul-Bariyah S, Ahmed D, Ikram M (2012) *Ocimum basilicum*: a review on phytochemical
- 10 and pharmacological studies. *Pak J Chem* 2 (2):78-85
- 11 Khakdan F, Alizadeh H, Ranjbar M (2018) Molecular cloning, functional characterization and
- 12 expression of a drought inducible phenylalanine ammonia-lyase gene (ObPAL) from
- 13 *Ocimum basilicum* L. *Plant Physiology and Biochemistry* 130:464-472
- 14 Khakdan F, Nasiri J, Ranjbar M, Alizadeh H (2017) Water deficit stress fluctuates expression
- 15 profiles of 4Cl, C3H, COMT, CVOMT and EOMT genes involved in the biosynthetic
- 16 pathway of volatile phenylpropanoids alongside accumulation of methylchavicol and
- 17 methyleugenol in different Iranian cultivars of basil. *Journal of plant physiology* 218:74-
- 18 83
- 19 Kim J, Choi B, Natarajan S, Bae H (2013) Expression analysis of kenaf cinnamate 4-hydroxylase
- 20 (C4H) ortholog during developmental and stress responses.
- 21 Kim YS, Kim YB, Kim Y, Lee MY, Park SU (2014) Overexpression of cinnamate 4-
- 22 hydroxylase and 4-coumaroyl CoA ligase prompted flavone accumulation in *Scutellaria*
- 23 *baicalensis* hairy roots. *Natural product communications* 9 (6):1934578X1400900618
- 24 Koopmann E, Logemann E, Hahlbrock K (1999) Regulation and functional expression of
- 25 cinnamate 4-hydroxylase from parsley. *Plant physiology* 119 (1):49-56
- 26 Korkina L (2007) Phenylpropanoids as naturally occurring antioxidants: from plant defense to
- 27 human health. *Cell Mol Biol* 53 (1):15-25
- 28 Lavhale SG, Kalunke RM, Giri AP (2018) Structural, functional and evolutionary diversity of 4-
- 29 coumarate-CoA ligase in plants. *Planta* 248 (5):1063-1078
- 30 Leng X, Jia H, Sun X, Shangguan L, Mu Q, Wang B, Fang J (2015) Comparative transcriptome
- 31 analysis of grapevine in response to copper stress. *Scientific reports* 5:17749
- 32 Li X, Kim JK, Park SU (2019) Molecular cloning and characterization of rosmarinic acid
- 33 biosynthetic genes and rosmarinic acid accumulation in *Ocimum basilicum* L. *Saudi*
- 34 *journal of biological sciences* 26 (3):469-472
- 35 Lim TK (2012) *Edible medicinal and non-medicinal plants*, vol 1. Springer,
- 36 Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
- 37 quantitative PCR and the 2- $\Delta\Delta$ CT method. *methods* 25 (4):402-408
- 38 Mandoulakani BA, Eyvazpour E, Ghadimzadeh M (2017) The effect of drought stress on the
- 39 expression of key genes involved in the biosynthesis of phenylpropanoids and essential
- 40 oil components in basil (*Ocimum basilicum* L.). *Phytochemistry* 139:1-7
- 41 Marchler-Bauer A, Derbyshire MK, Gonzales NR, Lu S, Chitsaz F, Geer LY, Geer RC, He J,
- 42 Gwadz M, Hurwitz DI (2014) CDD: NCBI's conserved domain database. *Nucleic acids*
- 43 *research* 43 (D1):D222-D226
- 44 McGuffin LJ, Bryson K, Jones DT (2000) The PSIPRED protein structure prediction server.
- 45 *Bioinformatics* 16 (4):404-405

- Meek CR Arthropod Population, Phenylalanine Ammonia Lyase Activity, and Fresh Weight of Sweet Basil (*Ocimum basilicum*) as Affected by Plant Age and *Bacillus thuringiensis* Treatment. In: Proceedings of the Oklahoma Academy of Science, 2015. pp 9-17
- Miraj S, Kiani S (2016) Study of pharmacological effect of *Ocimum basilicum*: A review. *Der Pharmacia Lettre* 8 (9):276-280
- Moura JCMS, Bonine CAV, de Oliveira Fernandes Viana J, Dornelas MC, Mazzafera P (2010) Abiotic and biotic stresses and changes in the lignin content and composition in plants. *Journal of integrative plant biology* 52 (4):360-376
- Nerio LS, Olivero-Verbel J, Stashenko E (2010) Repellent activity of essential oils: a review. *Bioresource technology* 101 (1):372-378
- Niinemets Ü, Kännaste A, Copolovici L (2013) Quantitative patterns between plant volatile emissions induced by biotic stresses and the degree of damage. *Frontiers in Plant Science* 4:262
- Park NI, Park JH, Park SU (2012) Overexpression of cinnamate 4-hydroxylase gene enhances biosynthesis of decursinol angelate in *Angelica gigas* hairy roots. *Molecular biotechnology* 50 (2):114-120
- Pereira A (2016) Plant abiotic stress challenges from the changing environment. *Frontiers in plant science* 7:1123
- Perin EC, da Silva Messias R, Borowski JM, Crizel RL, Schott IB, Carvalho IR, Rombaldi CV, Galli V (2019) ABA-dependent salt and drought stress improve strawberry fruit quality. *Food chemistry* 271:516-526
- Petersen TN, Brunak S, von Heijne G, Nielsen H (2011) SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nature methods* 8 (10):785-786
- Pinasseau L, Vallverdú-Queralt A, Verbaere A, Roques M, Meudec E, Le Cunff L, Péros J-P, Ageorges A, Sommerer N, Boulet J-C (2017) Cultivar diversity of grape skin polyphenol composition and changes in response to drought investigated by LC-MS based metabolomics. *Frontiers in plant science* 8:1826
- Purushothaman B, PrasannaSrinivasan R, Suganthi P, Ranganathan B, Gimbun J, Shanmugam K (2018) A comprehensive review on *Ocimum basilicum*. *Journal of Natural Remedies* 18 (3):71-85
- Rastogi S, Shah S, Kumar R, Vashisth D, Akhtar MQ, Kumar A, Dwivedi UN, Shasany AK (2019) *Ocimum* metabolomics in response to abiotic stresses: Cold, flood, drought and salinity. *PloS one* 14 (2):e0210903
- Reddy MS, Chen F, Shadle G, Jackson L, Aljoe H, Dixon RA (2005) Targeted down-regulation of cytochrome P450 enzymes for forage quality improvement in alfalfa (*Medicago sativa* L.). *Proceedings of the National Academy of Sciences* 102 (46):16573-16578
- Rezaie R, Mandoulakani BA, Fattahi M (2020) Cold stress changes antioxidant defense system, phenylpropanoid contents and expression of genes involved in their biosynthesis in *Ocimum basilicum* L. *Scientific reports* 10 (1):1-10
- Rossi L, Borghi M, Francini A, Lin X, Xie D-Y, Sebastiani L (2016) Salt stress induces differential regulation of the phenylpropanoid pathway in *Olea europaea* cultivars Frantoio (salt-tolerant) and Leccino (salt-sensitive). *Journal of plant physiology* 204:8-15
- Sadeghi M, Dehghan S, Fischer R, Wenzel U, Vilcinskis A, Kavousi HR, Rahnamaeian M (2013) Isolation and characterization of isochorismate synthase and cinnamate 4-hydroxylase during salinity stress, wounding, and salicylic acid treatment in *Carthamus tinctorius*. *Plant signaling & behavior* 8 (11):e27335

- 1 Saxena M, Saxena J, Nema R, Singh D, Gupta A (2013) Phytochemistry of medicinal plants.
2 Journal of pharmacognosy and phytochemistry 1 (6)
- 3 Sharma A, Shahzad B, Rehman A, Bhardwaj R, Landi M, Zheng B (2019) Response of
4 phenylpropanoid pathway and the role of polyphenols in plants under abiotic stress.
5 Molecules 24 (13):2452
- 6 Silva RRD, da Câmara CA, Almeida AV, Ramos CS (2012) Biotic and abiotic stress-induced
7 phenylpropanoids in leaves of the mango (*Mangifera indica* L., Anacardiaceae). Journal
8 of the Brazilian Chemical Society 23 (2):206-211
- 9 Simon JE, Quinn J, Murray RG (1990) Basil: a source of essential oils. Advances in new
10 crops:484-489
- 11 Solecka D (1997) Role of phenylpropanoid compounds in plant responses to different stress
12 factors. Acta Physiologiae Plantarum 19 (3):257-268
- 13 Soltani BM, Ehltling J, Hamberger B, Douglas CJ (2006) Multiple cis-regulatory elements
14 regulate distinct and complex patterns of developmental and wound-induced expression
15 of *Arabidopsis thaliana* 4CL gene family members. Planta 224 (5):1226-1238
- 16 Swedan E (2013) PAL gene activity and total phenolic compounds in some members of
17 lamiaceae. Journal of Applied Sciences Research 9 (2):1222-1227
- 18 Tahsili J, Sharifi M, Behmanesh M, Pourbozorgi-Rudsari N, Ziaei M (2012) Expression of 4
19 genes in *Ocimum basilicum* and their relationship with phenylpropanoids content.
- 20 Van Wyk B-E, Wink M (2018) Medicinal plants of the world. CABI,
- 21 Wang C-H, Yu J, Cai Y-X, Zhu P-P, Liu C-Y, Zhao A-C, Lü R-H, Li M-J, Xu F-X, Yu M-D
22 (2016) Characterization and functional analysis of 4-coumarate: CoA ligase genes in
23 mulberry. PloS one 11 (5):e0155814
- 24 Wang L, Shan T, Xie B, Ling C, Shao S, Jin P, Zheng Y (2019) Glycine betaine reduces chilling
25 injury in peach fruit by enhancing phenolic and sugar metabolisms. Food chemistry
26 272:530-538
- 27 Winkel-Shirley B (1999) Evidence for enzyme complexes in the phenylpropanoid and flavonoid
28 pathways. Physiologia Plantarum 107 (1):142-149
- 29 Xu B, Escamilla-Treviño LL, Sathitsuksanoh N, Shen Z, Shen H, Percival Zhang YH, Dixon
30 RA, Zhao B (2011) Silencing of 4-coumarate: coenzyme A ligase in switchgrass leads to
31 reduced lignin content and improved fermentable sugar yields for biofuel production.
32 New phytologist 192 (3):611-625
- 33 Xu Z, Zhang D, Hu J, Zhou X, Ye X, Reichel KL, Stewart NR, Syrenne RD, Yang X, Gao P
34 Comparative genome analysis of lignin biosynthesis gene families across the plant
35 kingdom. In: BMC bioinformatics, 2009. vol 11. BioMed Central, p S3
- 36 Yang L, Wang C, Guo W, Li X, Lu M, Yu C (2006) Differential expression of cell wall related
37 genes in the elongation zone of rice roots under water deficit. Russian Journal of Plant
38 Physiology 53 (3):390-395
- 39 Yoshimura K, Masuda A, Kuwano M, Yokota A, Akashi K (2008) Programmed proteome
40 response for drought avoidance/tolerance in the root of a C3 xerophyte (wild watermelon)
41 under water deficits. Plant and Cell Physiology 49 (2):226-241
- 42 Yu C-S, Cheng C-W, Su W-C, Chang K-C, Huang S-W, Hwang J-K, Lu C-H (2014)
43 CELLO2GO: a web server for protein subCELLular LOcalization prediction with
44 functional gene ontology annotation. PLoS One 9 (6):e99368

1
2
3
4
5
6
7
8
9
10
11
12
13
14

15 Table 1. Primers used for the molecular cloning and gene expression analysis.

Primer name	Sequence (5'→3')	Product Length (bp)	Ta (°C)	
Full-4CL-F	CCTTCTTCCTCTTTCTTCTTC	1,815	60	Molecular Cloning
Full-4CL-R	ATGAACAAGGTAGGATAAATGG			
Full-CVOMT-F	AGTTCACCATATAAATTACATCCA	1,214	60	
Full-CVOMT-R	AAGTATTCGTTGCCTTTACATA			
Full-C4H-F	CACCGAAAATGGATCTTCTC	1,631	59	
Full-C4H-R	CCAATACCAACACTACAACAT			
Full-ACTIN-F	CTCTTCCGATCTCTCTCTTAA	1,334	61	
Full-ACTIN-R	CATCAGAAATCACTTCATAGACT			
4CL-F	CGACTGCGAGAACCATAG	100	53	Gene Expression Study
4CL-R	TCCTTCACCCTGTCTACG			
CVOMT-F	GGGAGAAGTTAGCAAATAAGC	163	52	
CVOMT-R	TTAGCCATCGTTCCATTACC			
C4H-F	AATCTCCACGACGCCAAG	151	57	
C4H-R	CGACTTTAGCCTCCTCTTCC			
ACTIN-F	GACAACATACAACCTCCATCAT	126	58	
ACTIN-R	GTTATCTCCTTGCTCATTCTG			
F: Forward primer; R: Reverse primer; Ta: Annealing temperature				

16
17
18
19

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17

18
19
20
21
22
23
24
25
26

Table 2. The features of identified genes.


Gene Name	cDNA length (bp)	ORF length (bp)	Protein Length (aa)	MolecularWeight (kDa)
<i>4CL</i>	1,815	1,704	567	61.12
<i>CVOMT</i>	1,214	1,071	356	39.922
<i>C4H</i>	1,631	1,518	505	58.044
<i>ACTIN</i>	1,334	1,134	377	41.664

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24


Table 3. Functional characterization of identified genes.

Gene Name	Signal sequence	Localization site	Secondary structures	Functional domain
<i>4CL</i>	No	Cytoplasmic	41% α and 21% β	4-coumarate-CoA ligase
<i>CVOMT</i>	No	Cytoplasmic	56% α and 13% β	AdoMet-MTase/ Dimerization domain
<i>C4H</i>	No	Cytoplasmic	64% α and 8% β	Tans-cinnamate 4-monooxygenase
ACTIN	No	Cytoplasmic	38% α and 23% β	NBD_sugar-kinase_HSP70

11
12
13
14
15
16
17




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16



17 Figure 2. Multiple sequence alignment of identified 4CL of *O. basilicum* with its orthologs in
 18 other mono- and dicotyledon plant species. The black and grey colors show identical and
 19 conservative amino acids, respectively. The numbers above the alignment display the position of
 20 amino acids. Gaps (-) were inserted to improve the alignment.
 21


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18



19 Figure 3. Multiple sequence alignment of identified *C4H* of *O. basilicum* with its orthologs in
20 other mono- and dicotyledon plant species. The black and grey colors show identical and
21 conservative amino acids, respectively. The numbers above the alignment display the position of
22 amino acids.

23
24
25

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18



19 Figure 4. Multiple sequence alignment of identified CVOMT of *O. basilicum* with its orthologs in
20 other mono- and dicotyledon plant species. The black and grey colors show identical and
21 conservative amino acids, respectively. The numbers above the alignment display the position of
22 amino acids. Gaps (-) were inserted to improve the alignment.

23
24
25
26




Figure 5. Expression pattern of 4CL gene in red and green cultivars of basil under different abiotic stresses. Bars show standard errors. Letters above the error show significant differences. Means with the same letter are not significantly different. The Tukey's test was used for multiple comparisons of means at $P < 0.05$.





Figure 6. Expression pattern of *C4H* gene in red and green cultivars of basil under different abiotic stresses. Bars show standard errors. Letters above the error show significant differences. Means with the same letter are not significantly different. The Tukey's test was used for multiple comparisons of means at $P < 0.05$.

1
2
3
4
5
6
7
8
9
10
11
12



13 Figure 7. Expression pattern of *CVOMT* gene in red and green cultivars of basil under different
14 abiotic stresses. Bars show standard errors. Letters above the error show significant differences.
15 Means with the same letter are not significantly different. The Tukey's test was used for multiple
16 comparisons of means at $P < 0.05$.

17
18