Full wwPDB X-ray Structure Validation Report (i)

Jul 26, 2022 – 04:40 pm BST

PDB ID	:	8AEZ
Title	:	X-ray structure of the deglycosylated receptor binding domain of Env glycoprotein of Simian Foamy virus
Deposited on	:	2022-07-14
Resolution	:	2.57 Å(reported)

This wwPDB validation report is for manuscript review

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0267
CCP4	:	7.1.010 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


For Manuscript Review

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3676 (2.60-2.56)
Clashscore	141614	4049 (2.60-2.56)
Ramachandran outliers	138981	3979 (2.60-2.56)
Sidechain outliers	138945	3979 (2.60-2.56)
RSRZ outliers	127900	3614 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	3% 85% 11%
2	B	3	33% 67%
3	C	8	25% 75%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

For Manuscript Review

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	605	-	-	-	X

For Manuscript Review

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2945 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein gp130.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	328	2711	1744	454	497	16	0	0	0

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	553	ASP	-	expression tag	UNP K7YEW5
A	554	ASP	-	expression tag	UNP K7YEW5
A	555	ASP	-	expression tag	UNP K7YEW5
A	556	ASP	-	expression tag	UNP K7YEW5
A	557	LYS	-	expression tag	UNP K7YEW5

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	3	39	22	2	15	0	0	0


- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



For Manuscript Review

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	C	8	Total C N O 94 52 2 40	0	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O 14 8 1 5	0	0
4	A	1	Total C N O 14 8 1 5	0	0
4	A	1	Total C N O 14 8 1 5	0	0
4	A	1	Total C N O 14 8 1 5	0	0
4	A	1	Total C N O 14 8 1 5	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	31	Total O 31 31	0	0

For Manuscript Review

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Envelope glycoprotein gp130



• Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



• Molecule 3: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



For Manuscript Review

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α, β, γ	99.45Å 99.45Å 120.61Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.73 – 2.57 49.73 – 2.57	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.73-2.57) 99.9 (49.73-2.57)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.53 (at 2.58Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.213 , 0.253 0.212 , 0.247	Depositor DCC
R_{free} test set	1052 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å^2)	71.1	Xtriage
Anisotropy	0.004	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	(Not available) , (Not available)	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	2945	wwPDB-VP
Average B, all atoms (Å^2)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

For Manuscript Review

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	A	0.43	0/2791	0.63	0/3793

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2711	0	2598	18	0
2	B	39	0	34	0	0
3	C	94	0	79	0	0
4	A	70	0	65	1	0
5	A	31	0	0	0	0
All	All	2945	0	2776	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

For Manuscript Review

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:HIS:HE1	1:A:341:THR:HG21	1.44	0.82
1:A:225:HIS:CE1	1:A:341:THR:HG21	2.20	0.75
1:A:546:LYS:HE2	1:A:550:THR:HG22	1.69	0.73
1:A:362:TRP:HA	1:A:391:LYS:HD2	1.83	0.60
1:A:337:GLU:O	1:A:341:THR:HG23	2.10	0.52
1:A:226:ARG:NH2	1:A:337:GLU:OE2	2.35	0.50
1:A:361:GLU:O	1:A:391:LYS:HD3	2.11	0.50
1:A:403:CYS:HA	4:A:603:NAG:H82	1.94	0.49
1:A:279:ILE:HG12	1:A:298:MET:HB3	1.95	0.48
1:A:242:THR:HG22	1:A:244:GLN:HG3	1.99	0.44
1:A:351:ASN:ND2	1:A:502:GLU:OE1	2.48	0.44
1:A:320:ASP:OD1	1:A:535:ASN:ND2	2.41	0.43
1:A:393:TYR:HB2	1:A:489:ILE:HD11	2.00	0.43
1:A:271:ALA:HB1	1:A:537:ARG:CZ	2.49	0.42
1:A:361:GLU:O	1:A:391:LYS:CD	2.68	0.42
1:A:386:VAL:O	1:A:390:ASN:HB2	2.20	0.42
1:A:530:VAL:HG22	1:A:533:MET:HE3	2.02	0.41
1:A:391:LYS:NZ	1:A:494:TYR:CE1	2.89	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	324/340 (95%)	311 (96%)	12 (4%)	1 (0%)	41 62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	325	LYS

For Manuscript Review

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	298/313 (95%)	285 (96%)	13 (4%)	28 51

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	219	ASP
1	A	242	THR
1	A	263	ASP
1	A	292	LEU
1	A	308	ARG
1	A	364	LYS
1	A	382	ARG
1	A	410	VAL
1	A	514	SER
1	A	518	SER
1	A	528	LYS
1	A	544	ASP
1	A	550	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	230	GLN
1	A	492	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

For Manuscript Review

5.5 Carbohydrates (i)

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
2	NAG	B	1	1,2	14,14,15	0.25	0	17,19,21	0.93	1 (5%)
2	NAG	B	2	2	14,14,15	0.31	0	17,19,21	0.73	1 (5%)
2	BMA	B	3	2	11,11,12	0.30	0	15,15,17	0.61	0
3	NAG	C	1	3,1	14,14,15	0.30	0	17,19,21	0.77	1 (5%)
3	NAG	C	2	3	14,14,15	0.30	0	17,19,21	0.74	0
3	BMA	C	3	3	11,11,12	0.29	0	15,15,17	0.52	0
3	MAN	C	4	3	11,11,12	0.28	0	15,15,17	0.86	1 (6%)
3	MAN	C	5	3	11,11,12	0.36	0	15,15,17	0.83	1 (6%)
3	MAN	C	6	3	11,11,12	0.33	0	15,15,17	0.82	1 (6%)
3	MAN	C	7	3	11,11,12	0.39	0	15,15,17	0.91	1 (6%)
3	MAN	C	8	3	11,11,12	0.44	0	15,15,17	0.98	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	BMA	C	3	3	-	0/2/19/22	0/1/1/1
3	MAN	C	4	3	-	0/2/19/22	0/1/1/1
3	MAN	C	5	3	-	0/2/19/22	0/1/1/1
3	MAN	C	6	3	-	0/2/19/22	0/1/1/1
3	MAN	C	7	3	-	1/2/19/22	0/1/1/1
3	MAN	C	8	3	-	0/2/19/22	0/1/1/1

For Manuscript Review

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	C	4	MAN	C1-O5-C5	3.09	116.38	112.19
3	C	7	MAN	C1-O5-C5	2.98	116.24	112.19
3	C	5	MAN	C1-O5-C5	2.94	116.18	112.19
3	C	6	MAN	C1-O5-C5	2.88	116.09	112.19
2	B	1	NAG	C1-C2-N2	-2.78	105.74	110.49
3	C	1	NAG	C1-O5-C5	2.41	115.45	112.19
3	C	8	MAN	C1-O5-C5	2.18	115.15	112.19
2	B	2	NAG	C1-C2-N2	-2.06	106.96	110.49

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	7	MAN	C4-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

For Manuscript Review


Oligosaccharide Chain B




Bond lengths



Bond angles




Torsions



Rings

For Manuscript Review

Oligosaccharide Chain C



5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

For Manuscript Review

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	NAG	A	604	1	14,14,15	0.28	0	17,19,21	0.74	1 (5%)
4	NAG	A	601	1	14,14,15	0.30	0	17,19,21	0.85	1 (5%)
4	NAG	A	605	1	14,14,15	0.32	0	17,19,21	1.03	2 (11%)
4	NAG	A	602	1	14,14,15	0.30	0	17,19,21	0.48	0
4	NAG	A	603	1	14,14,15	0.35	0	17,19,21	0.85	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	604	1	-	0/6/23/26	0/1/1/1
4	NAG	A	601	1	-	0/6/23/26	0/1/1/1
4	NAG	A	605	1	-	0/6/23/26	0/1/1/1
4	NAG	A	602	1	-	0/6/23/26	0/1/1/1
4	NAG	A	603	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
4	A	603	NAG	C1-O5-C5	3.24	116.59	112.19
4	A	605	NAG	C1-O5-C5	2.99	116.24	112.19
4	A	601	NAG	C1-O5-C5	2.87	116.08	112.19
4	A	605	NAG	O5-C1-C2	2.57	115.35	111.29
4	A	604	NAG	C1-O5-C5	2.01	114.92	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	603	NAG	1	0

For Manuscript Review

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

For Manuscript Review

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	328/340 (96%)	0.40	11 (3%) 45 41	53, 72, 107, 120	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	285	ALA	3.1
1	A	329	LEU	3.0
1	A	263	ASP	3.0
1	A	457	TYR	2.9
1	A	528	LYS	2.9
1	A	292	LEU	2.8
1	A	415	PRO	2.8
1	A	447	LEU	2.4
1	A	543	ILE	2.3
1	A	413	THR	2.3
1	A	417	CYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
2	BMA	B	3	11/12	0.65	0.30	106,108,109,109	0

Continued on next page...

For Manuscript Review


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	MAN	C	8	11/12	0.80	0.23	102,103,104,104	0
3	MAN	C	7	11/12	0.86	0.22	93,96,97,100	0
2	NAG	B	2	14/15	0.91	0.15	93,95,99,102	0
3	MAN	C	6	11/12	0.92	0.34	100,102,102,103	0
3	MAN	C	4	11/12	0.93	0.17	84,85,87,90	0
2	NAG	B	1	14/15	0.93	0.14	80,83,86,90	0
3	BMA	C	3	11/12	0.94	0.12	75,80,85,89	0
3	NAG	C	2	14/15	0.95	0.18	63,65,68,71	0
3	MAN	C	5	11/12	0.96	0.23	93,93,95,97	0
3	NAG	C	1	14/15	0.99	0.19	57,58,59,60	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain B:


$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



For Manuscript Review

Electron density around Chain C:

$2mF_o$-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
4	NAG	A	605	14/15	0.79	0.49	121,123,123,124	0
4	NAG	A	603	14/15	0.81	0.37	104,106,107,107	0
4	NAG	A	601	14/15	0.82	0.17	88,90,92,92	0
4	NAG	A	604	14/15	0.84	0.20	106,107,108,108	0
4	NAG	A	602	14/15	0.87	0.35	101,102,103,104	0

For Manuscript Review

6.5 Other polymers (i)

There are no such residues in this entry.

For Manuscript Review