

Genome-wide identification and characterization of caffeoyl-coenzyme A

O-methyltransferase genes related to Fusarium head blight response in wheat

Guang Yang^{1#}, Wenqiu Pan^{1#}, Ruoyu Zhang¹, Yan Pan¹, Qifan Guo¹, Weining Song^{1,2}, Weijun Zheng^{1*}, Xiaojun Nie^{1,2*}

¹State Key Laboratory of Crop Stress Biology in Arid Areas, College of Agronomy and Yangling Branch of China Wheat Improvement Center, Northwest A&F University, Yangling 712100, Shaanxi, China

²ICARDA-NWSUAF Joint Research Centre, Northwest A&F University, Yangling 712100, Shaanxi, China

*Corresponding author:

Weijun Zheng: zhengweijun@nwfau.edu.cn; Xiaojun Nie: small@nwsuaf.edu.cn




Figure S1. Motifs of TaCCoAOMT genes.




Figure S2. Predicted cis-element analysis in promoter regions of TaCCoAOMT genes.




Figure S3. GO enrichment for TaCCoAOMT proteins.




Figure S4. Main haplotype and frequency of TaCCoAOMT genes in A subgenome of *Triticum*.




Figure S5. Main haplotype and frequency of TaCCoAOMT genes in B subgenome of Triticum.




Figure S6. Main haplotype and frequency of TaCCoAOMT genes in D subgenome of *Triticum*.