

Article type: Original Article

Title: Clinical significance of prolymphocytes in Chinese patients with chronic lymphocytic leukemia including those treated with BTK inhibitors

Authors: *Ling-Xiao Xing ^{a, b}, *Yan Wang ^{a, b}, *Ze Jin ^{a, b}, Ye-Qin Sha ^{a, b}, Tong-Lu Qiu ^{a, b}, Li Wang ^{a, b}, Lei Fan ^{a, b}, Hua-Yuan Zhu ^{a, b}, Wei Xu ^{a, b}, #Jian-Yong Li ^{a, b}, #Yi Miao ^{a, b}

^a Department of Hematology, the First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, Nanjing, China, 210029

^b Key Laboratory of Hematology of Nanjing Medical University, Nanjing, China, 210029

*These authors contributed equally to this work.

#Corresponding author:

Yi Miao

Department of Hematology, the First Affiliated Hospital of Nanjing Medical University, Jiangsu Province Hospital, Nanjing, China, 210029

TEL: 86-25-83781120

FAX: 86-25-83781120

E-mail: miaoyi1992@hotmail.com

Jian-Yong Li

Department of Hematology, the First Affiliated Hospital of Nanjing Medical
University, Jiangsu Province Hospital, Nanjing, China, 210029


TEL: 86-25-83781120

FAX: 86-25-83781120


E-mail: lijianyonglm@126.com

Supplemental fig 1. The optimal cutoffs of prolymphocytes percentage provided by X-tile analysis to predict treatment free survival(A) and overall survival(B)

(A) treatment free survival



(B) overall survival



Supplemental fig 2. Treatment free survival(A, B) and overall survival(C, D) of patients with prolymphocytes percentage higher than 1% based on the cutoffs of 5% and 10%

