

Table 2

Variables	RAS wild type group (n=140)	RAS mutant group (n=128)	p-value
Age (mean)	66.3 (28-89)	69.3 (27-93)	0.037
Gender (male / female)	85 (60.3) / 55 (39.7)	71 (55.9) / 57 (44.1)	0.468
Tumor location (Right side / Left side)	42 (29.8) / 98 (70.2)	46 (36.2) / 82 (63.8)	0.263
Histology (Well or Moderate / Others)	126 (90.1) / 14 (9.9)	103 (80.3) / 25 (19.7)	0.023
Depth of tumor invasion (T1 · T2 / T3 · T4)	55 (40.0) / 84 (60)	39 (29.9) / 89 (70.1)	0.085
Lymph node metastasis (+ / −)	42 (30.6) / 98 (69.4)	40 (30.7) / 88 (69.3)	0.970
Lymph invasion (+ / −)	45 (31.9) / 95 (68.1)	50 (39.4) / 78 (60.6)	0.203
Venous invasion (+ / −)	97 (68.8) / 43 (31.2)	84(66.1) / 44 (33.9)	0.643
CEA level (high / normal)	35 (25.0) / 104 (75.0)	45 (35.4) / 81 (64.6)	0.063
CA19-9 level (high / normal)	14 (10.0) / 125(90.0)	20 (15.8) / 108 (84.3)	0.159
Adjuvant chemotherapy (− / +)	110 (78.6) / 30 (21.4)	107 (83.6) / 21 (16.4)	0.351