

Table 1: Association between RNF125 expression and clinicopathologic features in HCC cases

Characteristics	RNF125 expression		<i>P</i> -value
	High (n=40)	Low (n=40)	
Sex			0.402
female	6 (7.5%)	10 (12.5%)	
male	34 (42.5%)	30 (37.5%)	
Age			0.494
>60	14 (17.5%)	18 (22.5%)	
≤60	26 (32.5%)	22 (27.5%)	
TNM staging			0.013*
I-II	29 (36.2%)	17 (21.2%)	
III-IV	11 (13.8%)	23 (28.7%)	
Tumor size(cm)			0.042*
>5	25 (31.2%)	34 (42.5%)	
≤5	15 (18.8%)	6 (7.5%)	
Tumor number			0.422
multiple	7 (8.8%)	11 (13.8%)	
single	33 (41.2%)	29 (36.2%)	
Vascular invasion			0.025*
no	25 (31.2%)	14 (17.5%)	
yes	15 (18.8%)	26 (32.5%)	
AFP(ng/ml)			0.65
<400	15 (18.8%)	18 (22.5%)	
≥400	25 (31.2%)	22 (27.5%)	
HBsAg			0.18
negative	23 (28.7%)	16 (20%)	
positive	17 (21.2%)	24 (30%)	
Liver cirrhosis			0.074
no	24 (30%)	15 (18.8%)	
yes	16 (20%)	25 (31.2%)	

AFP, alpha fetal protein; HBsAg, hepatitis B surface antigen; TNM, tumor node metastasis. Chi-

square test was performed for statistical analysis. Bold italics indicate statistically significant values (* $p < 0.05$)