

Table 2: Therapeutic drugs and efficacy for delirium

	Effective patients (%)	Days(%) IDCSC Score <4	ICDSC score (before)	ICDSC score (after)	Change (%) IDCSC score	p^*
Overall (n=97)	14 (14%)	19% [28%]	6.8 [1.1]	5.6 [1.6]	-17% [22%]	<0.001
Haloperidol (n=40)	5 (13%)	21% [33%]	6.9 [1.1]	5.9 [1.6]	-14% [24%]	<0.001
Quetiapine (n=33)	2 (6%)	16% [21%]	6.7 [1.0]	5.8 [1.5]	-15% [21%]	0.002
Chlorpromazine (n=13)	4 (31%)	19% [26%]	6.9 [1.3]	4.8 [1.4]	-30% [21%]	<0.001
Olanzapine (n=8)	1 (13%)	14% [26%]	6.3 [0.8]	5.2 [1.2]	-18% [17%]	0.033
Risperidone (n=2)	1 (50%)	33% [47%]	6.5 [2.1]	5.5 [3.0]	-19% [21%]	0.374
Olanzapine + Quetiapine (n=1)	1 (100%)	67% [-]	4.0 [-]	2.7 [-]	-33% [-]	-

Values are mean [standard deviation], *unpaired t-test