

In(OTf)₃-Catalyzed Reorganization/Cycloaddition of Two Imine units and Subsequently Modular Assembly of Acridinium Photocatalysts

Jiang Nan,^{*1, 2} Guanjie Huang,¹ Shilei Liu,¹ Jing Wang,¹ Yangmin Ma,¹ and Xinjun Luan^{*3}

¹ Shaanxi Key Laboratory of Chemical Additives for Industry, College of Chemistry and Chemical Engineering, Shaanxi University of Science and Technology, Xi'an 710021 (China)

² The Youth Innovation Team of Shaanxi Universities, Shaanxi Key Laboratory of Chemical Additives for Industry, College of Chemistry and Chemical Engineering, Shaanxi University of Science and Technology, Xi'an 710021 (China)

Email: nanjiang@sust.edu.cn

³ Key Laboratory of Synthetic and Natural Functional Molecule of the Ministry of Education, College of Chemistry & Materials Science, Northwest University, Xi'an 710021 (China)

E-mail: xluan@nwu.edu.cn

Supplementary information

Table of Contents:


A. General information	2
B. Preparation of substrates	3
C. Synthesis of dihydroacridines	3
D. Synthesis of acridinium photocatalysts.....	3
E. Mechanism studies	24
F. Synthetic transformation of products.....	26
G. References.....	28
H. Spectrophotometric and electrochemical data	29
I. NMR spectra.....	53

A. General information


All reagents were used as received unless otherwise noted. Analytical thin-layer chromatography was performed with 0.25 mm coated commercial silica gel plates (TLC Silica Gel 60 F₂₅₄); visualization of the developed chromatogram was performed by fluorescence. Flash Chromatography was performed with silica gel (300-400 mesh). Proton-1 nuclear magnetic resonance (¹H NMR) data were acquired at 400 MHz on a Bruker Ascend 400 (400 MHz) spectrometer, and chemical shifts are reported in delta (δ) units, in parts per million (ppm) downfield from tetramethylsilane. Splitting patterns are designated as s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet, coupling constants *J* are quoted in Hz. Carbon-13 nuclear magnetic resonance (¹³C NMR) data were acquired at 100 MHz on a Bruker Ascend 400 spectrometer, chemical shifts are reported in ppm relative to the center line of a triplet at 77.0 ppm for CDCl₃. Fluorine-19 nuclear magnetic resonance (¹⁹F NMR) data were acquired at 376 MHz on a Bruker Ascend 400 spectrometer. Infrared spectra (IR) data were recorded on a TENSOR 27 FT-IR spectrometer and recorded in wave numbers (cm⁻¹). High resolution mass spectra were acquired on a Bruker Daltonics MicroTof-Q II mass spectrometer. UV-Vis spectra were determined on a Hitachi U-2900 spectrometer. Fluorescence spectra were acquired on a Hitachi F-4600 fluorescence spectrometer with a 10-mm quartz cuvette. The Substrates 1a-1z, 1a'-1i' ^{[1][2][3][4]} was prepared according to the literature methods, Naphthylamines and aromatic aldehydes required for the synthesis of 1a can be directly purchased from Well-known chemical reagent companies such as Energy, Innochem, Sigma-Aldrich, Acros, Alfa Aesar, or TCI.

Photoreactor Configuration

The reaction device consists of a quartz beaker with a 240V blue LED lamp winding, a stirrer with a temperature detector and an external tin box. In order to control the reaction temperature, two high-power fans are installed on both sides of the reactor. Once reaction can accommodate 5 quartz sealing tubes.




B. Preparation of substrates




A round bottom flask equipped with a stir bar was charged with the mixture of naphthalen-2-amines (0.57 g, 4.0 mmol) and MgSO_4 (0.96 g, 2 equiv.), then added 15 mL CH_2Cl_2 and stirred the mixture for 5 minutes. After that, added benzaldehydes (0.43 g, 4.0 mmol) and stirred at room temperature overnight. When the starting materials were completely converted, the reaction mixture was filtered and concentrated in vacuo. Then the residue was recrystallized to afford the product **1**.

C. Synthesis of dihydroacridines



An oven-dried vial equipped with a stir bar was charged with N-(naphthalen-2-yl)-1-phenylmethanimines **1** (0.4 mmol) and $\text{In}(\text{OTf})_3$ (22.5 mg, 0.04 mmol). The reaction mixture was placed under a positive pressure of argon and subjected to three evacuation/backfilling cycles under high vacuum. Ethylene glycol (typically, 0.8 mL) was added and the reaction mixture was stirred at 120 °C for 12 hours in an oil bath. After the reaction was completed, it was cooled to room temperature. The mixture was diluted with EtOAc and washed with saturated aqueous NaHCO_3 , then concentrated in vacuo. The residue was purified by silica gel chromatography to afford the dihydroacridine products **2**.


D. Synthesis of acridinium photocatalysts



Under argon atmosphere, $\text{Pd}(\text{OAc})_2$ (0.01 mmol), xantphos (0.02 mmol), $t\text{BuOK}$ (2 equiv.) were added to the pressure tube equipped with a stir bar and dissolved with 1 mL toluene, then aryl iodides (0.3 mmol) was slowly added and the mixture stirred for 5 minutes. After that, product **2** (0.2 mmol) dissolved with 1 mL toluene was slowly added and the mixture stirred at 120 °C for 4 hours. After the reaction was completed, it was filtered and concentrated in vacuo.


The residue from previous step was dissolved with 5 mL CH_3CN and added to a round bottom flask equipped with a stir bar, it was stirred at 0 °C and 2,3-dichloro-5,6-dicyano-1,4-benz-oquinone (90.8 mg, 2 equiv.) was slowly added. The mixture was stirred at room temperature for 2 hours. After the reaction was completed, the mixture was diluted with CH_2Cl_2 and washed with pure water, then concentrated in vacuo. The residue was

dissolved with 8 mL CH₂Cl₂ and added to a round bottom flask equipped with a stir bar, it was stirred with 10 mL NaBF₄ aqueous solution for 8 hours. After the reaction was completed, it was washed with pure water and the organic phase was concentrated in vacuo. The residue was purified by silica gel chromatography to afford the acridinium photocatalysts **8**.




(6aR,14R)-14-Phenyl-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2a)

White solid (62.1 mg, 87% yield). PE / EA = 10:1, *R*_f = 0.32. ¹H NMR (400 MHz, CDCl₃): δ 8.50 (d, *J* = 8.5 Hz, 2H), 7.85 (d, *J* = 8.1 Hz, 2H), 7.75 (d, *J* = 8.7 Hz, 2H), 7.65 (t, *J* = 7.8 Hz, 2H), 7.59 (d, *J* = 7.8 Hz, 2H), 7.42 (t, *J* = 7.6 Hz, 2H), 7.20 (t, *J* = 7.7 Hz, 2H), 7.12 (d, *J* = 8.7 Hz, 2H), 7.06 (t, *J* = 7.5 Hz, 1H), 6.80 (s, 1H), 6.46 (s, 1H). ¹³C NMR (100 MHz, CDCl₃): δ 146.1, 136.0, 132.2, 130.3, 128.8, 128.4, 128.3, 127.9, 126.9, 126.2, 123.0, 122.2, 116.8, 114.6, 39.1. IR (KBr): 3467, 3017, 1631, 1521, 1509, 1477, 1420, 1285, 1206, 1155, 871 cm⁻¹. HRMS (ESI) *m/z* Calcd for C₂₇H₂₀N [M+H]⁺ 358.1596, found 358.1589.



(6aR,14R)-14-(p-tolyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2b)


White solid (49.1 mg, 66% yield). PE / EA = 10:1, *R*_f = 0.28. ¹H NMR (400 MHz, CDCl₃): δ 8.47 (d, *J* = 8.5 Hz, 2H), 7.84 (d, *J* = 8.1 Hz, 2H), 7.75 (d, *J* = 8.6 Hz, 3H), 7.63 (t, *J* = 7.8 Hz, 3H), 7.45 (d, *J* = 7.8 Hz, 2H), 7.40 (t, *J* = 7.6 Hz, 2H), 6.99 (d, *J* = 7.7 Hz, 3H), 6.75 (s, 1H), 6.51 (s, 1H), 2.19 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 143.2, 135.9, 135.6, 132.1, 130.3, 129.0, 128.8, 128.2, 127.6, 126.8, 123.0, 122.1, 116.7, 114.8, 38.6, 20.9. IR (KBr): 3462, 3023, 2898, 1621, 1549, 1538, 1485, 1453, 1430, 1388, 1295, 1050 cm⁻¹. HRMS (ESI) *m/z* Calcd for C₂₈H₂₂N [M+H]⁺ 372.1752, found 372.1746.



(6aR,14R)-14-(4-(tert-butyl)phenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2c)


White solid (56.9 mg, 69% yield). PE / EA = 10:1, *R*_f = 0.29. ¹H NMR (400 MHz, DMSO): δ 9.61 (s, 1H), 8.64 (d, *J* = 7.8 Hz, 2H), 7.91–7.80 (m, 4H), 7.67–7.55 (m, 4H), 7.54–7.45 (m, 2H), 7.41–7.30 (m, 2H), 7.21–7.06 (m, 2H), 6.78 (s, 1H), 1.11 (s, 9H). ¹³C NMR (100 MHz, DMSO): δ

148.5, 144.7, 136.9, 132.3, 130.0, 129.0, 128.3, 127.6, 127.0, 125.3, 122.9, 122.7, 117.7, 114.2, 38.3, 34.3, 31.4. IR (KBr): 3469, 3019, 2998, 1659, 1578, 1561, 1474, 1451, 1391, 1289, 1255, 729 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{31}\text{H}_{28}\text{N}$ $[\text{M}+\text{H}]^+$ 414.2222, found 414.2213.




(6aR,14R)-14-(4-methoxyphenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2d)

White solid (48.8 mg, 63% yield). PE / EA = 8:1, R_f = 0.27. ^1H NMR (400 MHz, DMSO): δ 9.56 (s, 1H), 8.56 (d, J = 8.5 Hz, 2H), 7.85–7.71 (m, 4H), 7.57–7.46 (m, 4H), 7.38 (d, J = 8.7 Hz, 2H), 7.29 (t, J = 7.5 Hz, 2H), 6.71–6.59 (m, 3H), 3.52 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 157.7, 140.3, 136.7, 132.2, 129.9, 128.9, 128.3, 127.0, 122.9, 122.8, 117.7, 114.3, 113.9, 55.2, 37.9. IR (KBr): 3461, 3031, 2933, 1631, 1555, 1534, 1473, 1441, 1289, 1187, 746 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{28}\text{H}_{22}\text{NO}$ $[\text{M}+\text{H}]^+$ 388.1701, found 388.1695.



(6aR,14R)-14-(4-(methylthio)phenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2e)


Yellow solid (45.2 mg, 56% yield). PE / EA = 10:1, R_f = 0.32. ^1H NMR (400 MHz, CDCl_3): δ 8.41 (d, J = 8.5 Hz, 2H), 7.82 (d, J = 8.0 Hz, 2H), 7.75 (d, J = 8.7 Hz, 2H), 7.60 (t, J = 7.1 Hz, 2H), 7.44 (d, J = 8.2 Hz, 2H), 7.38 (t, J = 7.5 Hz, 2H), 7.17 (d, J = 8.7 Hz, 2H), 7.04 (d, J = 8.5 Hz, 2H), 6.71 (s, 1H), 6.55 (s, 1H), 2.33 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 143.2, 135.9, 135.6, 132.0, 130.2, 128.8, 128.4, 128.2, 126.9, 126.8, 123.0, 122.0, 116.7, 114.4, 38.5, 15.9. IR (KBr): 3458, 3031, 1626, 1523, 1515, 1445, 1413, 1289, 1062, 745 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{28}\text{H}_{22}\text{NS}$ $[\text{M}+\text{H}]^+$ 404.1473, found 404.1467.



(6aR,14R)-14-([1,1'-biphenyl]-4-yl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2f)


White solid (52.8 mg, 61% yield). PE / EA = 10:1, R_f = 0.32. ^1H NMR (400 MHz, CDCl_3): δ 8.56 (d, J = 8.6 Hz, 2H), 7.89 (d, J = 8.2 Hz, 2H), 7.78 (d, J = 8.7 Hz, 2H), 7.73–7.62 (m, 4H), 7.51–7.38 (m, 8H), 7.34 (d, J = 7.4 Hz, 1H), 7.14 (d, J = 8.7 Hz, 2H), 6.88 (s, 1H), 6.48 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 145.1, 140.9, 139.0, 136.1, 132.2, 130.4, 128.9, 128.6, 128.4, 128.2, 127.2, 127.1, 127.0, 127.0, 123.1, 122.2, 116.8, 114.6, 38.7. IR (KBr): 3451, 3021, 1642, 1577, 1518, 1470, 1422, 1268, 1042 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{33}\text{H}_{24}\text{N}$ $[\text{M}+\text{H}]^+$ 434.1909,

found 434.1902.




(6aR,14R)-14-(4-chlorophenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2g)

White solid (56.3 mg, 72% yield). PE / EA = 10:1, R_f = 0.28. ^1H NMR (400 MHz, CDCl_3): δ 8.37 (d, J = 8.4 Hz, 2H), 7.82 (d, J = 8.0 Hz, 2H), 7.75 (d, J = 8.7 Hz, 2H), 7.60 (t, J = 8.6 Hz, 2H), 7.44 (d, J = 7.4 Hz, 2H), 7.39 (t, J = 7.5 Hz, 2H), 7.17 (d, J = 8.6 Hz, 2H), 7.10 (d, J = 8.8 Hz, 2H), 6.72 (s, 1H), 6.55 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 144.5, 135.9, 131.9, 131.7, 130.2, 129.1, 128.9, 128.6, 128.4, 127.0, 123.1, 121.8, 116.7, 114.1, 38.4. IR (KBr): 3475, 3027, 1619, 1518, 1502, 1466, 1421, 1290, 733 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{27}\text{H}_{19}\text{ClN}$ $[\text{M}+\text{H}]^+$ 392.1206, found 392.1198.



(6aR,14R)-14-(4-fluorophenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2h)


Yellow solid (49.6 mg, 66% yield). PE / EA = 10:1, R_f = 0.32. ^1H NMR (400 MHz, CDCl_3): δ 8.43 (d, J = 8.5 Hz, 2H), 7.86 (d, J = 8.1 Hz, 2H), 7.76 (d, J = 8.7 Hz, 2H), 7.64 (t, J = 7.9 Hz, 2H), 7.50 (t, J = 6.9 Hz, 2H), 7.43 (t, J = 7.6 Hz, 2H), 7.14 (d, J = 8.6 Hz, 2H), 6.86 (t, J = 8.6 Hz, 2H), 6.77 (s, 1H), 6.49 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 161.1 (d, J = 244.5 Hz), 141.9 (d, J = 3.5 Hz), 135.9, 132.0, 130.2, 129.2, 129.2, 128.9, 128.5, 127.0, 123.1, 121.9, 116.8, 115.1 (d, J = 21.3 Hz), 114.4, 38.2. ^{19}F NMR (376 MHz, DMSO): δ -116.9. IR (KBr): 3442, 3012, 1614, 1569, 1528, 1467, 1414, 1349, 1248, 1207 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{27}\text{H}_{19}\text{FN}$ $[\text{M}+\text{H}]^+$ 376.1502, found 376.1443.



(6aR,14R)-14-(4-(trifluoromethyl)phenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2i)


White solid (61.2 mg, 72% yield). PE / EA = 10:1, R_f = 0.33. ^1H NMR (400 MHz, CDCl_3): δ 8.42 (d, J = 8.4 Hz, 2H), 7.88 (d, J = 8.0 Hz, 2H), 7.80 (d, J = 8.6 Hz, 2H), 7.73–7.62 (m, 4H), 7.45 (d, J = 7.6 Hz, 4H), 7.19 (d, J = 8.5 Hz, 2H), 6.84 (s, 1H), 6.58 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 149.8, 136.0, 132.0, 130.3, 129.0, 128.8, 128.3 (q, J = 32.3 Hz), 128.0, 127.1, 125.4 (q, J = 3.8 Hz), 124.2 (q, J = 270.1 Hz), 123.3, 121.8, 116.8, 113.8, 39.0. ^{19}F NMR (376 MHz, CDCl_3): δ -62.4. IR (KBr): 3450, 3017, 1634, 1558, 1512, 1460, 1438, 1332, 1261, 1212, 1108

cm⁻¹. HRMS (ESI) m/z Calcd for C₂₈H₁₉F₃N [M+H]⁺ 426.1470, found 426.1466.




(6aR,14R)-14-(4-((trifluoromethyl)thio)phenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2j)

White solid (74.1 mg, 81% yield). PE / EA = 10:1, R_f = 0.32. ¹H NMR (400 MHz, CDCl₃): δ 8.38 (d, *J* = 8.5 Hz, 2H), 7.85 (d, *J* = 8.4 Hz, 2H), 7.77 (d, *J* = 8.6 Hz, 2H), 7.63 (t, *J* = 8.4 Hz, 2H), 7.55 (d, *J* = 8.2 Hz, 2H), 7.46–7.37 (m, 4H), 7.17 (d, *J* = 8.7 Hz, 2H), 6.79 (s, 1H), 6.55 (s, 1H). ¹³C NMR (100 MHz, CDCl₃): δ 148.8, 136.3, 136.1, 132.0, 130.2, 129.5 (q, *J* = 306.3 Hz), 128.9, 128.8, 128.7, 127.1, 123.2, 121.8, 121.6, 116.8, 113.7, 38.6. ¹⁹F NMR (376 MHz, CDCl₃): δ -42.6. IR (KBr): 3469, 3018, 1623, 1562, 1254, 1229, 1038, 738 cm⁻¹. HRMS (ESI) m/z Calcd for C₂₈H₁₉F₃NS [M+H]⁺ 458.1190, found 458.1182.




4-((6aR,14R)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridin-14-yl)benzonitrile (2k)

Yellow solid (53.4 mg, 70% yield). PE / EA = 7:1, R_f = 0.29. ¹H NMR (400 MHz, DMSO): δ 9.70 (s, 1H), 8.60 (d, *J* = 8.5 Hz, 2H), 7.86–7.80 (m, 6H), 7.63–7.54 (m, 4H), 7.41 (d, *J* = 8.8 Hz, 2H), 7.33 (t, *J* = 7.6 Hz, 2H), 6.87 (s, 1H). ¹³C NMR (100 MHz, DMSO): δ 152.9, 136.9, 132.7, 132.1, 129.9, 129.0, 128.9, 128.8, 127.3, 123.2, 122.5, 119.2, 117.7, 112.9, 109.3, 38.8. IR (KBr): 3479, 3020, 2237, 1616, 1541, 1517, 1441, 1420, 1260, 1197 cm⁻¹. HRMS (ESI) m/z Calcd for C₂₈H₁₉N₂ [M+H]⁺ 383.1548, found 383.1542.




(6aR,14R)-14-(4-nitrophenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2l)

Yellow solid (57.0 mg, 71% yield). PE / EA = 4:1, R_f = 0.31. ¹H NMR (400 MHz, DMSO): δ 9.71 (s, 1H), 8.61 (d, *J* = 8.6 Hz, 2H), 8.03 (d, *J* = 8.5 Hz, 2H), 7.91 (d, *J* = 8.4 Hz, 2H), 7.87–7.78 (m, 4H), 7.57 (t, *J* = 7.9 Hz, 2H), 7.41 (d, *J* = 8.8 Hz, 2H), 7.34 (t, *J* = 7.6 Hz, 2H), 6.94 (s, 1H). ¹³C NMR (100 MHz, DMSO): δ 154.9, 146.1, 136.9, 132.0, 129.9, 129.0, 128.9, 127.3, 124.0, 123.2, 122.5, 117.7, 112.7, 38.7. IR (KBr): 3460, 3321, 3024, 1587, 1522, 1447, 1408, 1368, 1269, 1043 cm⁻¹. HRMS (ESI) m/z Calcd for C₂₇H₁₉N₂O₂ [M+H]⁺ 403.1447, found 403.1441.




(6aR,14R)-14-(4-(1H-1,2,4-triazol-1-yl)phenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2m)

Yellow solid (49.2 mg, 58% yield). PE / EA = 5:1, R_f = 0.30. ^1H NMR (400 MHz, DMSO): δ 9.67 (s, 1H), 9.06 (s, 1H), 8.65 (d, J = 8.6 Hz, 2H), 8.15 (s, 1H), 7.87–7.81 (m, 6H), 7.64–7.56 (m, 4H), 7.43 (d, J = 8.8 Hz, 2H), 7.34 (t, J = 7.7 Hz, 2H), 6.86 (s, 1H). ^{13}C NMR (100 MHz, DMSO): δ 152.6, 147.6, 142.6, 136.8, 135.1, 132.1, 129.9, 129.0, 129.0, 128.7, 127.1, 123.0, 122.6, 120.1, 117.7, 113.5, 38.3. IR (KBr): 3916, 3022, 1756, 1588, 1524, 1472, 1402, 1277, 1147, 1020 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{21}\text{N}_4$ $[\text{M}+\text{H}]^+$ 425.1766, found 425.1752.




(6aR,14R)-14-(3-bromophenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2n)

White solid (59.9 mg, 69% yield). PE / EA = 10:1, R_f = 0.31. ^1H NMR (400 MHz, CDCl_3): δ 8.41 (d, J = 8.5 Hz, 2H), 7.85 (d, J = 8.1 Hz, 2H), 7.76 (d, J = 8.7 Hz, 2H), 7.71–7.62 (m, 3H), 7.56 (d, J = 7.8 Hz, 1H), 7.43 (t, J = 7.5 Hz, 2H), 7.21–7.11 (m, 3H), 7.06 (t, J = 7.8 Hz, 1H), 6.74 (s, 1H), 6.51 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 148.3, 135.9, 131.9, 130.8, 130.2, 129.8, 129.4, 128.9, 128.6, 127.1, 126.3, 123.2, 122.8, 121.8, 116.8, 113.8, 38.9. IR (KBr): 3448, 3029, 1644, 1560, 1478, 1434, 1269, 1238, 578 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{27}\text{H}_{19}\text{BrN}$ $[\text{M}+\text{H}]^+$ 436.0701, found 436.0696.




(6aR,14R)-14-(3-iodophenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2o)

White solid (70.6 mg, 73% yield). PE / EA = 10:1, R_f = 0.30. ^1H NMR (400 MHz, CDCl_3): δ 8.37 (d, J = 8.5 Hz, 2H), 7.89–7.80 (m, 3H), 7.75 (d, J = 8.7 Hz, 2H), 7.62 (t, J = 7.4 Hz, 2H), 7.56 (d, J = 7.7 Hz, 1H), 7.44–7.32 (m, 3H), 7.14 (d, J = 8.6 Hz, 2H), 6.89 (t, J = 7.8 Hz, 1H), 6.68 (s, 1H), 6.52 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 148.4, 136.7, 135.9, 135.3, 131.9, 130.2, 123.0, 128.9, 128.6, 127.0, 127.0, 123.2, 121.8, 116.8, 113.9, 94.9, 38.9. IR (KBr): 3467, 3024, 1631, 1554, 1474, 1429, 1059, 637 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{19}\text{IN}$ $[\text{M}+\text{H}]^+$ 484.0562, found 484.0557.




(6aR,14S)-14-(2-bromophenyl)-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2p)

White solid (59.2 mg, 68% yield). PE / EA = 10:1, R_f = 0.30. ^1H NMR (400 MHz, CDCl_3): δ 8.97 (d, J = 8.6 Hz, 2H), 7.84–7.73 (m, 5H), 7.65 (t, J = 7.7 Hz, 2H), 7.45–7.35 (m, 3H), 7.18 (d, J = 8.6 Hz, 2H), 6.99 (t, J = 7.6 Hz, 1H), 6.91 (s, 1H), 6.82 (t, J = 7.6 Hz, 1H), 6.58 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 147.8, 135.8, 132.9, 132.4, 132.0, 130.2, 128.6, 128.5, 127.9, 126.8, 123.6, 123.2, 119.7, 116.9, 115.9, 38.8. IR (KBr): 3451, 3031, 1616, 1529, 1511, 1478, 1434, 1264, 1220, 657 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{27}\text{H}_{19}\text{BrN}$ $[\text{M}+\text{H}]^+$ 436.0701, found 436.0696.




(6aR,14R)-14-(naphthalen-2-yl)-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2q)

White solid (45.6 mg, 56% yield). PE / EA = 10:1, R_f = 0.32. ^1H NMR (400 MHz, CDCl_3): δ 8.54 (d, J = 8.5 Hz, 2H), 8.00 (s, 1H), 7.84–7.68 (m, 7H), 7.68–7.52 (m, 5H), 7.42–7.32 (m, 4H), 7.19 (d, J = 8.7 Hz, 2H), 6.92 (s, 1H), 6.61 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 143.4, 135.9, 133.1, 132.2, 132.0, 130.3, 128.8, 128.5, 128.4, 128.0, 127.4, 126.9, 126.7, 125.7, 125.6, 125.3, 123.0, 122.1, 116.8, 114.3, 39.4. IR (KBr): 3463, 3032, 1637, 1559, 1538, 1466, 1439, 1261, 1198 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{31}\text{H}_{22}\text{N}$ $[\text{M}+\text{H}]^+$ 408.1752, found 408.1744.




(6aR,14S)-14-(thiophen-2-yl)-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2r)

Yellow solid (50.8 mg, 70% yield). PE / EA = 10:1, R_f = 0.29. ^1H NMR (400 MHz, CDCl_3): δ 8.45 (d, J = 8.5 Hz, 2H), 7.87 (d, J = 8.0 Hz, 2H), 7.77 (d, J = 8.7 Hz, 2H), 7.66 (t, J = 8.6 Hz, 2H), 7.44 (t, J = 7.5 Hz, 2H), 7.13 (d, J = 8.7 Hz, 2H), 7.09 (s, 1H), 6.97 (d, J = 4.5 Hz, 1H), 6.78–6.70 (m, 2H), 6.59 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 149.2, 136.0, 131.9, 130.2, 128.8, 128.6, 127.1, 126.3, 124.1, 123.8, 123.2, 122.0, 116.7, 113.7, 33.8. IR (KBr): 3472, 3022, 2915, 1667, 1559, 1462, 1279, 1241, 688 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{25}\text{H}_{18}\text{NS}$ $[\text{M}+\text{H}]^+$ 364.1160, found 364.1153.




(6aR,14R)-14-(benzofuran-6-yl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2s)

White solid (41.2 mg, 52% yield). PE / EA = 10:1, R_f = 0.30. ^1H NMR (400 MHz, CDCl_3): δ 8.48 (d, J = 8.5 Hz, 2H), 7.84 (d, J = 8.1 Hz, 2H), 7.78 (d, J = 8.6 Hz, 2H), 7.64 (t, J = 8.4 Hz, 2H), 7.41 (t, J = 7.4 Hz, 2H), 7.35–7.24 (m, 2H), 7.16 (d, J = 8.7 Hz, 2H), 7.13–7.02 (m, 2H), 7.00 (s, 1H), 6.63 (s, 1H), 6.15 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 160.2, 136.2, 132.3, 130.1, 128.8, 128.6, 127.0, 123.1, 122.3, 122.2, 120.4, 116.6, 111.1, 110.7, 103.2, 33.5. IR (KBr): 3461, 2979, 1633, 1523, 1466, 1400, 1259, 1065, 742 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{20}\text{NO}$ $[\text{M}+\text{H}]^+$ 398.1545, found 398.1537.




(6aR,14R)-14-(3,5-dimethylphenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2t)

White solid (57.8 mg, 75% yield). PE / EA = 10:1, R_f = 0.33. ^1H NMR (400 MHz, CDCl_3): δ 8.47 (d, J = 8.6 Hz, 2H), 7.81 (d, J = 8.0, 1.2 Hz, 2H), 7.73 (d, J = 8.6 Hz, 2H), 7.60 (t, J = 8.2 Hz, 2H), 7.38 (t, J = 8.2 Hz, 2H), 7.19–7.14 (m, 4H), 6.67 (d, J = 8.2 Hz, 2H), 6.49 (s, 1H), 2.19 (s, 6H). ^{13}C NMR (100 MHz, CDCl_3): δ 146.1, 137.5, 135.9, 132.2, 130.3, 128.7, 128.2, 128.0, 126.7, 125.7, 122.9, 122.2, 116.8, 114.9, 39.1, 21.5. IR (KBr): 3461, 3015, 2989, 1642, 1543, 1519, 1451, 1384, 1266, 998 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{24}\text{N}$ $[\text{M}+\text{H}]^+$ 386.1909, found 386.1903.




(6aR,14R)-14-(3,5-dimethoxyphenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2u)

Yellow solid (64.2 mg, 77% yield). PE / EA = 10:1, R_f = 0.29. ^1H NMR (400 MHz, CDCl_3): δ 8.49 (d, J = 8.5 Hz, 2H), 7.81 (d, J = 8.0 Hz, 2H), 7.70–7.57 (m, 4H), 7.40 (t, J = 7.4 Hz, 2H), 6.99 (d, J = 8.7 Hz, 2H), 6.79 (s, 2H), 6.73 (s, 1H), 6.50 (s, 1H), 6.19 (s, 1H), 3.67 (s, 6H). ^{13}C NMR (100 MHz, CDCl_3): δ 160.6, 148.4, 136.0, 132.2, 130.3, 128.8, 128.3, 126.8, 123.0, 122.2, 116.8, 114.2, 106.9, 97.2, 55.1, 39.2. IR (KBr): 3455, 3010, 2987, 1661, 1579, 1460, 1429, 1278, 1219, 1088 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{24}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 418.1807, found 418.1801.




(6aR,14R)-14-(3,4-dimethylphenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2v)

White solid (54.6 mg, 71% yield). PE / EA = 10:1, R_f = 0.31. ^1H NMR (400 MHz, CDCl_3): δ 8.56–8.44 (m, 2H), 7.84 (t, J = 7.7 Hz, 2H), 7.77 (t, J = 7.6 Hz, 2H), 7.70–7.57 (m, 2H), 7.46–7.36 (m, 2H), 7.36–7.26 (m, 2H), 7.25–7.13 (m, 2H), 7.01–6.90 (m, 1H), 6.73 (s, 1H), 6.51 (s, 1H), 2.21–2.06 (m, 6H). ^{13}C NMR (100 MHz, CDCl_3): δ 143.6, 136.3, 135.9, 134.3, 132.2, 130.3, 129.4, 129.1, 128.7, 128.1, 126.8, 125.2, 122.9, 122.2, 116.7, 115.0, 38.7, 19.9, 19.1. IR (KBr): 3459, 3008, 2989, 1645, 1533, 1520, 1454, 1382, 1280, 1048 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{24}\text{N}$ $[\text{M}+\text{H}]^+$ 386.1909, found 386.1901.




(6aR,14R)-14-(3,4-dimethoxyphenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2w)

White solid (65.1 mg, 78% yield). PE / EA = 10:1, R_f = 0.27. ^1H NMR (400 MHz, CDCl_3): δ 8.48 (d, J = 8.6 Hz, 2H), 7.84 (d, J = 8.1 Hz, 2H), 7.72 (d, J = 8.7 Hz, 2H), 7.62 (t, J = 7.8 Hz, 2H), 7.40 (t, J = 7.5 Hz, 2H), 7.13 (d, J = 8.7 Hz, 2H), 7.10–6.99 (m, 2H), 6.75 (s, 1H), 6.66 (d, J = 8.1 Hz, 1H), 6.59 (s, 1H), 3.72 (s, 3H), 3.70 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 149.0, 147.4, 138.8, 136.1, 132.2, 130.3, 128.8, 128.2, 126.8, 123.0, 122.1, 119.8, 116.7, 114.7, 111.8, 111.0, 55.7, 55.7, 38.3. IR (KBr): 3445, 3023, 2966, 1654, 1545, 1532, 1473, 1424, 1257, 1206, 1169 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{24}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 418.1807, found 418.1802.




(6aR,14S)-14-(2-bromo-4-methoxyphenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2x)

White solid (50.2 mg, 54% yield). PE / EA = 10:1, R_f = 0.27. ^1H NMR (400 MHz, DMSO): δ 9.70 (s, 1H), 8.76 (d, J = 8.6 Hz, 2H), 7.81 (t, J = 9.0 Hz, 4H), 7.62–7.48 (m, 3H), 7.42 (d, J = 8.7 Hz, 2H), 7.32 (t, J = 7.4 Hz, 2H), 6.96 (s, 1H), 6.72–6.59 (m, 2H), 3.54 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 158.5, 140.6, 137.0, 132.4, 132.3, 129.9, 129.1, 128.7, 126.9, 123.1, 123.0, 119.3, 118.0, 117.1, 116.2, 114.4, 55.7, 37.8. IR (KBr): 3458, 3014, 2987, 1671, 1568, 1531, 1466, 1418, 1281, 1234, 1045, 949, 641 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{28}\text{H}_{21}\text{BrNO}$ $[\text{M}+\text{H}]^+$ 486.0807, found 486.0802.




(6aR,14R)-14-(4-bromo-3-fluorophenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2y)

White solid (56.2 mg, 62% yield). PE / EA = 7:1, R_f = 0.29. ^1H NMR (400 MHz, DMSO): δ 9.68 (s, 1H), 8.62 (d, J = 8.5 Hz, 2H), 7.85–7.78 (m, 4H), 7.69 (d, J = 10.1 Hz, 1H), 7.58 (t, J = 7.7 Hz, 2H), 7.46–7.30 (m, 6H), 6.82 (s, 1H). ^{13}C NMR (100 MHz, DMSO): δ 158.4 (d, J = 245.1 Hz), 150.0 (d, J = 5.3 Hz), 136.8, 133.7, 132.0, 129.9, 129.0, 128.9, 127.3, 125.7 (d, J = 3.0 Hz), 123.2, 122.5, 117.7, 115.8 (d, J = 21.9 Hz), 113.0, 105.6 (d, J = 20.7 Hz), 38.1. ^{19}F NMR (376 MHz, DMSO): δ -107.3. IR (KBr): 3471, 3007, 1626, 1541, 1511, 1474, 1445, 1333, 1261, 1234, 652 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{27}\text{H}_{18}\text{BrFN}$ $[\text{M}+\text{H}]^+$ 454.0607, found 454.0602.




(6aR,14S)-14-(2-bromo-4-methylphenyl)-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2z)

Yellow solid (49.4 mg, 55% yield). PE / EA = 10:1, R_f = 0.29. ^1H NMR (400 MHz, CDCl_3): δ 8.95 (d, J = 8.6 Hz, 2H), 7.78 (d, J = 8.0 Hz, 2H), 7.71 (d, J = 8.7 Hz, 2H), 7.67–7.58 (m, 3H), 7.38 (t, J = 7.5 Hz, 2H), 7.23 (s, 1H), 7.12 (d, J = 8.6 Hz, 2H), 6.86 (s, 1H), 6.76 (d, J = 8.2 Hz, 1H), 6.51 (s, 1H), 2.06 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 144.8, 137.9, 135.8, 133.1, 132.4, 131.5, 130.2, 129.6, 128.5, 128.5, 126.7, 123.7, 123.1, 119.4, 116.9, 115.9, 38.4, 20.2. IR (KBr): 3479, 3018, 2972, 1664, 1528, 1466, 1422, 1387, 1228, 1014, 677 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{28}\text{H}_{21}\text{BrN}$ $[\text{M}+\text{H}]^+$ 450.0857, found 450.0849.




(6aR,14R)-2,12-dimethoxy-14-phenyl-6a,7,14,14a-tetrahydrodibenzo[a,j]acridine (2a')

White solid (50.8 mg, 61% yield). PE / EA = 7:1, R_f = 0.28. ^1H NMR (400 MHz, CDCl_3): δ 7.76–7.69 (m, 4H), 7.69–7.64 (m, 2H), 7.60 (d, J = 7.5 Hz, 2H), 7.20 (t, J = 7.5 Hz, 2H), 7.09–7.03 (m, 3H), 7.04–6.98 (m, 2H), 6.50 (s, 1H), 6.44 (s, 1H), 4.09 (s, 6H). ^{13}C NMR (100 MHz, CDCl_3): δ 158.4, 146.1, 136.1, 133.4, 130.2, 128.3, 128.1, 128.0, 126.1, 125.5, 114.3, 114.1, 113.8, 102.5, 55.4, 40.3. IR (KBr): 3455, 3009, 2986, 1654, 1545, 1473, 1424, 1257, 1206, 1109 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{24}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 418.1807, found 418.1799.




(6aR,14R)-2,12-dibromo-14-phenyl-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2b')

White solid (68.6 mg, 67% yield). PE / EA = 8:1, R_f = 0.31. ^1H NMR (400 MHz, DMSO): δ 9.82 (s, 1H), 8.90 (s, 2H), 7.79 (d, J = 8.8 Hz, 2H), 7.75 (d, J = 8.6 Hz, 2H), 7.59 (d, J = 7.1 Hz, 2H), 7.45–7.35 (m, 4H), 7.15 (t, J = 7.6 Hz, 2H), 6.98 (t, J = 7.4 Hz, 1H), 6.73 (s, 1H). ^{13}C NMR (100 MHz, DMSO): δ 147.6, 137.5, 133.7, 131.1, 128.8, 128.6, 127.9, 126.6, 126.1, 125.3, 121.2, 118.2, 113.5, 38.4. IR (KBr): 3465, 3010, 1679, 1519, 1485, 1453, 1276, 1227, 689 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{27}\text{H}_{18}\text{Br}_2\text{N}$ $[\text{M}+\text{H}]^+$ 513.9806, found 513.9801.




(6aR,14R)-3,11-dimethyl-14-phenyl-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2c')

White solid (44.5 mg, 58% yield). PE / EA = 8:1, R_f = 0.31. ^1H NMR (400 MHz, CDCl_3): δ 8.34 (d, J = 8.6 Hz, 2H), 7.66 (d, J = 8.6 Hz, 2H), 7.59 (s, 2H), 7.50 (d, J = 7.7 Hz, 2H), 7.43 (d, J = 8.7 Hz, 2H), 7.17–7.10 (m, 4H), 7.01 (t, J = 7.2 Hz, 1H), 6.70 (s, 1H), 6.44 (s, 1H), 2.52 (s, 6H). ^{13}C NMR (100 MHz, CDCl_3): δ 146.2, 135.4, 132.3, 130.4, 130.3, 128.9, 128.2, 127.9, 127.8, 127.6, 126.0, 122.0, 116.8, 114.6, 39.1, 21.3. IR (KBr): 3463, 3011, 2991, 1625, 1568, 1525, 1431, 1377, 1278, 1225, 994 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{29}\text{H}_{24}\text{N}$ $[\text{M}+\text{H}]^+$ 386.1909, found 386.1901.




(6aR,14R)-3,11-dicyclopropyl-14-phenyl-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2d')

White solid (68.2 mg, 78% yield). PE / EA = 10:1, R_f = 0.32. ^1H NMR (400 MHz, CDCl_3): δ 8.30 (d, J = 8.7 Hz, 2H), 7.62 (d, J = 8.7 Hz, 2H), 7.52–7.44 (m, 4H), 7.29 (s, 2H), 7.16–7.07 (m, 4H), 6.98 (t, J = 7.4 Hz, 1H), 6.64 (s, 1H), 6.44 (s, 1H), 2.11–1.98 (m, 2H), 1.08–0.98 (m, 4H), 0.85–0.74 (m, 4H). ^{13}C NMR (100 MHz, CDCl_3): δ 146.2, 138.2, 135.3, 130.5, 130.3, 128.2, 127.8, 127.6, 126.0, 125.6, 124.8, 122.2, 116.9, 114.6, 39.2, 15.3, 9.0, 8.8. IR (KBr): 3545, 3401, 3003, 2922, 1599, 1520, 1485, 1401, 943 804, 749, 510 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{33}\text{H}_{28}\text{N}$ $[\text{M}+\text{H}]^+$ 438.2222, found 438.2218.




(6aR,14R)-3,11-dibromo-14-phenyl-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2e')

White solid (69.6 mg, 68% yield). PE / EA = 8:1, R_f = 0.28. ^1H NMR (400 MHz, DMSO): δ 9.74 (s, 1H), 8.52 (d, J = 9.1 Hz, 2H), 8.04 (s, 2H), 7.75 (d, J = 8.7 Hz, 2H), 7.59 (d, J = 9.1 Hz, 2H), 7.53 (d, J = 7.7 Hz, 2H), 7.38 (d, J = 8.8 Hz, 2H), 7.09 (t, J = 7.7 Hz, 2H), 6.95 (t, J = 7.5 Hz, 1H), 6.67 (s, 1H). ^{13}C NMR (100 MHz, DMSO): δ 152.1, 141.7, 136.0, 135.5, 135.4, 134.4, 133.4, 132.6, 132.5, 131.2, 123.0, 123.6, 120.6, 118.8, 43.2. IR (KBr): 3462, 3032, 1618, 1555, 1528, 1461, 1421, 1266, 1212, 677 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{27}\text{H}_{18}\text{Br}_2\text{N}$ $[\text{M}+\text{H}]^+$ 513.9806, found 513.9798.




(6aR,14R)-3,11,14-triphenyl-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2f')

Yellow solid (60.9 mg, 60% yield). PE / EA = 8:1, R_f = 0.31. ^1H NMR (400 MHz, CDCl_3): δ 8.52 (d, J = 8.8 Hz, 2H), 8.02 (d, J = 2.0 Hz, 2H), 7.88 (d, J = 8.6 Hz, 2H), 7.80 (d, J = 8.7 Hz, 2H), 7.76 (d, J = 9.6 Hz, 4H), 7.57 (d, 2H), 7.52 (t, J = 7.7 Hz, 4H), 7.44–7.37 (m, 2H), 7.22–7.15 (m, 4H), 7.04 (t, J = 6.9 Hz, 1H), 6.79 (s, 1H), 6.56 (s, 1H). ^{13}C NMR (100 MHz, CDCl_3): δ 146.0, 141.1, 135.9, 135.7, 131.4, 130.6, 128.8, 128.7, 128.4, 127.8, 127.1, 127.0, 126.6, 126.4, 126.2, 122.7, 117.1, 114.6, 39.3. IR (KBr): 3451, 3019, 1638, 1559, 1541, 1477, 1438, 1409, 1259, 1223 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{39}\text{H}_{28}\text{N}$ $[\text{M}+\text{H}]^+$ 510.2222, found 510.2218.




(6aR,14R)-14-phenyl-3,11-di(thiophen-3-yl)-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2g')

Yellow solid (84.4 mg, 83% yield). PE / EA = 10:1, R_f = 0.29. ^1H NMR (400 MHz, DMSO): δ 9.81 (s, 1H), 8.72 (d, J = 9.0 Hz, 2H), 8.26 (s, 2H), 8.07–7.98 (m, 4H), 7.93 (d, J = 8.8 Hz, 2H), 7.80–7.66 (m, 6H), 7.53 (d, J = 8.8, 1.9 Hz, 2H), 7.17 (t, J = 7.7 Hz, 2H), 7.00 (t, J = 7.4 Hz, 1H), 6.87 (s, 1H). ^{13}C NMR (100 MHz, DMSO): δ 148.0, 141.9, 136.7, 131.4, 130.3, 129.8, 128.8, 128.6, 128.1, 127.5, 126.7, 126.4, 125.8, 125.6, 123.6, 120.9, 118.2, 114.2, 39.0. IR (KBr): 3467, 3031, 2955, 1704, 1561, 1518, 1444, 1408, 1272, 1230, 1015, 655 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{35}\text{H}_{24}\text{NS}_2$ $[\text{M}+\text{H}]^+$ 522.1350, found 522.1342.




(7aR,17R)-17-phenyl-7a,8,17,17a-tetrahydronaphtho[2,3-a:2',3'-j]acridine (2h')

Yellow solid (70.4 mg, 77% yield). PE / EA = 7:1, R_f = 0.27. ^1H NMR (400 MHz, DMSO): δ 9.79 (s, 1H), 9.17 (s, 2H), 8.43 (s, 2H), 8.19 (d, J = 8.5 Hz, 2H), 7.97 (t, J = 9.6 Hz, 4H), 7.80 (d, J = 7.7 Hz, 2H), 7.52 (t, J = 7.6 Hz, 2H), 7.46–7.38 (m, 4H), 7.08 (t, J = 7.6 Hz, 2H), 6.98 (s, 1H), 6.87 (t, J = 7.4 Hz, 1H). ^{13}C NMR (100 MHz, DMSO): δ 148.1, 135.5, 132.4, 130.7, 129.6, 129.4, 129.0, 128.5, 128.3, 128.3, 127.3, 126.2, 126.2, 124.8, 120.3, 119.4, 111.7, 39.1. IR (KBr): 3451, 3031, 1644, 1558, 1511, 1470, 1438, 1280, 1231 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{35}\text{H}_{24}\text{N}$ $[\text{M}+\text{H}]^+$ 458.1909, found 458.1903.




(6aR,14R)-3,11-dibromo-14-(4-(trifluoromethyl)phenyl)-6a,7,14,14a-tetrahydrodibenzo[*a,j*]acridine (2i')

Yellow solid (98.6 mg, 85% yield). PE / EA = 7:1, R_f = 0.31. ^1H NMR (400 MHz, DMSO): δ 9.62 (s, 1H), 8.25 (d, J = 9.2 Hz, 2H), 7.74 (s, 2H), 7.55–7.44 (m, 4H), 7.35 (d, J = 9.2 Hz, 2H), 7.21 (d, J = 8.8 Hz, 2H), 7.10 (d, J = 8.0 Hz, 2H), 6.55 (s, 1H). ^{13}C NMR (100 MHz, DMSO): δ 151.4, 137.1, 131.3, 130.7, 130.7, 129.9, 128.4, 128.2, 127.3 (q, J = 31.7 Hz), 125.6 (q, J = 4.5 Hz), 124.9, 124.5 (q, J = 270.3 Hz), 119.0, 116.1, 113.2, 38.6. ^{19}F NMR (376 MHz, DMSO): δ -61.1. IR (KBr): 3463, 3023, 1631, 1518, 1477, 1422, 1324, 1285, 1206 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{28}\text{H}_{17}\text{Br}_2\text{FN}$ $[\text{M}+\text{H}]^+$ 581.9680, found 581.9672.




7,14-diphenyldibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8a)

Yellow solid (81.8 mg, 79% yield). DCM / MeOH = 100:1, R_f = 0.24. ^1H NMR (400 MHz, CDCl_3): δ 8.26 (d, J = 9.3 Hz, 2H), 8.00 (d, J = 7.8 Hz, 2H), 7.94–7.82 (m, 4H), 7.79–7.70 (m, 4H), 7.68–7.57 (m, 4H), 7.39–7.22 (m, 6H). ^{13}C NMR (100 MHz, CDCl_3): δ 156.2, 142.7, 140.7, 140.4, 138.3, 132.4, 132.0, 131.7, 131.5, 130.7, 129.9, 129.4, 129.3, 129.1, 128.9, 128.3, 128.0, 125.0, 117.1. ^{19}F NMR (376 MHz, DMSO): δ -154.2. IR (KBr): 3465, 3048, 1586, 1461, 1345, 1221, 1084 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{33}\text{H}_{22}\text{N}^+$ $[\text{M}]^+$ 432.1747 found 432.1744.




7-phenyl-14-(p-tolyl)dibenzo[a,j]acridin-7-ium tetrafluoroborate (8b)

Red solid (76.6 mg, 72% yield). DCM / MeOH = 100:1, R_f = 0.25. ^1H NMR (400 MHz, DMSO): δ 8.56 (d, J = 9.4 Hz, 2H), 8.24 (d, J = 7.8 Hz, 2H), 8.05–7.98 (m, 3H), 7.94–7.88 (m, 2H), 7.78 (t, J = 7.9 Hz, 2H), 7.70 (d, J = 7.8 Hz, 2H), 7.49 (d, J = 7.8 Hz, 2H), 7.45–7.30 (m, 6H), 2.68 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 155.9, 143.0, 141.3, 140.5, 139.0, 138.1, 132.6, 132.3, 132.0, 131.9, 130.5, 129.4, 129.3, 129.2, 128.8, 128.8, 128.3, 124.8, 117.9, 21.8. ^{19}F NMR (376 MHz, CDCl_3): δ -154.4. IR (KBr): 3661, 2981, 1686, 1531, 1371, 1242, 1072, 752 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{34}\text{H}_{24}\text{N}^+$ $[\text{M}]^+$ 446.1903 found 446.1898.



14-(4-(tert-butyl)phenyl)-7-phenyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8c)


Yellow solid (82.9 mg, 72% yield). DCM / MeOH = 100:1, R_f = 0.27. ^1H NMR (400 MHz, CDCl_3): δ 8.28 (d, J = 9.4 Hz, 2H), 8.02 (d, J = 7.9 Hz, 2H), 7.94–7.89 (m, 3H), 7.80 (d, J = 8.1 Hz, 2H), 7.71–7.63 (m, 4H), 7.51 (d, J = 8.1 Hz, 2H), 7.36 (t, J = 9.1 Hz, 4H), 7.24 (t, J = 8.2 Hz, 2H), 1.59 (s, 9H). ^{13}C NMR (100 MHz, CDCl_3): δ 156.6, 155.0, 142.6, 140.6, 138.3, 137.9, 132.5, 132.1, 131.8, 130.0, 129.3, 129.3, 129.0, 128.9, 128.5, 128.3, 127.9, 125.1, 117.1, 35.3, 31.6. ^{19}F NMR (376 MHz, CDCl_3): δ -154.3. IR (KBr): 3463, 2959, 1634, 1520, 1371, 1213, 1055, 827, 758 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{37}\text{H}_{30}\text{N}^+$ $[\text{M}]^+$ 488.2373 found 488.2369.



14-(4-methoxyphenyl)-7-phenyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8d)


Yellow solid (64.8 mg, 59% yield). DCM / MeOH = 100:1, R_f = 0.23. ^1H NMR (400 MHz, CDCl_3): δ 8.23 (t, J = 8.0 Hz, 2H), 8.06–7.81 (m, 7H), 7.65 (d, J = 7.4 Hz, 2H), 7.53–7.24 (m, 10H), 4.09 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 161.8, 156.3, 142.6, 140.2, 138.3, 132.6, 132.4, 131.9, 131.6, 130.9, 129.8, 129.3, 129.0, 128.9, 128.5, 128.0, 125.2, 117.2, 117.1, 55.7. ^{19}F

NMR (376 MHz, CDCl₃): δ -154.3. IR (KBr): 3671, 2897, 1669, 1534, 1361, 1254, 1206, 1102, 742 cm⁻¹. HRMS (ESI) m/z Calcd for C₃₄H₂₄NO⁺ [M]⁺ 462.1852 found 462.1843.




14-(4-(methylthio)phenyl)-7-phenyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8e)

Yellow solid (76.8 mg, 68% yield). DCM / MeOH = 100:1, R_f = 0.28. ¹H NMR (400 MHz, DMSO): δ 8.53 (d, J = 9.7 Hz, 2H), 8.23 (d, J = 7.5 Hz, 2H), 8.02–7.67 (m, 9H), 7.50 (d, J = 7.3 Hz, 2H), 7.46–7.37 (m, 4H), 7.33 (d, J = 9.3 Hz, 2H), 2.70 (s, 3H). ¹³C NMR (100 MHz, DMSO): δ 155.1, 143.0, 142.9, 140.5, 138.9, 136.9, 132.6, 132.3, 131.9, 130.5, 130.1, 129.4, 129.1, 128.8, 128.3, 124.8, 117.8, 15.1. ¹⁹F NMR (376 MHz, CDCl₃): δ -154.4. IR (KBr): 3471, 3056, 1750, 1590, 1519, 1371, 1052, 821, 755 cm⁻¹. HRMS (ESI) m/z Calcd for C₃₄H₂₄NS⁺ [M]⁺ 478.1624 found 478.1605.



14-(4-nitrophenyl)-7-phenyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8f)


Yellow solid (64.3 mg, 57% yield). DCM / MeOH = 100:1, R_f = 0.23. ¹H NMR (400 MHz, DMSO): δ 8.72 (d, J = 8.2 Hz, 2H), 8.61 (d, J = 9.4 Hz, 2H), 8.29 (d, J = 7.8 Hz, 2H), 8.10–8.02 (m, 3H), 7.98 (d, J = 8.2 Hz, 2H), 7.95–7.88 (m, 2H), 7.82 (t, J = 7.5 Hz, 2H), 7.48 (t, J = 8.0 Hz, 2H), 7.41 (d, J = 9.4 Hz, 2H), 7.23 (d, J = 8.7 Hz, 2H). ¹³C NMR (100 MHz, DMSO): δ 152.9, 149.3, 147.2, 143.1, 140.8, 138.8, 132.7, 132.4, 132.0, 131.7, 130.7, 129.7, 129.2, 128.7, 128.5, 127.0, 124.4, 117.9. ¹⁹F NMR (376 MHz, CDCl₃): δ -154.2. IR (KBr): 3688, 3049, 1636, 1514, 1371, 1216, 751 cm⁻¹. HRMS (ESI) m/z Calcd for C₃₃H₂₁N₂O₂⁺ [M]⁺ 477.1598 found 477.1589.



14-(4-cyanophenyl)-7-phenyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8g)


Yellow solid (69.6 mg, 64% yield). DCM / MeOH = 100:1, R_f = 0.27. ¹H NMR (400 MHz, CDCl₃): δ 8.21 (d, J = 9.4 Hz, 2H), 7.99 (d, J = 7.8 Hz, 4H), 7.92–7.83 (m, 5H), 7.74 (d, J = 6.2 Hz, 2H), 7.65 (t, J = 7.4 Hz, 2H), 7.33–7.25 (m, 4H), 7.21 (d, J = 8.7 Hz, 2H). ¹³C NMR (100

MHz, CDCl₃): δ 153.2, 145.4, 142.9, 140.2, 138.3, 134.6, 132.6, 131.9, 131.5, 131.2, 130.1, 129.3, 129.0, 128.6, 128.4, 127.9, 124.7, 118.1, 117.2, 114.2. ¹⁹F NMR (376 MHz, CDCl₃): δ -154.3. IR (KBr): 3918, 3068, 1733, 1521, 1371, 1052, 758 cm⁻¹. HRMS (ESI) m/z Calcd for C₃₄H₂₁N₂⁺ [M]⁺ 457.1699 found 457.1694.




14-(3,4-dimethylphenyl)-7-phenyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8h)

Yellow solid (68.9 mg, 63% yield). DCM / MeOH = 100:1, R_f = 0.28. ¹H NMR (400 MHz, CDCl₃): δ 8.30 (d, *J* = 9.3 Hz, 2H), 8.04 (d, *J* = 7.9 Hz, 2H), 7.94 (d, *J* = 6.1 Hz, 3H), 7.74–7.65 (m, 4H), 7.55 (d, *J* = 7.7 Hz, 1H), 7.45 (d, *J* = 8.7 Hz, 2H), 7.38–7.27 (m, 6H), 2.62 (s, 3H), 2.41 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 156.7, 142.5, 140.5, 139.8, 138.2, 138.0, 132.7, 132.4, 132.0, 131.8, 129.9, 129.3, 129.2, 129.0, 128.2, 127.9, 126.4, 125.0, 117.0, 20.0, 19.9. ¹⁹F NMR (376 MHz, DMSO): δ -154.4. IR (KBr): 3669, 3050, 2991, 1673, 1562, 1370, 1276, 1033, 759 cm⁻¹. HRMS (ESI) m/z Calcd for C₃₅H₂₆N⁺ [M]⁺ 460.2060 found 460.2055.



14-(3,4-dimethoxyphenyl)-7-phenyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8i)


Yellow solid (68.4 mg, 59% yield). DCM / MeOH = 100:1, R_f = 0.23. ¹H NMR (400 MHz, DMSO): δ 8.57 (d, *J* = 9.6 Hz, 2H), 8.26 (d, *J* = 7.9 Hz, 2H), 8.07–7.77 (m, 7H), 7.53–7.42 (m, 5H), 7.37 (d, *J* = 9.5 Hz, 2H), 7.19–7.08 (m, 2H), 4.07 (s, 3H), 3.63 (s, 3H). ¹³C NMR (100 MHz, DMSO): δ 155.8, 152.1, 151.6, 142.9, 140.5, 138.9, 132.9, 132.5, 132.3, 132.0, 130.4, 129.4, 129.3, 128.8, 128.3, 125.0, 122.1, 117.7, 115.1, 113.1, 56.6. ¹⁹F NMR (376 MHz, CDCl₃): δ -154.3. IR (KBr): 3677, 3049, 2889, 1724, 1643, 1551, 1326, 1224, 1022, 741 cm⁻¹. HRMS (ESI) m/z Calcd for C₃₅H₂₆NO₂⁺ [M]⁺ 492.1958 found 492.1952.



14-(3,5-dimethylphenyl)-7-phenyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8j)


Yellow solid (83.2 mg, 76% yield). DCM / MeOH = 100:1, R_f = 0.24. ¹H NMR (400 MHz, CDCl₃): δ 8.27 (d, *J* = 9.3 Hz, 2H), 8.01 (d, *J* = 7.8 Hz, 2H), 7.94–7.86 (m, 3H), 7.71–7.61 (m,

4H), 7.46 (s, 1H), 7.42 (d, $J = 8.7$ Hz, 2H), 7.36–7.26 (m, 4H), 7.16 (s, 2H), 2.43 (s, 6H). ^{13}C NMR (100 MHz, CDCl_3): δ 156.8, 142.5, 141.7, 140.6, 140.4, 138.2, 132.4, 132.3, 132.0, 131.8, 130.0, 129.4, 129.2, 129.1, 128.2, 128.0, 126.4, 124.8, 117.0, 21.5. ^{19}F NMR (376 MHz, CDCl_3): δ -154.2. IR (KBr): 3647, 2977, 1633, 1551, 1370, 1249, 1208, 805 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{35}\text{H}_{26}\text{N}^+ [\text{M}]^+$ 460.2060 found 460.2049.




14-(3,5-dimethoxyphenyl)-7-phenyldibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8k)

Yellow solid (71.6 mg, 69% yield). DCM / MeOH = 100:1, $R_f = 0.25$. ^1H NMR (400 MHz, CDCl_3): δ 8.28 (d, $J = 9.5$ Hz, 2H), 8.03 (d, $J = 7.8$ Hz, 2H), 7.97–7.87 (m, 3H), 7.74–7.64 (m, 6H), 7.46–7.33 (m, 4H), 6.92 (s, 1H), 6.80 (d, $J = 2.2$ Hz, 2H), 3.82 (s, 6H). ^{13}C NMR (100 MHz, CDCl_3): δ 163.7, 156.1, 142.6, 142.1, 140.4, 138.4, 132.3, 131.9, 131.7, 129.9, 129.3, 129.1, 129.0, 128.3, 128.1, 124.8, 117.0, 106.9, 103.0, 56.0. ^{19}F NMR (376 MHz, CDCl_3): δ -154.5. IR (KBr): 3675, 3061, 2878, 1641, 1553, 1338, 1239, 1055, 776 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{35}\text{H}_{26}\text{NO}_2^+ [\text{M}]^+$ 492.1958 found 492.1951.




14-(2-bromo-4-methylphenyl)-7-phenyldibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8l)

Yellow solid (68.3 mg, 56% yield). DCM / MeOH = 100:1, $R_f = 0.27$. ^1H NMR (400 MHz, CDCl_3): δ 8.27 (d, $J = 9.3$ Hz, 2H), 8.03 (d, $J = 7.9$ Hz, 2H), 7.95–7.86 (m, 4H), 7.79 (s, 1H), 7.70 (t, $J = 7.4$ Hz, 2H), 7.62–7.52 (m, 3H), 7.43 (d, $J = 8.7$ Hz, 2H), 7.36 (t, $J = 8.1$ Hz, 4H), 2.68 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 154.1, 143.3, 142.7, 140.5, 138.3, 138.2, 135.4, 132.3, 132.1, 132.0, 131.7, 131.6, 131.3, 130.2, 129.1, 129.0, 128.7, 128.0, 127.6, 125.0, 122.0, 117.2, 21.4. ^{19}F NMR (376 MHz, CDCl_3): δ -154.4. IR (KBr): 3654, 3031, 2978, 1664, 1527, 1370, 1221, 648 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{34}\text{H}_{23}\text{BrN}^+ [\text{M}]^+$ 524.1008 found 524.1003.




14-(2-bromo-4-methoxyphenyl)-7-phenyldibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8m)

Yellow solid (52.6 mg, 42% yield). DCM / MeOH = 100:1, R_f = 0.24. ^1H NMR (400 MHz, CDCl_3): δ 8.27 (d, J = 9.2 Hz, 2H), 8.03 (d, J = 7.9 Hz, 2H), 7.94–7.88 (m, 4H), 7.71 (t, J = 7.4 Hz, 2H), 7.59 (d, J = 7.8 Hz, 2H), 7.51–7.40 (m, 6H), 7.32–7.24 (m, 2H), 4.07 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 162.2, 154.0, 142.6, 140.5, 138.3, 133.2, 132.7, 132.2, 132.1, 132.0, 131.3, 130.1, 129.3, 129.1, 128.9, 128.7, 127.9, 127.6, 125.3, 122.8, 120.6, 117.1, 116.7, 56.1. ^{19}F NMR (376 MHz, CDCl_3): δ -154.3. IR (KBr): 3663, 3048, 2878, 1674, 1537, 1280, 1209, 1064 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{34}\text{H}_{23}\text{BrNO}^+ [\text{M}]^+$ 540.0958 found 540.0951.




14-(naphthalen-2-yl)-7-phenyldibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8n)

Yellow solid (49.8 mg, 44% yield). DCM / MeOH = 100:1, R_f = 0.28. ^1H NMR (400 MHz, CDCl_3): δ 8.28 (d, J = 8.7 Hz, 3H), 8.15 (d, J = 8.3 Hz, 1H), 8.10 (s, 1H), 8.00 (d, J = 7.9 Hz, 2H), 7.96–7.90 (m, 3H), 7.87 (d, J = 8.2 Hz, 1H), 7.80–7.64 (m, 5H), 7.57 (t, J = 7.5 Hz, 2H), 7.37 (d, J = 9.3 Hz, 2H), 7.29 (d, J = 8.8 Hz, 2H), 7.05 (t, J = 7.9 Hz, 2H). ^{13}C NMR (100 MHz, CDCl_3): δ 156.0, 142.7, 140.4, 138.3, 137.8, 134.5, 133.8, 132.5, 132.0, 131.7, 131.5, 129.9, 129.3, 129.2, 129.2, 128.9, 128.3, 128.1, 127.9, 127.5, 126.3, 125.2, 117.1. ^{19}F NMR (376 MHz, CDCl_3): δ -154.2. IR (KBr): 3662, 3048, 1666, 1540, 1347, 1261, 1206 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{37}\text{H}_{24}\text{N}^+ [\text{M}]^+$ 482.1904 found 482.1898.




3,7,11,14-tetraphenyldibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8o)

Yellow solid (95.1 mg, 71% yield). DCM / MeOH = 100:1, R_f = 0.28. ^1H NMR (400 MHz, DMSO): δ 8.67–8.56 (m, 4H), 8.07–7.92 (m, 8H), 7.88 (d, J = 7.6 Hz, 4H), 7.75–7.64 (m, 4H), 7.55 (t, J = 7.5 Hz, 4H), 7.51–7.39 (m, 4H), 7.36 (d, J = 9.1 Hz, 2H). ^{13}C NMR (100 MHz, DMSO): δ 155.4, 142.9, 140.8, 140.8, 140.4, 139.0, 138.3, 133.3, 132.4, 132.2, 132.0, 131.5, 129.6, 129.5, 129.0, 128.7, 128.1, 127.8, 127.5, 126.4, 124.8, 118.3. ^{19}F NMR (376 MHz, CDCl_3): δ -154.3. IR (KBr): 3665, 3049, 1581, 1473, 1331, 1251, 1043 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{45}\text{H}_{30}\text{N}^+ [\text{M}]^+$ 584.2373 found 584.2367.




7,14-di-p-tolyldibenzo[a,j]acridin-7-ium tetrafluoroborate (8p)

Yellow solid (81.1 mg, 74% yield). DCM / MeOH = 100:1, R_f = 0.29. ^1H NMR (400 MHz, DMSO): δ 8.56 (d, J = 9.4 Hz, 2H), 8.24 (d, J = 7.8 Hz, 2H), 7.85–7.74 (m, 6H), 7.70 (d, J = 7.7 Hz, 2H), 7.49 (d, J = 7.7 Hz, 2H), 7.44–7.35 (m, 6H), 2.68 (s, 6H). ^{13}C NMR (100 MHz, DMSO): δ 155.8, 143.1, 142.3, 141.2, 140.4, 138.1, 136.5, 132.6, 132.6, 132.3, 130.5, 129.4, 129.2, 128.8, 128.5, 128.2, 124.8, 117.9, 21.8, 21.5. ^{19}F NMR (376 MHz, CDCl_3): δ -154.3. IR (KBr): 3673, 3051, 2959, 1737, 1541, 1373, 1241, 752 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{35}\text{H}_{26}\text{N}^+$ $[\text{M}]^+$ 460.2060 found 460.2053.




7-(4-chlorophenyl)-14-(p-tolyl)dibenzo[a,j]acridin-7-ium tetrafluoroborate (8q)

Yellow solid (75.8 mg, 67% yield). DCM / MeOH = 100:1, R_f = 0.27. ^1H NMR (400 MHz, DMSO): δ 8.57 (d, J = 9.4 Hz, 2H), 8.27 (d, J = 7.9 Hz, 2H), 8.10 (d, J = 8.3 Hz, 2H), 7.96 (d, J = 8.4 Hz, 2H), 7.78 (t, J = 7.5 Hz, 2H), 7.70 (d, J = 7.7 Hz, 2H), 7.5–7.34 (m, 8H), 2.68 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 156.0, 143.0, 141.3, 140.6, 138.0, 137.7, 137.1, 132.6, 132.1, 130.8, 130.5, 129.4, 129.4, 129.2, 128.8, 128.3, 124.8, 117.9, 21.8. ^{19}F NMR (376 MHz, CDCl_3): δ -154.2. IR (KBr): 3917, 3060, 1734, 1520, 1370, 1051, 824, 757 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{34}\text{H}_{23}\text{ClN}^+$ $[\text{M}]^+$ 480.1514 found 480.1505.




7-(4-(methoxycarbonyl)phenyl)-14-(p-tolyl)dibenzo[a,j]acridin-7-ium tetrafluoroborate (8r)

Yellow solid (68.6 mg, 58% yield). DCM / MeOH = 100:1, R_f = 0.27. ^1H NMR (400 MHz, DMSO): δ 8.53 (d, J = 8.8 Hz, 4H), 8.24 (d, J = 7.9 Hz, 2H), 8.06 (d, J = 8.0 Hz, 2H), 7.76 (t, J = 7.4 Hz, 2H), 7.68 (d, J = 7.8 Hz, 2H), 7.46 (d, J = 7.7 Hz, 2H), 7.43–7.32 (m, 6H), 4.04 (s, 3H), 2.66 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 165.8, 156.2, 142.7, 142.6, 141.3, 140.7, 138.0, 133.1, 132.7, 132.6, 130.6, 129.6, 129.4, 129.3, 129.2, 128.8, 128.3, 124.8, 117.8, 53.3, 21.7. ^{19}F NMR (376 MHz, DMSO): δ -143.7. IR (KBr): 3364, 2921, 2853, 1722, 1642, 1517, 1371, 1214, 1054, 531 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{36}\text{H}_{26}\text{NO}_2^+$ $[\text{M}]^+$ 504.1958 found 504.1949.




7-(3-nitrophenyl)-14-(p-tolyl)dibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8s)

Yellow solid (78.5 mg, 68% yield). DCM / MeOH = 100:1, R_f = 0.25. ^1H NMR (400 MHz, DMSO): δ 8.96 (s, 1H), 8.84 (d, J = 8.4 Hz, 1H), 8.58 (d, J = 9.5 Hz, 2H), 8.40 (d, J = 8.0 Hz, 1H), 8.34–8.24 (m, 3H), 7.80 (t, J = 7.3 Hz, 2H), 7.73 (d, J = 7.7 Hz, 2H), 7.50 (d, J = 8.6 Hz, 4H), 7.46–7.36 (m, 4H), 2.69 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 156.3, 150.2, 143.1, 141.4, 140.8, 139.5, 138.0, 135.4, 133.5, 132.7, 132.7, 130.6, 129.4, 129.3, 129.3, 128.8, 128.4, 127.1, 124.9, 124.8, 118.0, 21.8. ^{19}F NMR (376 MHz, CDCl_3): δ -153.7. IR (KBr): 3467, 3087, 1602, 1529, 1359, 1214, 1055, 822, 744 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{34}\text{H}_{23}\text{N}_2\text{O}_2^+$ $[\text{M}]^+$ 491.1754 found 491.1744.




7-(3,5-dimethylphenyl)-14-(p-tolyl)dibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8t)

Yellow solid (81.8 mg, 73% yield). DCM / MeOH = 100:1, R_f = 0.29. ^1H NMR (400 MHz, DMSO): δ 8.58 (d, J = 9.4 Hz, 2H), 8.25 (d, J = 7.8 Hz, 2H), 7.82–7.64 (m, 5H), 7.57–7.38 (m, 10H), 2.69 (s, 3H), 2.56 (s, 6H). ^{13}C NMR (100 MHz, DMSO): δ 155.8, 142.9, 141.6, 141.3, 140.4, 138.8, 138.1, 133.6, 132.7, 132.6, 130.5, 129.4, 129.4, 129.2, 128.9, 128.2, 126.1, 124.8, 118.0, 21.8, 21.4. ^{19}F NMR (376 MHz, CDCl_3): δ -154.3. IR (KBr): 3475, 2922, 1605, 1519, 1370, 1212, 1053, 823, 756 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{36}\text{H}_{28}\text{N}^+$ $[\text{M}]^+$ 474.2216 found 474.2207.




7-(3,5-dimethoxyphenyl)-14-(p-tolyl)dibenzo[a,j]acridin-7-ium tetrafluoroborate (8u)

Yellow solid (66.5 mg, 56% yield). DCM / MeOH = 100:1, R_f = 0.26. ^1H NMR (400 MHz, DMSO): δ 8.61 (d, J = 9.4 Hz, 2H), 8.28 (d, J = 7.8 Hz, 2H), 7.80 (t, J = 7.2 Hz, 2H), 7.73 (d, J = 7.7 Hz, 2H), 7.59 (d, J = 9.4 Hz, 2H), 7.51 (d, J = 7.6 Hz, 2H), 7.45–7.37 (m, 4H), 7.22 (s, 2H), 7.12 (s, 1H), 3.94 (s, 6H), 2.70 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 162.7, 155.9, 142.8, 141.3, 140.5, 140.3, 138.0, 132.7, 132.7, 130.5, 129.4, 129.3, 129.2, 128.8, 128.3, 124.7, 118.1, 107.3, 103.7, 56.5, 21.8. ^{19}F NMR (376 MHz, CDCl_3): δ -154.3. IR (KBr): 3661, 3057, 2899, 1728, 1553, 1326, 1218, 1035, 754 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{36}\text{H}_{28}\text{NO}_2^+$ $[\text{M}]^+$ 506.2115 found 506.2107.




7-(3,5-bis(trifluoromethyl)phenyl)-14-(p-tolyl)dibenzo[a,j]acridin-7-ium tetrafluoroborate (8v)

Yellow solid (85.5 mg, 64% yield). DCM / MeOH = 100:1, R_f = 0.28. ^1H NMR (400 MHz, DMSO): δ 8.82 (s, 3H), 8.58 (d, J = 9.5 Hz, 2H), 8.30 (d, J = 7.9 Hz, 2H), 7.80 (t, J = 7.4 Hz, 2H), 7.72 (d, J = 7.6 Hz, 2H), 7.53 (d, J = 9.4 Hz, 2H), 7.48 (d, J = 7.6 Hz, 2H), 7.46–7.36 (m, 4H), 2.69 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 156.3, 143.2, 141.5, 141.1, 140.4, 137.9, 134.0 (q, J = 34.2 Hz), 132.7, 132.7, 130.7, 129.5, 129.3, 129.2, 128.7, 128.5, 126.8 (q, J = 3.7 Hz), 124.8, 123.2 (q, J = 271.6 Hz), 118.0, 21.8. ^{19}F NMR (376 MHz, CDCl_3): δ -62.7, -154.2. IR (KBr): 3468, 2980, 1612, 1521, 1369, 1278, 1186, 1053, 747 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{36}\text{H}_{22}\text{F}_6\text{N}^+$ $[\text{M}]^+$ 582.1651 found 582.1648.



7-(naphthalen-2-yl)-14-(p-tolyl)dibenzo[a,j]acridin-7-ium tetrafluoroborate (8w)


Yellow solid (71.2 mg, 61% yield). DCM / MeOH = 100:1, R_f = 0.27. ^1H NMR (400 MHz, DMSO): δ 8.62–8.49 (m, 4H), 8.37 (d, J = 8.2 Hz, 1H), 8.26 (d, J = 7.8 Hz, 2H), 8.21 (d, J = 7.9 Hz, 1H), 8.00 (d, J = 8.1 Hz, 1H), 7.95–7.83 (m, 2H), 7.83–7.76 (m, 2H), 7.74 (d, J = 7.3 Hz, 2H), 7.54 (d, J = 7.3 Hz, 2H), 7.49–7.40 (m, 6H), 2.71 (s, 3H). ^{13}C NMR (100 MHz, DMSO): δ 156.0, 143.2, 141.3, 140.5, 138.1, 136.3, 134.3, 133.7, 132.7, 132.6, 132.2, 130.5, 129.4, 129.3, 129.2, 128.9, 128.7, 128.6, 128.3, 125.3, 124.9, 118.1, 21.8. ^{19}F NMR (376 MHz, CDCl_3): δ -154.2. IR (KBr): 3648, 3039, 2931, 1674, 1528, 1370, 1216, 749 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{38}\text{H}_{26}\text{N}^+$ $[\text{M}]^+$ 496.2060 found 496.2052.




7-(dibenzo[*b,d*]thiophen-2-yl)-14-(*p*-tolyl)dibenzo[*a,j*]acridin-7-ium tetrafluoroborate (8x)

Yellow solid (70.2 mg, 55% yield). DCM / MeOH = 100:1, R_f = 0.26. ^1H NMR (400 MHz, CDCl_3): δ 8.61 (s, 1H), 8.35 (d, J = 8.4 Hz, 1H), 8.27 (d, J = 7.9 Hz, 1H), 8.17 (d, J = 9.5 Hz, 2H), 8.02–7.93 (m, 3H), 7.74 (d, J = 8.5 Hz, 1H), 7.66–7.56 (m, 5H), 7.53–7.48 (m, 3H), 7.45 (d, J = 8.8 Hz, 2H), 7.37 (d, J = 9.4 Hz, 2H), 7.30 (d, J = 7.6 Hz, 2H), 2.70 (s, 3H). ^{13}C NMR (100 MHz, CDCl_3): δ 156.7, 143.0, 142.7, 141.0, 140.5, 140.2, 138.0, 137.8, 135.0, 134.3, 132.4, 132.1, 129.8, 129.4, 129.3, 129.3, 128.8, 128.4, 127.9, 125.7, 125.6, 125.4, 125.2, 123.0, 122.9, 121.8, 117.2, 21.7. ^{19}F NMR (376 MHz, CDCl_3): δ -154.2. IR (KBr): 3468, 2924, 1723, 1600, 1519, 1473, 1370, 1216, 1055, 823, 602 cm^{-1} . HRMS (ESI) m/z Calcd for $\text{C}_{40}\text{H}_{26}\text{NS}^+$ $[\text{M}]^+$ 552.1780 found 552.1774.


E. Mechanism studies



An oven-dried vial equipped with a stir bar was charged with naphthalen-2-amine **3** (28.6 mg, 0.2 mmol), $\text{In}(\text{OTf})_3$ (11.2 mg, 0.02 mmol) and dissolved with 1 mL ethylene glycol, then stirred the mixture for 5 minutes. After that, benzaldehyde **4** (10.6 mg, 0.1 mmol) was slowly added. Under argon atmosphere, the mixture was stirred at 120 °C for 12 hours in an oil bath. After the reaction was completed, it was cooled to room temperature. The mixture was diluted with 25 mL CH_2Cl_2 and washed with saturated aqueous NaHCO_3 , then concentrated in vacuo. The residue was purified by silica gel chromatography to afford the product **2a** (16.1 mg, 45% yield).




An oven-dried vial equipped with a stir bar was charged with N-(naphthalen-2-yl)-1-phenylmethanimine **1a** (46.2 mg, 0.2 mmol), In(OTf)₃ (11.2 mg, 0.02 mmol), naphthalen-2-ol **6** (28.8 mg, 0.2 mmol) and dissolved with 1.5 mL ethylene glycol. Under argon atmosphere, the mixture was stirred at 120 °C for 12 hours in an oil bath. After the reaction was completed, the mixture was diluted with 25 mL EtOAc and washed with saturated aqueous NaHCO₃, then concentrated in vacuo. The residue was purified by silica gel chromatography to afford the product **2a** (23.2 mg, 65% yield) and product **7** (6.2 mg, 17% yield).




14-phenyl-14H-dibenzo[*a,j*]xanthene (**7**)

White solid (12.2 mg, 17% yield). PE / EA = 50:1, *R*_f = 0.34. ¹H NMR (400 MHz, CDCl₃): δ 8.49 (d, *J* = 7.4 Hz, 2H), 7.98–7.78 (m, 4H), 7.74–7.41 (m, 8H), 7.24 (t, *J* = 7.4 Hz, 2H), 7.08 (t, *J* = 7.4 Hz, 1H), 6.58 (s, 1H). ¹³C NMR (100 MHz, CDCl₃): δ 148.8, 145.1, 131.5, 131.1, 128.9, 128.8, 128.5, 128.3, 126.8, 126.4, 124.3, 122.8, 118.1, 117.4, 38.1. The identity of **10** was confirmed based on reported NMR spectra.^[5]




An oven-dried vial equipped with a stir bar was charged with (E)-N-(6-bromonaphthalen-2-yl)-1-phenylmethanimine **1e'** (30.9 mg, 0.1 mmol), (E)-N-(naphthalen-2-yl)-1-(4-(trifluoromethyl)phenyl)methanimine **1i** (29.9 mg, 0.1 mmol), In(OTf)₃ (5.7 mg, 0.02 mmol) and dissolved with 0.4 mL ethylene glycol. Under argon atmosphere, the mixture was stirred at 120 °C for 12 hours in an oil bath. After the reaction was completed, it was cooled to room temperature. The mixture was diluted with 25 mL EtOAc and washed with saturated aqueous NaHCO₃, then concentrated in vacuo. The residue was purified by silica gel chromatography to afford the product **2a** (2.9 mg, 8% yield), **2i** (13.6 mg, 32% yield), **2e'** (9.4 mg, 24% yield), **2i'** (11.1 mg, 19% yield).



An oven-dried vial equipped with a stir bar was charged with naphthalen-2-amine **3** (14.1 mg, 0.1 mmol), In(OTf)₃ (11.2 mg, 0.02 mmol), (2-aminophenyl)(phenyl)methanol **5** (20.0 mg, 0.1

mmol) and dissolved with 0.4 mL ethylene glycol. Under argon atmosphere, the mixture was stirred at 120 °C for 12 hours in an oil bath. After the reaction was completed, the mixture was diluted with 25 mL EtOAc and washed with saturated aqueous NaHCO₃, then concentrated in vacuo. The residue was purified by silica gel chromatography to afford the product **2j'** (23.6 mg, 77% yield).




(S)-12-phenyl-7,12-dihydrobenzo[a]acridine (**2j'**)

White solid (17.8 mg, 58% yield). PE / DCM = 8:1, *R*_f = 0.31. ¹H NMR (400 MHz, CDCl₃): δ 8.01 (d, *J* = 8.5 Hz, 1H), 7.81 (d, *J* = 8.1 Hz, 1H), 7.76 (d, *J* = 8.7 Hz, 1H), 7.51–7.44 (m, 2H), 7.40–7.30 (m, 3H), 7.26–7.08 (m, 5H), 6.99 (t, *J* = 7.4 Hz, 1H), 6.86 (d, *J* = 7.9 Hz, 1H), 6.33 (s, 1H), 6.03 (s, 1H). ¹³C NMR (100 MHz, CDCl₃): δ 147.1, 137.8, 136.8, 132.2, 129.8, 129.4, 128.7, 128.6, 128.6, 127.1, 127.1, 126.8, 126.2, 123.9, 122.8, 122.3, 121.5, 116.7, 114.1, 113.6, 44.1. IR (KBr): 3448, 3015, 1629, 1519, 1467, 1420, 1285, 1206, 1155 cm⁻¹. HRMS (ESI) *m/z* Calcd for C₃₁H₂₀N [M+H]⁺ 308.1439, found 308.1433.

F. Synthetic transformation of products




An oven-dried vial equipped with a stir bar was charged with photoredox catalyst **8c** (5.8 mg, 10 mmol%), [1,1'-biphenyl]-2-carboxylic acid **9** (19.8 mg, 0.1 mmol) and dissolved with 1 mL MeCN. Under oxygen atmosphere, 1,8-Diazabicyclo[5.4.0]-undec-7-ene was slowly added to the tube (3.1 mg, 0.02 mmol) and the mixture was stirred in photoreactor at room temperature for 20 hours. After the reaction was completed, the reaction mixture was concentrated in vacuo. The residue was purified by silica gel chromatography to afford the product **10** (12.5 mg, 64% yield).




6H-benzo[c]chromen-6-one (**10**)

White solid (12.5 mg, 64% yield). PE/ EA = 20:1, *R*_f = 0.34. ¹H NMR (400 MHz, CDCl₃): δ 8.41 (dd, *J* = 8.0, 1.6 Hz, 1H), 8.12 (d, *J* = 8.1 Hz, 1H), 8.06 (d, *J* = 7.9 Hz, 1H), 7.83 (t, *J* = 8.3 Hz, 1H), 7.59 (t, *J* = 7.7 Hz, 1H), 7.49 (t, *J* = 8.5 Hz, 1H), 7.41–7.32 (m, 2H). ¹³C NMR (100 MHz, CDCl₃): δ 161.1, 151.3, 134.8, 134.7, 130.5, 130.4, 128.8, 124.5, 122.7, 121.6, 121.3, 118.0, 117.7. The identity of **10** was confirmed based on reported NMR spectra.^[6]




An oven-dried vial equipped with a stir bar was charged with photoredox catalyst **8c** (5.8 mg, 10 mmol%), pyrazole **12** (13.6 mg, 0.2 mmol) and 2,2,6,6-Tetramethylpiperidine-1-oxyl (3.1 mg, 0.02 mmol). Under oxygen atmosphere, Anisole **11** (10.8 mg, 0.1 mmol) dissolved with 1 mL CH₂Cl₂ and slowly added to the tube and the mixture was stirred in photoreactor at room temperature for 20 hours. After the reaction was completed, the reaction mixture concentrated in vacuo. The residue was purified by silica gel chromatography to afford the product **13** (9.5 mg, 54% yield).



1-(4-methoxyphenyl)-1H-1,2,4-triazole (**13**)

Yellow oil (9.5 mg, 54% yield). PE / EA = 10:1, *R*_f = 0.32. ¹H NMR (400 MHz, CDCl₃): δ 7.91 (d, *J* = 2.7 Hz, 1H), 7.84–7.64 (m, 3H), 7.07 (d, *J* = 9.4 Hz, 2H), 6.53 (s, 1H), 3.94 (s, 3H). ¹³C NMR (100 MHz, CDCl₃): δ 158.3, 140.6, 126.8, 120.9, 114.5, 107.1, 55.5. The identity of **13** was confirmed based on reported NMR spectra.^[7]

Scale-up reaction (2.5 mmol)



An oven-dried vial equipped with a stir bar was charged with N-(naphthalen-2-yl)-1-phenylmethanimines **1a** (1.16 g, 2.5 mmol) and In(OTf)₃ (281.0 mg, 0.5 mmol). The reaction mixture was placed under a positive pressure of argon and subjected to three evacuation/backfilling cycles under high vacuum. Ethylene glycol (typically, 15 mL) was added and the reaction mixture was stirred at 120 °C for 12 hours in an oil bath. After the reaction was completed, it was cooled to room temperature. The mixture was diluted with EtOAc and washed with saturated aqueous NaHCO₃, then concentrated in vacuo. The residue was purified by silica gel chromatography to afford the dihydroacridine products **2a** (731.8 mg, 82% yield).

Under an Ar atmosphere, product **2a** (731.3 mg, 2.0 mmol), Pd(OAc)₂ (23.9 mg, 0.1 mmol), xantphos (115.8 mg, 0.2 mmol) and *t*-BuOK (459.2 mg, 2 equiv.) were added to the pressure tube equipped with a stir bar and dissolved with 15 mL toluene, then iodobenzene (611.1 mg, 3.0 mmol) was slowly added and the mixture stirred for 10 minutes. After that, product **2a** dissolved with 10 mL toluene was slowly added and the mixture stirred at 120 °C for 4 hours. After the reaction was completed, the reaction mixture was filtered and concentrated in vacuo.


The residue from previous step was dissolved with 30 mL CH₃CN and added to a round bottom flask equipped with a stir bar, it was stirred at 0 °C and 2,3-dichloro-5,6-dicyano-1,4-benzquinone (926.6 mg, 2 equiv.) was slowly added at 0 °C. The reaction was stirred at room

temperature for 2 hours. After the reaction was completed, the mixture concentrated in vacuo, the mixture was diluted with 50 mL CH₂Cl₂ and washed with pure water, the organic phases were dried over anhydrous MgSO₄ and concentrated in vacuo. The residue was dissolved in CH₂Cl₂ (50 mL) and added to a round bottom flask equipped with a stir bar, it was stirred with NaBF₄ aqueous solution (40 mL) for 8 hours. After the reaction was completed, it was washed with pure water. The organic phase was dried over anhydrous MgSO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography to afford the product **8a** (789.2 mg, 76% yield).




G. References

1. Hu, Y., Nan, J., Yin, J., Huang, G., Ren, X., Ma, Y. Rhodium-Catalyzed Dehydrogenative Annulation of N-Arylmethanimines with Vinylene Carbonate for Synthesizing Quinolines. *Org. Lett.* **23**, 8527–8532 (2021).
2. Suresh, R., Sakthinathan, S. P., Kamalakkannan, D., Ranganathan, K., Sathiyamoorthi, K., Mala, V., Arulkumaran, R., Vijayakumar, S., Sundararajan, R., Vanangamudi, G., Subramanian, M., Thirunarayanan, G., Vanaja, G., Kanagambal, P. Solvent-free synthesis of azomethines, spectral correlations and antimicrobial activities of some E-benzylidene-4-chlorobenzenamines *Bull. Chem. Soc. Ethiop.* **29**, 275–290 (2015).
3. Saha, S., Rajput, L., Joseph, S., Mishra, M. K., Ganguly, S., Desiraju, G. R. *CrystEngComm*. IR spectroscopy as a probe for C–H···X hydrogen bonded supramolecular synthons. **17**, 1273–1290 (2015).
4. Sek, D., Siwy, M., Bijak, K., Filapek, M., Malecki, G., Nowak, E., Sanetra, J., Jedryka, A. J., Laba, K., Lapkowski, M., Balcerzak, E. Optical and electrochemical properties of novel thermally stable Schiff bases bearing naphthalene unit. *J. Electroanal. Chem.* **751**, 128–136 (2015).
5. Sek, D., Siwy, M., Grucela, M., Małeck, G., Nowak, E. M., Lewinska, G., Santera, J., Laba, K., Lapkowski, M., Kotowicz, S., Balcerzak, E. S. New anthracene-based Schiff bases: Theoretical and experimental investigations of photophysical and electrochemical properties *Spectrochim. Acta. A Mol. Biomol. Spectrosc.* **175**, 24–35 (2017).
6. Niu, K., Zhou, P., Ding, L., Hao, Y., Liu, Y., Song, H., Wang, Q. Photoelectrochemical Decarboxylative C–H Alkylation of Quinoxalin-2(1H)-ones. *ACS. Sustain. Chem. Eng.* **9**, 16820–16828 (2021).
7. Romero, N. A., Margrey, K. A., Tay, N. E., Nicewicz, D. A. Site-selective arene C–H amination via photoredox catalysis. *Science*. **349**, 1326–1330 (2015).



H. Spectrophotometric and electrochemical data





Spectrophotometric and electrochemical data of **8a** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $\text{nBu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8b** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8c** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $\text{nBu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8d** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $\text{nBu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8e** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8f** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8g** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8h** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8i** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8j** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8k** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8I** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8m** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 485$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8n** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $\text{nBu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38 \text{ V vs SCE}$ in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 485 \text{ nm}$, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5} \text{ M}$, $\lambda_{\text{ex}} = 340 \text{ nm}$). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **80** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8p** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8q** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M nBu₄N·PF₆ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference (Fc/Fc⁺ = 0.38 V vs SCE in DCM). top right: UV-visible absorption (red line) and emission (λ_{ex} = 447 nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K (c ≈ 10⁻⁵ M, λ_{ex} = 340 nm). Experimental decays in yellow, exponential fits in red, residual traces in green.




Spectrophotometric and electrochemical data of **8r** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8s** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 485$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8t** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M nBu₄N·PF₆ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference (Fc/Fc⁺ = 0.38 V vs SCE in DCM). top right: UV-visible absorption (red line) and emission (λ_{ex} = 447 nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K (c ≈ 10⁻⁵ M, λ_{ex} = 340 nm). Experimental decays in yellow, exponential fits in red, residual traces in green.






Spectrophotometric and electrochemical data of **8u** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $\text{nBu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.



Spectrophotometric and electrochemical data of **8v** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $n\text{Bu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 485$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.




Spectrophotometric and electrochemical data of **8w** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $\text{nBu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.




Spectrophotometric and electrochemical data of **8x** top left: Cyclic voltammograms measured in a deaerated DCM with 0.1 M $\text{nBu}_4\text{N}^+\text{PF}_6^-$ with a scan rate of 100 mV/s, a platinum working electrode and with ferrocene as the internal reference ($\text{Fc}/\text{Fc}^+ = 0.38$ V vs SCE in DCM). top right: UV-visible absorption (red line) and emission ($\lambda_{\text{ex}} = 447$ nm, black line) spectra, in DCM. bottom: Fitting of luminescence decays in DCM at 298 K ($c \approx 10^{-5}$ M, $\lambda_{\text{ex}} = 340$ nm). Experimental decays in yellow, exponential fits in red, residual traces in green.

I. NMR spectra


^1H NMR of **2a** (400 MHz, CDCl_3)




^{13}C NMR of **2a** (100 MHz, CDCl_3)




^1H NMR of **2b** (400 MHz, CDCl_3)




^{13}C NMR of **2b** (100 MHz, CDCl_3)




¹H NMR of 2c (400 MHz, DMSO)




¹³C NMR of 2c (100 MHz, DMSO)




^1H NMR of **2d** (400 MHz, DMSO)




^{13}C NMR of **2d** (100 MHz, DMSO)




^1H NMR of **2e** (400 MHz, CDCl_3)




^{13}C NMR of **2e** (100 MHz, CDCl_3)




^1H NMR of **2f** (400 MHz, CDCl_3)




^{13}C NMR of **2f** (100 MHz, CDCl_3)




^1H NMR of **2g** (400 MHz, CDCl_3)




^{13}C NMR of **2g** (100 MHz, CDCl_3)




^1H NMR of **2h** (400 MHz, CDCl_3)




^{13}C NMR of **2h** (100 MHz, CDCl_3)




^{19}F NMR of **2h** (376 MHz, CDCl_3)




^1H NMR of **2i** (400 MHz, CDCl_3)



^{13}C NMR of **2i** (100 MHz, CDCl_3)



^{19}F NMR of **2i** (376 MHz, CDCl_3)



Chemical structure of (S)-1-(2,2,2-trifluorophenyl)-1,2,3,4-tetrahydronaphthalen-1-ylamine is shown. The structure features a tetrahydronaphthalene core with a trifluorophenyl group (SCF₃) attached to the 1-position and an amine group (NH) at the 1-position. The amine proton is labeled 'H'.

The ¹H NMR spectrum (400 MHz, CDCl₃) shows the following peaks and integrations:


- Aromatic region (6.6–8.4 ppm):
 - Peak at ~8.3 ppm (integration: 2.1H)
 - Peak at ~8.0 ppm (integration: 2.1H)
 - Peak at ~7.8 ppm (integration: 2.1H)
 - Peak at ~7.6 ppm (integration: 2.1H)
 - Peak at ~7.4 ppm (integration: 4.1H)
 - Peak at ~7.2 ppm (integration: 2.0H)
 - Peak at ~7.0 ppm (integration: 1.0H)
 - Peak at ~6.8 ppm (integration: 1.0H)
- Aliphatic region (1.0–2.1 ppm):
 - Peak at ~1.7 ppm (integration: 2.1H)
 - Peak at ~1.5 ppm (integration: 2.1H)
 - Peak at ~1.3 ppm (integration: 2.1H)
 - Peak at ~1.1 ppm (integration: 2.1H)
 - Peak at ~0.9 ppm (integration: 1.0H)
 - Peak at ~0.7 ppm (integration: 1.0H)

Chemical structure of 1-(4-(trifluoromethyl)phenyl)-2,3-dihydro-1H-benzo[e][1,2-b:4,5-b']difuran is shown. The structure features a central nitrogen atom (N) bonded to a hydrogen atom (H) and a 4-(trifluoromethyl)phenyl group (SCF₃). The nitrogen atom is also bonded to two fused benzene rings, forming a difuran system.


¹³C NMR spectrum (f1 (ppm)) showing chemical shifts (ppm) for the compound. The spectrum displays a series of peaks corresponding to the carbon atoms in the molecule, with the following chemical shifts (ppm) listed above the peaks:

- 148.8
- 136.3
- 136.1
- 135.9
- 130.9
- 130.2
- 128.9
- 128.8
- 128.7
- 127.1
- 123.2
- 121.8
- 121.6
- 116.8
- 113.7
- 38.6


^{19}F NMR of **2j** (376 MHz, CDCl_3)




^1H NMR of **2k** (400 MHz, DMSO)




^{13}C NMR of **2k** (100 MHz, DMSO)




^1H NMR of **2l** (400 MHz, DMSO)




^{13}C NMR of **2l** (100 MHz, DMSO)




^1H NMR of **2m** (400 MHz, DMSO)




^{13}C NMR of **2m** (100 MHz, DMSO)




^1H NMR of **2n** (400 MHz, CDCl_3)




^{13}C NMR of **2n** (100 MHz, CDCl_3)




^1H NMR of **2o** (400 MHz, CDCl_3)




^{13}C NMR of **2o** (100 MHz, CDCl_3)




^1H NMR of **2p** (400 MHz, CDCl_3)




^{13}C NMR of **2p** (100 MHz, CDCl_3)




^1H NMR of **2q** (400 MHz, CDCl_3)





^{13}C NMR of **2q** (100 MHz, CDCl_3)




^1H NMR of **2r** (400 MHz, CDCl_3)




¹³C NMR of **2r** (100 MHz, CDCl₃)

¹H NMR of **2s** (400 MHz, CDCl₃)


^{13}C NMR of **2s** (100 MHz, CDCl_3)




^1H NMR of **2t** (400 MHz, CDCl_3)




^{13}C NMR of **2t** (100 MHz, CDCl_3)




^1H NMR of **2u** (400 MHz, CDCl_3)




^{13}C NMR of **2u** (100 MHz, CDCl_3)




^1H NMR of **2v** (400 MHz, CDCl_3)




^{13}C NMR of **2v** (100 MHz, CDCl_3)




^1H NMR of **2w** (400 MHz, CDCl_3)




^{13}C NMR of **2w** (100 MHz, CDCl_3)




^1H NMR of **2x** (400 MHz, DMSO)




^{13}C NMR of **2x** (100 MHz, DMSO)




^1H NMR of **2y** (400 MHz, DMSO)




^{13}C NMR of **2y** (100 MHz, DMSO)




^{19}F NMR of **2y** (376 MHz, DMSO)




^1H NMR of **2z** (400 MHz, CDCl_3)




^{13}C NMR of **2z** (100 MHz, CDCl_3)




¹H NMR of **2a'** (400 MHz, CDCl₃)




¹³C NMR of **2a'** (100 MHz, CDCl₃)




^1H NMR of **2b'** (400 MHz, DMSO)




^{13}C NMR of **2b'** (100 MHz, DMSO)




^1H NMR of **2c'** (400 MHz, CDCl_3)




^{13}C NMR of **2c'** (100 MHz, CDCl_3)




¹H NMR of **2d'** (400 MHz, CDCl₃)




¹³C NMR of **2d'** (100 MHz, CDCl₃)




^1H NMR of **2e'** (400 MHz, DMSO)




^{13}C NMR of **2e'** (100 MHz, DMSO)




^1H NMR of **2f'** (400 MHz, CDCl_3)




^{13}C NMR of **2f'** (100 MHz, CDCl_3)




¹H NMR of **2g'** (400 MHz, DMSO)




¹³C NMR of **2g'** (100 MHz, DMSO)




^1H NMR of **2h'** (400 MHz, DMSO)




^{13}C NMR of **2h'** (100 MHz, DMSO)




^1H NMR of **2i'** (400 MHz, DMSO)




^{13}C NMR of **2i'** (100 MHz, DMSO)




^{19}F NMR of **2i'** (376 MHz, DMSO)




^1H NMR of **8a** (400 MHz, CDCl_3)




^{13}C NMR of **8a** (100 MHz, CDCl_3)




^{19}F NMR of **8a** (376 MHz, CDCl_3)




¹H NMR of **8b** (400 MHz, DMSO)




¹³C NMR of **8b** (100 MHz, DMSO)




^{19}F NMR of **8b** (376 MHz, CDCl_3)




^1H NMR of **8c** (400 MHz, CDCl_3)




^{13}C NMR of **8c** (100 MHz, CDCl_3)




^{19}F NMR of **8c** (376 MHz, CDCl_3)




^1H NMR of **8d** (400 MHz, CDCl_3)




^{13}C NMR of **8d** (100 MHz, CDCl_3)




^{19}F NMR of **8d** (376 MHz, CDCl_3)




^1H NMR of **8e** (400 MHz, DMSO)




^{13}C NMR of **8e** (100 MHz, DMSO)




^{19}F NMR of **8e** (376 MHz, CDCl_3)




¹H NMR of **8f** (400 MHz, DMSO)




¹³C NMR of **8f** (100 MHz, DMSO)




^{19}F NMR of **8f** (376 MHz, CDCl_3)




^1H NMR of **8g** (400 MHz, CDCl_3)




^{13}C NMR of **8g** (100 MHz, CDCl_3)




^{19}F NMR of **8g** (376 MHz, CDCl_3)




^1H NMR of **8h** (400 MHz, CDCl_3)




^{13}C NMR of **8h** (100 MHz, CDCl_3)




^{19}F NMR of **8h** (376 MHz, CDCl_3)




^1H NMR of **8i** (400 MHz, DMSO)




^{13}C NMR of **8i** (100 MHz, DMSO)




^{19}F NMR of **8i** (376 MHz, CDCl_3)




^1H NMR of **8j** (400 MHz, CDCl_3)




^{13}C NMR of **8j** (100 MHz, CDCl_3)




^{19}F NMR of **8j** (376 MHz, CDCl_3)




^1H NMR of **8k** (400 MHz, CDCl_3)




^{13}C NMR of **8k** (100 MHz, CDCl_3)




^{19}F NMR of **8k** (376 MHz, CDCl_3)




^1H NMR of **8l** (400 MHz, CDCl_3)




^{13}C NMR of **8l** (100 MHz, CDCl_3)




^{19}F NMR of **8l** (376 MHz, CDCl_3)




^1H NMR of **8m** (400 MHz, CDCl_3)




^{13}C NMR of **8m** (100 MHz, CDCl_3)




^{19}F NMR of **8m** (376 MHz, CDCl_3)




^1H NMR of **8n** (400 MHz, CDCl_3)




^{13}C NMR of **8n** (100 MHz, CDCl_3)




^{19}F NMR of **8n** (376 MHz, CDCl_3)




^1H NMR of **8o** (400 MHz, DMSO)




^{13}C NMR of **8o** (100 MHz, DMSO)




^{19}F NMR of **8o** (376 MHz, CDCl_3)




¹H NMR of **8p** (400 MHz, DMSO)




¹³C NMR of **8p** (100 MHz, DMSO)




^{19}F NMR of **8p** (376 MHz, CDCl_3)




^1H NMR of **8q** (400 MHz, DMSO)




^{13}C NMR of **8q** (100 MHz, DMSO)




^{19}F NMR of **8q** (376 MHz, CDCl_3)




¹H NMR of **8r** (400 MHz, DMSO)




¹³C NMR of **8r** (100 MHz, DMSO)




^{19}F NMR of **8r** (376 MHz, DMSO)




^1H NMR of **8s** (400 MHz, DMSO)




^{13}C NMR of **8s** (100 MHz, DMSO)




^{19}F NMR of **8s** (376 MHz, CDCl_3)




¹H NMR of **8t** (400 MHz, DMSO)




¹³C NMR of **8t** (100 MHz, DMSO)




^{19}F NMR of **8t** (376 MHz, CDCl_3)




^1H NMR of **8u** (400 MHz, DMSO)



¹³C NMR of **8u** (100 MHz, DMSO)



¹⁹F NMR of **8u** (376 MHz, CDCl_3)



Chemical structure of compound 10: A fluorene derivative with a 4-methylphenyl group at position 9, a 4-(trifluoromethyl)phenyl group at position 1, and a BF_4^- counterion.

^1H NMR spectrum (CDCl₃):


- Chemical shift range:** 0.0 to 12.0 ppm.
- Peak assignments and integration:**
 - Aromatic protons (7.4–8.8 ppm): Integration values include 3.0H, 2.0H, 2.0H, 2.0H, 2.1H, 2.0H, 2.0H, 4.1H.
 - CF_3 groups (~11.2 ppm): Integration value 3.0H.
 - Methoxy group (~3.8 ppm): Integration value 3.0H.
 - Methyl group (~2.3 ppm): Integration value 3.0H.

Chemical structure of the cation: 1-methyl-2-(pentafluorophenyl)-3,4,5,6-tetrahydronaphthalene-1-ylboronate. The structure shows a naphthalene ring system with a methyl group (Me) at position 1, a pentafluorophenyl group at position 2, and a boronate group at position 3. The boronate group is shown as a boron atom bonded to the naphthalene ring and a pentafluorophenyl group, with a negative charge on the boron atom. The counterion is BF_4^- .


^1H NMR spectrum (ppm):

- 156.3, 143.2, 141.5, 140.4, 137.9, 134.1, 133.8, 132.7, 130.7, 129.5, 129.3, 129.2, 128.7, 128.5, 126.8, 124.8, 124.6, 121.9, 118.0 (Aromatic region)
- 14.5, 14.1, 13.8, 13.5, 13.2, 12.9, 12.8 (Aliphatic region)


^{19}F NMR of **8v** (376 MHz, CDCl_3)




^1H NMR of **8w** (400 MHz, DMSO)




^{13}C NMR of **8w** (100 MHz, DMSO)




^{19}F NMR of **8w** (376 MHz, CDCl_3)




^1H NMR of **8x** (400 MHz, CDCl_3)




^{13}C NMR of **8x** (100 MHz, CDCl_3)




^{19}F NMR of **8x** (376 MHz, CDCl_3)




^1H NMR of **10** (400 MHz, CDCl_3)




¹³C NMR of **10** (100 MHz, CDCl₃)




¹H NMR of **13** (400 MHz, CDCl₃)




^{13}C NMR of **13** (100 MHz, CDCl_3)




^1H NMR of **7** (400 MHz, CDCl_3)



^{13}C NMR of **7** (100 MHz, CDCl_3)



^1H NMR of **2j'** (400 MHz, CDCl_3)



^{13}C NMR of **2j'** (100 MHz, CDCl_3)

