


**Production of virus-like particles of porcine circovirus type 2 in baculovirus
expression system and its application for antibody detection**


Yanwei Li, Pingping Yu, Yaxuan Bao, Yuwen Ren, Shaowei Zhao, Xuexian Zhang*
Beijing Kemufeng Biopharmaceutical Co., Ltd, No.25 Xiangrui Street Daxing District,
Beijing 102600 China

*Corresponding author: Xuexian Zhang

E-mail: xuexianz@hotmail.com



Coomassie blue stain shows overexpression of the purified protein. Lanes 1 is uninfected sf9 cell lysates; lanes 2 and 3 are cell lysates infected with pFastBac™Dual plasmid and recombinant baculovirus recombinant baculovirus passage 1, respectively.



Identification of recombinant protein expression by Western blot. Lanes 1 is uninfected Sf9 cell lysate; 2, Sf9 cells infected with pFastBac™Dual plasmid; lanes 3 to 7 are sf9 cell lysates at 24 h, 30 h, 36h, 42 h, 48 h after infection with recombinant baculovirus recombinant baculovirus passage 1, Lanes 4 to 11 are hi5 cell lysates at 24 h, 30 h, 36h, 42 h, 48 h after infection with recombinant baculovirus recombinant baculovirus passage 1; Lanes 12 is uninfected hi5 cell lysate.