

Table S 1: Correlations of Child Feeding practice

				Nutritional status of the child	Supplementary food started at recommended age	Mealtime frequency	Type of supplementary food started
Kendall's tau_b	Nutritional status of the child	Correlation Coefficient		1.000	-.275**	.597**	.842**
		Sig. (2-tailed)		.	.000	.000	.000
		N		206	206	206	206
		Bootstrap ^c	Bias	.000	.011	.000	.001
			Std. Error	.000	.061	.033	.012
			99% Confidence Interval	1.000	-.415	.485	.808
			Lower Upper	1.000	-.109	.686	.880
	Supplementary food started at recommended age	Correlation Coefficient		-.275**	1.000	-.295**	-.352**
		Sig. (2-tailed)		.000	.	.000	.000
		N		206	206	206	206
		Bootstrap ^c	Bias	.011	.000	.003	.010
			Std. Error	.061	.000	.051	.069
			99% Confidence Interval	-.415	1.000	-.414	-.522
			Lower Upper	-.109	1.000	-.168	-.161
	Mealtime frequency while home child feeding	Correlation Coefficient		.597**	-.295**	1.000	.595**
		Sig. (2-tailed)		.000	.000	.	.000
		N		206	206	206	206
		Bootstrap ^c	Bias	.000	.003	.000	-.002
			Std. Error	.033	.051	.000	.030
			99% Confidence Interval	.485	-.414	1.000	.512
			Lower Upper	.686	-.168	1.000	.671
	Type of supplementary food started	Correlation Coefficient		.842**	-.352**	.595**	1.000
		Sig. (2-tailed)		.000	.000	.000	.
		N		206	206	206	206
		Bootstrap ^c	Bias	.001	.010	-.002	.000
			Std. Error	.012	.069	.030	.000
			99% Confidence Interval	.808	-.522	.512	1.000
			Lower Upper	.880	-.161	.671	1.000

** . Correlation is significant at the 0.01 level (2-tailed).

* . Correlation is significant at the 0.05 level (2-tailed).

c. Unless otherwise noted, bootstrap results are based on 206 bootstrap samples