

Supplementary File 3

Reference list of included studies

1. Sherrington C, Lord SR. Home exercise to improve strength and walking velocity after hip fracture: a randomized controlled trial. *Arch Phys Med Rehabil.* 1997;**78**(2):208-212. doi:10.1016/s0003-9993(97)90265-3
2. Hauer K, Specht N, Schuler M, Bärtsch P, Oster P. Intensive physical training in geriatric patients after severe falls and hip surgery. *Age Ageing.* 2002;**31**(1):49-57. doi:10.1093/ageing/31.1.49
3. Sherrington C, Lord SR, Herbert RD. A randomised trial of weight-bearing versus non-weight-bearing exercise for improving physical ability in inpatients after hip fracture. *Aust J Physiother.* 2003;**49**(1):15-22. doi:10.1016/s0004-9514(14)60184-7
4. Binder EF, Brown M, Sinacore DR, Steger-May K, Yarasheski KE, Schechtman KB. Effects of extended outpatient rehabilitation after hip fracture: a randomized controlled trial. *JAMA.* 2004;**292**(7):837-846. doi:10.1001/jama.292.7.837
5. Sherrington C, Lord SR, Herbert RD. A randomized controlled trial of weight-bearing versus non-weight-bearing exercise for improving physical ability after usual care for hip fracture. *Arch Phys Med Rehabil.* 2004;**85**(5):710-716. doi:10.1016/s0003-9993(03)00620-8
6. Suetta C, Magnusson SP, Rosted A, et al. Resistance training in the early postoperative phase reduces hospitalization and leads to muscle hypertrophy in elderly hip surgery patients--a controlled, randomized study. *J Am Geriatr Soc.* 2004;**52**(12):2016-2022. doi:10.1111/j.1532-5415.2004.52557.x
7. Mangione KK, Craik RL, Tomlinson SS, Palombaro KM. Can elderly patients who have had a hip fracture perform moderate- to high-intensity exercise at home?. *Phys Ther.* 2005;**85**(8):727-739. PMID: 16048421
8. Mendelsohn ME, Overend TJ, Connelly DM, Petrella RJ. Improvement in aerobic fitness during rehabilitation after hip fracture. *Arch Phys Med Rehabil.* 2008;**89**(4):609-617. doi:10.1016/j.apmr.2007.09.036

9. Minna Mard, Johanna Vaha, Heinonen A. The effects of muscle strength and power training on mobility among older hip fracture patients. *Advances in Physiotherapy*. 2008;**10**(4): 195-202. doi: 10.1080/14038190801999570
10. Portegijs E, Kallinen M, Rantanen T, et al. Effects of resistance training on lower-extremity impairments in older people with hip fracture. *Arch Phys Med Rehabil*. 2008;**89**(9):1667-1674. doi:10.1016/j.apmr.2008.01.026
11. Mangione KK, Craik RL, Palombaro KM, Tomlinson SS, Hofmann MT. Home-based leg-strengthening exercise improves function 1 year after hip fracture: a randomized controlled study. *J Am Geriatr Soc*. 2010;**58**(10):1911-1917. doi:10.1111/j.1532-5415.2010.03076.x
12. Kui DQ. The effect of early weight-bearing and walking on elderly patients after internal fixation of femoral neck fractures. *Chin J Phys Med Rehabil*. 2011;(08): 614-616. doi:10.3760/cma.j.issn.0254-1424.2011.08.015
13. Yan Y. Balanced functional exercise in the rehabilitation of elderly patients with femoral neck fractures. *Chin J Mod Nur*. 2012;(24): 2938-2939.
14. Singh N A, Quine S, Clemson L M, et al. Effects of high-intensity progressive resistance training and targeted multidisciplinary treatment of frailty on mortality and nursing home admissions after hip fracture: a randomized controlled trial. *J Am Med Dir Assoc*. 2012;**13**(1): 24-30. doi:10.1016/j.jamda.2011.08.005
15. Sylliaas H, Brovold T, Wyller T B, et al. Prolonged strength training in older patients after hip fracture: a randomized controlled trial. *Age Ageing*. 2012;**41**(2): 206-212. doi:10.1093/ageing/afr164
16. Morishima Y, Mizushima T, Yamauchi K, et al. Effects of home-based interval walking training on thigh muscle strength and aerobic capacity in female total hip arthroplasty patients: a randomized, controlled pilot study. *Plos One*. 2014;**9**(9): e108690. doi:10.1371/journal.pone.0108690
17. Xing S, Zhang LL, Tang SY, et al. The design and application of household music rehabilitation exercises in the post-operation for elderly patient with hip fracture. *Orthop J Chin*. 2016;**24**(05): 470-473.
18. Zhao X. Application of home furnishing music rehabilitation gymnastics in

- discharged elderly patients with hip fracture. *Chin Nurs Res.* 2016;**30**(24): 3056-3058. doi:10.3969/j.issn.1009-6493.2016.24.039
19. van Ooijen MW, Roerdink M, Trekop M, Janssen TW, Beek PJ. The efficacy of treadmill training with and without projected visual context for improving walking ability and reducing fall incidence and fear of falling in older adults with fall-related hip fracture: a randomized controlled trial. *BMC Geriatr.* 2016;**16**(1):215. doi:10.1186/s12877-016-0388-x
 20. Xu YF. Rehabilitation effects of the comprehensive intervention of home-based balance training on the elderly with femoral neck fracture. *Chin J Mod Nurs.* 2017;**23**(13): 1797-1800. doi:10.3760/cma.j.issn.1674-2907.2017.13.022
 21. Zhang T, Xie Q, Wang J, et al. Effect of Intensive Core Muscles Training on General Function in Patients after Hip Arthroplasty. *Chin J Rehabil Theor and Prac.* 2017;**23**(10): 1171-1175. doi:10.3969/j.issn.1006-9771.2017.10.010
 22. Monticone M, Ambrosini E, Brunati R, et al. How balance task-specific training contributes to improving physical function in older subjects undergoing rehabilitation following hip fracture: a randomized controlled trial. *Clin Rehabil.* 2018;**32**(3): 340-351. doi:10.1177/0269215517724851
 23. Dong GM, Zhang W. Effect of Relaxation Training combined with Muscle Group Intensive Training on Patients with Hip Replacement. *Qilu J Nurs.* 2019;**25**(20): 19-22. doi:10.3969/j.issn.1006-7256.2019.20.006
 24. Kang ZM, Xi GQ, Chen YF, et al. Effect of hip rehabilitation exercise on functional rehabilitation of hip fracture patients after operation. *Chin J Prim Med Pharm.* 2019;**26**(16): 2041-2044. doi:10.3760/cma.j.issn.1008-6706.2019.16.032
 25. Wang G, Liu HJ, Li ZW, et al. Role of progressive resistance exercise in the rehabilitation of femoral intertrochanteric fracture after operation. *Journal of Zhengzhou University (Medical Sciences).* 2019;**54**(04): 615-618.
 26. Wu WY, Xu WG. Evaluation of the effect of early total weight-bearing training after cementless total hip arthroplasty. *Chin J Tiss Engin Res.* 2019;**23**(18): 2827-2832. doi:10.3969/j.issn.2095-4344.1196
 27. Stasi S, Papathanasiou G, Chronopoulos E, et al. The Effect of Intensive Abductor

- Strengthening on Postoperative Muscle Efficiency and Functional Ability of Hip-Fractured Patients: A Randomized Controlled Trial. *Indian J Orthop.* 2019;**53**(3): 407-419. doi:10.4103/ortho.IJOrtho_183_18
28. Cai Y, Wang J, Cen WX, et al. Effect of progressive resistance training on rehabilitation of patients with osteoporotic femoral neck fracture after hip arthroplasty. *Chin J Frontiers Med Sci.* 2020;**12**(11): 68-71.
 29. Chi K, Liu B, Yang XM. Application of early total weight bearing training in biological total hip replacement for elderly patients with unstable intertrochanteric fracture of femur. *Chin J Frontiers Med Sci.* 2020;**12**(12): 42-45.
 30. Li JH. Effect of intensive training of core muscle group on overall functional recovery in patients undergoing hip arthroplasty. *IMHGN.* 2020; **26**(12): 1765-1767. doi:10.3760/cma.j.issn.1007-1245.2020.12.035
 31. Qin DJ, Zhang Y, Zhang HF. Application of home furnishing music rehabilitation gymnastics in elderly patients discharged from hospital after intertrochanteric fracture of the femur. *Henan Med Res.* 2020;**29**(35): 6689-6691. doi:10.3969/j.issn.1004-437X.2020.35.067
 32. Sun ZL, Xu R. The effect of core muscle group training on rehabilitation and complications of elderly patients with femoral neck fracture after joint replacement. *Anhui J Med and Pharm.* 2020;**24**(07): 1359-1362. doi:10.3969/j.issn.1009-6469.2020.07.022
 33. Wang K, Zhang Z, Du YK. Effect of early weight-bearing and walking on patients with femoral neck fracture after internal fixation. *Contemp Med.* 2020;**26**(35): 50-53. doi:10.3969/j.issn.1009-4393.2020.35.020
 34. Xu LJ. Effect of home-based balance exercises on postoperative hip function, balance and compliance in elderly patients with femoral neck fractures. *Henan Med Res.* 2020;**29**(33): 6323-6324. doi:10.3969/j.issn.1004-437X.2020.33.074
 35. Kim P, Lee H, Choi W, et al. Effect of 4 Weeks of Anti-Gravity Treadmill Training on Isokinetic Muscle Strength and Muscle Activity in Adults Patients with a Femoral Fracture: A Randomized Controlled Trial. *Int J Environ Res Public*

Health. 2020;**17**(22). doi:10.3390/ijerph17228572

36. Oh M K, Yoo J I, Byun H, et al. Efficacy of Combined Antigravity Treadmill and Conventional Rehabilitation After Hip Fracture in Patients with Sarcopenia. *J Gerontol A Biol Sci Med Sci*. 2020; 75(10): 173-181. doi:10.1093/gerona/glaa158
37. Ding XY, Li J. Application of " Four More" family rehabilitation exercise in elderly patients with hip fracture. *Int J Nurs*. 2021;**40**(11):1973-1975. doi:10.3760/cma.j.cn221370-20200330-00571
38. Guo YT, Wang JW. Application of early rehabilitation exercise combined with family balance training in postoperative rehabilitation for elderly patients with femoral neck fracture. *Chin J Convale Med*. 2021;**30**(03): 266-268.
39. Liu Q, Li SS, Li XZ, et al. Effect of home-based balance training on hip function, lower limb mobility and balance in elderly patients with Pauwels type III femoral neck fractures. *Chin Gen Pract Nurs*. 2021;**19**(36):5110-5113. doi:10.12104/j.issn.1674-4748.2021.36.017
40. Wang DX, Lin Y, Zhang T, et al. Application of self-made rehabilitation exercises in postoperative rehabilitation of elderly patients with hip fracture. *Chin Evid Based Nurs*. 2021; **7**(15):2059-2061. doi:10.12102/j.issn.2095-8668.2021.15.015
41. Xu XX, Yang LF, Jing S, et al. Application effect of progressive resistance exercise training in elderly patients with intertrochanteric fracture after internal fixation. *Chin Med Herald*. 2021;**18**(09): 99-103.
42. Corna S, Arcolin I, Giardini M, et al. Addition of aerobic training to conventional rehabilitation after hip fracture: a randomized, controlled, pilot feasibility study. *Clin Rehabil*, 2021;**35**(4): 568-577. doi:10.1177/0269215520968694
43. Paulsson M, Ekholm C, Jonsson E, et al. Immediate Full Weight-Bearing Versus Partial Weight-Bearing After Plate Fixation of Distal Femur Fractures in Elderly Patients. A Randomized Controlled Trial. *Geriatr Orthop Surg Rehabil*. 2021;**12**: 21514593211055889. doi:10.1177/21514593211055889
44. Li J, Li J, Zhang QL. Effect of core muscle strengthening training on functional recovery and Barthel Index in hip replacement patients. *Jilin J Med*. 2022; **43**(03): 785-786. doi:10.3969/j.issn.1004-0412.2022.03.084

45. Yan YY, Zhang Y. Effect of progressive resistance exercise training on hip function after internal fixation of femoral intertrochanteric fractures in elderly patients. *Clin Res Prac.* 2022;**7**(20): 169-171.
46. Overgaard J A, Kallemose T, Mangione K K, et al. Six Versus 12 Weeks of Outpatient Physical Therapy Including Progressive Resistance Training in Cognitively Intact Older Adults After Hip Fracture: A Multicenter Randomized Controlled Trial. *J Gerontol A Biol Sci Med Sci.* 2022;**77**(7): 1455-1462. doi:10.1093/gerona/glab256