

Supplementary Figures for Regionally high risk increase for precipitation extreme events under global warming

Cristian Martinez-Villalobos and J. David Neelin

December 16, 2022




Figure S1: **a.** Multi-model mean of the absolute change in power law range exponent τ_P comparing 2075-2099 (τ_P^{GW}) and 1990-2014 (τ_P^{his}). **b.** **(c.)** Multi-model mean fractional change of the mean over wet days $\bar{P}_{wet}$ (fraction of wet days) comparing 2075-2099 vs 1990-2014. In **b** and **c** changes are normalized by each model global mean temperature change prior to aggregating. Dots show regions in which at least 80% of the models agree in the sign of the change.




Figure S2: **a.** Analytical risk ratios according to equation 13 in main text (crosses), and approximation for large $\frac{P}{P_L}$ (assuming no change in τ_P) (equation 3 in main text) corresponding to the same changes in parameters as in Figure 6 in main text (see caption). **b.** Risk ratio for different values of P_r as a function of $\gamma = \frac{\delta P_L}{P_L}$ according to approximation given in equation 3 in main text. In this example $\tau_P^{his} = \tau_P^{fut} = \tau_P = 0.5$, $P_L^{his} = P_L = 20mm$, and $P_L^{fut} = (1 + \gamma)P_L$.