

Table 2 - Clinicopathological features of the studied population *versus* ACOSOG Z0011

Characteristics	Studied population (N=82)	Z001 Trial (N=420)	p value
Age (yr)			
Median (min, max)	55 (31, 82)	56 (24, 92)	0.567
Missing	0	7	
Age (yr)			
<=50, N(%)	29 (35.4)	135 (32.1)	0.638
>50, N(%)	53 (64.6)	278 (66.2)	
Missing	0	7 (1.7)	
Clinical T stage, N(%)			
T1	62 (75.6)	284 (67.6)	0.039
T2	19 (23.2)	134 (31.9)	
T3	1 (1.2)	0	
Missing	0	2 (0.5)	
Receptor Status, N(%)			
ER+ and/or PgR+	75 (91.5)	320 (76.2)	0.069
ER-/PgR-	7 (8.5)	63 (15)	
Missing	0	37 (8.8)	
LVI, N(%)			
Yes	13 (15.9)	129 (30.7)	<0.001
No	69 (84.2)	189 (45)	
Missing	0	102 (24.3)	
Tumor type, N(%)			
Infiltrating ductal	66 (80.5)	344 (81.9)	0.921
Infiltrating lobular	5 (6.1)	27 (6.4)	
Other	10 (12.2)	45 (10.7)	
Missing	1 (1.2)	4 (1)	