



# Supplemental Figure 1




**Figure S1: Characterization of male mito-Keima mice fed western diet. A-B. Fat and lean mass. C-F. NAFLD criteria and composite histology scores. G-K. Plasma glucose, insulin, cholesterol, NEFA and AST levels. L-N. Liver mRNA levels for *Mcp1*, *Cd68* and *Scd1*. Data are the mean  $\pm$  s.e.m. for  $n = 5-10$  mice per group. Data were analyzed by 2-way ANOVA followed by multiple comparison testing. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ , \*\*\*\* $p < 0.0001$ .**

## Supplemental Figure 2




**Figure S2: Characterization of female mito-Keima mice fed western diet. A-D.** Body weight, change in body weight, and fat and lean mass. **E-F.** Liver triglyceride levels and relative rates of mitophagy. Data are the mean  $\pm$  s.e.m. for  $n = 5-8$  mice per group. Data were analyzed by 2-way ANOVA followed by multiple comparison testing. \* $p < 0.05$ , \*\*\* $p < 0.001$ .

### Supplemental Figure 3




**Figure S3: NAFLD progression in response to short-term dietary challenge is unaffected in female liver-specific PARKIN knockout mice.** **A.** Body weight and composition (fat and lean mass) for WT and LKO mice after six weeks WD feeding. **B.** Liver triglyceride levels expressed as mg triglyceride per g liver. **C.** Plasma triglyceride levels. **D.** Plasma cholesterol levels. **E.** Plasma ALT levels. **F.** Plasma AST levels. Data are the mean ± s.e.m. for n = 9-11 mice per group. Data were analyzed by Student's t-test. \*p<0.05

## Supplemental Figure 4




**Figure S4: NAFLD progression in response to short-term dietary challenge is unaffected in female liver-specific PARKIN knockout mice. A.** Representative images of H&E-stained liver sections from six-week WD fed WT and LKO mice at 10X, 20X and 60X. **B.** NAFLD activity score (NAS) consisting of steatosis (0-3), inflammation (0-3) and ballooning (0-2) grading and the composite (summed criteria) NAS. **C.** Liver gene expression measured by QPCR for noted gene markers of inflammation and fibrosis. Target gene expression was calculated relative to *Gapdh* and expressed as fold-change relative to WT. Data are the mean  $\pm$  s.e.m. for n = 9-11 mice per group. Data were analyzed by Student's t-test. \*p<0.05

## Supplemental Figure 5



**Figure S5: NAFLD progression in response to long-term dietary challenge is unaffected in female liver-specific PARKIN knockout mice.** **A.** Body weight and composition (fat and lean mass) for WT and LKO mice after 20 weeks WD feeding. **B.** Liver triglyceride levels expressed as mg triglyceride per g liver. **C.** Plasma triglyceride levels. **D.** Plasma cholesterol levels. **E.** Plasma ALT levels. **F.** Plasma AST levels. Data are the mean ± s.e.m. for n = 9-11 mice per group. Data were analyzed by Student's t-test. \*p<0.05

## Supplemental Figure 6



**Figure S6: NAFLD progression in response to long-term dietary challenge is unaffected in female liver-specific PARKIN knockout mice. A.** Representative images of H&E-stained liver sections from 20-week WD fed WT and LKO mice at 10X, 20X and 60X. **B.** NAFLD activity score (NAS) consisting of steatosis (0-3), inflammation (0-3) and ballooning (0-2) grading and the composite (summed criteria) NAS. **C.** Liver gene expression measured by QPCR for noted gene markers of inflammation and fibrosis. Target gene expression was calculated relative to *Gapdh* and expressed as fold-change relative to WT. Data are the mean  $\pm$  s.e.m. for n = 9-11 mice per group. Data were analyzed by Student's t-test. \*p<0.05