

Table1. Characteristics and outcomes of patients

	cLNU	OBCR	P value
No. of patients(n)	50	48	
Age(years) ($\bar{x}\pm s$)	67 $\pm$ 10.5(38-75)	66 $\pm$ 11.8(45-78)	0.93
Side(R/L)(n)	26/24	20/28	0.86
Gender(F/M)(n)	30/20	25/23	0.92
Median ASA score(n)			0.91
1/2/3	2/47/1	3/45/0	
Location of tumor(n)			0.89
Renal pelvis	43	38	
ureter	7	10	
operative time(min) ($\bar{x}\pm s$)	98.5 $\pm$ 40.3	132.4 $\pm$ 60.2	<0.01*
blood loss(ml) ($\bar{x}\pm s$)	60.4 $\pm$ 20.3	150.6 $\pm$ 50.2	<0.01*
Length of hospital stay(d) ($\bar{x}\pm s$)	5.3 $\pm$ 2.2	8.1 $\pm$ 2.3	<0.01*
Recovery of bowel function(d) ($\bar{x}\pm s$)	1.8 $\pm$ 0.8	2.1 $\pm$ 0.7	0.89
incision length(cm) ($\bar{x}\pm s$)	6.3 $\pm$ 1.2	11.5 $\pm$ 3.2	<0.01*
Pathological stage(n)			0.86
T1/T2/T3	6/23/1	5/18/1	
Tumor grade(n)			0.83
G1/G2/G3	5/22/3	8/18/4	
Follow-up(m) ($\bar{x}\pm s$)	27.5 $\pm$ 9.5	30.2 $\pm$ 8.2	0.92
Tumor recurrence(n)			
Intravesical tumor recurrence	3	2	0.32
Extra-bladder recurrence	3	1	0.22
ASA=American Society of Anesthesiologists, *Statistically significant			