

Supplementary Information


PDI augments kainic acid-induced seizure activity and neuronal death by inhibiting PP2A-GRIA2-PICK1-mediated AMPA receptor internalization in the mouse hippocampus

Duk-Shin Lee^{1,2}, Tae-Hyun Kim^{1,2}, Hana Park^{1,2} and Ji-Eun Kim^{1,2*}


¹ Department of Anatomy and Neurobiology, College of Medicine, Hallym University, Chuncheon 24252, South Korea

² Institute of Epilepsy Research, College of Medicine, Hallym University, Chuncheon 24252, South Korea


* Correspondence to: J. -E. Kim, Department of Anatomy and Neurobiology, College of Medicine, Hallym University, Chuncheon, Kangwon-Do 24252, South Korea; Tel: +82-33-248-2522; Fax: +82-33-248-2525; E-mail: jieunkim@hallym.ac.kr




Supplementary Figure 1. Full-length gel images of Western blots in Figure 1.




Supplementary Figure 2. Full-length gel images of Western blots in Figure 2.




Supplementary Figure 3. Full-length gel images of Western blots in Figure 3.




Supplementary Figure 4. Full-length gel images of Western blots in Figure 4.



Supplementary Figure 5. Full-length gel images of Western blots in Figure 6.



Supplementary Figure 6. Full-length gel images of Western blots in Figure 7.



Supplementary Figure 7. Full-length gel images of Western blots in Figure 8.