

Supplementary file

This supplementary file contains tables on adverse events occurrence within 1 hour (table 1) and 24 hours (table 2).

Supplementary table 1. Overview of adverse events occurrence within 1 hours after infusion.

	Percentage of patients			
	Never seen	less than 25%	25% or more	Total
Adverse events within 1h				
Allergic reaction	15 (29%)	30 (59%)	6 (12%)	51 (100%)
Arrhythmia	26 (51%)	24 (47%)	1 (2.0%)	51 (100%)
Dehydration	44 (86%)	7 (14%)	0 (0%)	51 (100%)
Diarrhea	25 (49%)	24 (47%)	2 (3.9%)	51 (100%)
Dyspnea	19 (37%)	29 (57%)	3 (5.9%)	51 (100%)
Fever	23 (45%)	21 (41%)	7 (14%)	51 (100%)
Fluid overload	26 (51%)	17 (33%)	8 (16%)	51 (100%)
Flushing	10 (20%)	25 (49%)	16 (31%)	51 (100%)
Haemoglobinuria	36 (71%)	10 (20%)	5 (9.8%)	51 (100%)
Haemolysis	41 (80%)	8 (16%)	2 (3.9%)	51 (100%)
Hypertension	19 (37%)	27 (53%)	5 (9.8%)	51 (100%)
Infection	40 (78%)	8 (16%)	3 (5.9%)	51 (100%)
Nausea	7 (14%)	27 (53%)	17 (33%)	51 (100%)
Renal dysfunction	32 (63%)	19 (37%)	0 (0%)	51 (100%)
Taste changes	13 (25%)	12 (24%)	26 (51%)	51 (100%)
Thoracic pressure	31 (61%)	19 (37%)	1 (2.0%)	51 (100%)
Vomiting	13 (25%)	38 (75%)	0 (0%)	51 (100%)
Total	420 (48%)	345 (40%)	102 (12%)	867 (100%)

Note: One respondent did not answer this question.

Supplementary table 2. Overview of adverse events occurrence within 24 hours after infusion.

	Percentage of patients			
	Never seen	less than 25%	25% or more	Total
Adverse events within 24h				
Allergic reaction	29 (57%)	20 (39%)	2 (3.9%)	51 (100%)
Arrhythmia	31 (61%)	19 (37%)	1 (2.0%)	51 (100%)
Dehydration	38 (75%)	13 (25%)	0 (0%)	51 (100%)
Diarrhea	13 (25%)	32 (63%)	6 (12%)	51 (100%)
Dyspnea	33 (65%)	17 (33%)	1 (2.0%)	51 (100%)
Fever	15 (29%)	31 (61%)	5 (9.8%)	51 (100%)
Fluid overload	19 (37%)	25 (49%)	7 (14%)	51 (100%)
Flushing	26 (51%)	23 (45%)	2 (3.9%)	51 (100%)
Haemoglobinuria	34 (67%)	13 (25%)	4 (7.8%)	51 (100%)
Haemolysis	43 (84%)	7 (14%)	1 (2.0%)	51 (100%)
Hypertension	22 (43%)	26 (51%)	3 (5.9%)	51 (100%)
Infection	29 (57%)	18 (35%)	4 (7.8%)	51 (100%)
Nausea	5 (9.8%)	28 (55%)	18 (35%)	51 (100%)
Renal dysfunction	26 (51%)	23 (45%)	2 (3.9%)	51 (100%)
Taste changes	12 (24%)	21 (41%)	18 (35%)	51 (100%)
Thoracic pressure	39 (76%)	12 (24%)	0 (0%)	51 (100%)
Vomiting	8 (16%)	35 (69%)	8 (16%)	51 (100%)
Total	422 (49%)	363 (42%)	82 (9.5%)	867 (100%)

Note: One respondent did not answer this question.