

Effects of maxillary expansion in mixed dentition with Invisalign First System and Hass: A prospective cohort study

Lanxin Lu^{1,2,3,+}, Lingling Zhang^{1,2,3,+}, Chengri Li^{1,2,3}, Fang Yi^{1,2,3}, Lei Lei^{1,2,3}, and Yanqin Lu^{1,2,3}

¹ Xiangya Stomatological Hospital & Xiangya School of Stomatology, Central South University, Changsha, 410008, Hunan, China

² Hunan Key Laboratory of Oral Health Research, China

³Hunan Clinical Research Center of Oral Major Diseases and Oral Health, China

⁺ Lanxin Lu and Lingling Zhang contributing equally to this article as co-first authors.

Corresponding Author:

Y. Lu, Professor, PhD.,

Department of Orthodontics, Hunan Key Laboratory of Oral Health Research & Hunan 3D Printing

Engineering Research Center of Oral Care & Hunan Clinical Research Center of Oral Major Diseases

and Oral Health & Xiangya Stomatological Hospital & Xiangya School of Stomatology, Central South

University, Changsha, 410008, Hunan, China.

Telephone: +86 13973102658

Email: 213031@csu.edu.cn

Journal name: *Clinical Oral Investigations*




Fig s1 Hass



Fig s2 Invisalign First System




Fig s3 Colour map generated from T0 model

Table s1 Baseline data of the two treated groups of patients after PSM

	Baseline data after PSM		<i>P</i> value
	First group	Hass group	
Sex			
male	8	8	1.00
female	9	9	
Age			
CS1	7	6	0.34
CS2	10	8	
CS3	0	3	
Complexity			
DI low	1	1	0.84
DI medium	3	5	
DI high	13	11	
Angle's classification			
Class I	5	4	1.00
Class II	12	13	