

Table 1: Epidemiological characteristics of COVID-19 patients. Data are shown as median (IQR) or n (%). ICU=intensive care unit. P values comparing ICU group and non-ICU group were obtained from χ^2 test, Fisher's exact test, or Mann-Whitney U test. * $P < 0.05$.

	No. (%)			
	ICU (n=45)	Non-ICU (n=156)	Total (n=201)	P Value
Age, median (IQR), y	57.0 (46.0-66.0)	40.0 (31.0-53.0)	45.0 (33.8-59.0)	<0.0001*
Sex				
Male	27 (60.0%)	75 (48.1%)	102 (50.7%)	0.178
Female	18 (40.0%)	81 (51.9%)	99 (49.3%)	
Occupation				
Office worker	18 (40.0%)	78 (50.0%)	96 (47.8%)	0.006*
Unemployed	3 (6.7%)	23 (14.7%)	26 (12.9%)	
Retired	16 (35.6%)	32 (20.5%)	48 (23.9%)	
Self-employed laborer	5 (11.1%)	8 (5.1%)	13 (6.5%)	
Peasant	3 (6.7%)	2 (1.3%)	5 (2.5%)	
Student	0 (0.0%)	13 (8.3%)	13 (6.5%)	
Exposure with Wuhan				
Yes	26 (57.8%)	66 (42.3%)	92 (45.8%)	0.048*
No	19 (42.2%)	90 (57.7%)	109 (54.2%)	
Duration of complaint	5.0 (3.0-8.0)	5.0 (3.0-8.0)	5.0 (3.0-8.0)	0.4904
Incubation	6.0 (4.0-9.0)	6.0 (3.0-7.3)	6.0 (3.8-8.0)	0.6978
Outcome				
Discharge	43 (26.7%)	156 (3.3%)	63 (31.3%)	0.049*
Death	2 (2.2%)	0 (0.0%)	1 (0.5%)	