

Table 1. Demographic and clinical characteristics of patients with obsessive-compulsive disorder (OCD) and healthy controls (HCs).

Characteristics	Adult OCD sample (n = 690)	Adult HC sample (n = 646)	Pediatric OCD sample (n = 175)	Pediatric HC sample (n = 142)
Demographic Characteristics				
Age (years)	31.6± 9.78	30.8± 9.97	14.5±2.3	14.3±2.4
Male N (%)	397 (25.6)	380 (24.2)	97 (27.8)	77 (22.1)
Clinical Characteristics				
OCD illness severity score	25± 7.11		20.8±8.0	
Childhood-onset N (%)	351 (51.7)			
Duration of illness	12.4± 11.1		3.0±2.5	
Medication use at time of scan N (%)	261 (37.8)		105 (60)	
Lifetime diagnosis				
Anxiety	76 (11.02)		48 (27.4)	
Major depression	84 (12.17)		18 (10.3)	
Current comorbid disorders				
Anxiety	69 (10.0)		29 (16.6)	
Major depression	77 (11.2)		6 (3.4)	
OCD symptom dimension				
Aggressive/checking	411 (59.6)		73 (41.7)	
Contamination/cleaning	355 (51.5)		62 (35.4)	
Symmetry/ordering	370 (53.6)		68 (38.9)	
Sexual/religious	228 (33.0)		55 (31.43)	
Hoarding	114 (16.5)		47 (26.9)	
Symptom score was indicated by total score on the adult and child version of the Yale-Brown Obsessive Compulsive Scales.				
OCD symptom dimensions were measured with the YBOCS symptom checklist.				