

Table 3. Participant sociodemographics and clinical characteristics.

	RESPONDENTS (CONJOINT ANALYSIS) (<i>n</i> = 67)
SEX	
Women	63%
Men	37%
AGE	
Mean, years old (SD)	50 (10.4)
18-44	42%
45-54	42%
≥ 55	16%
PREVIOUS INJECTABLE TREATMENTS	
Yes	61%
No	39%
Not sure	-
CONCOMITANT DISEASES	
Mean, <i>n</i> (SD)	3.3 (2.0)
Joint pain	64%
Cholesterol	39%
Pituitary hormone deficiency (TSH, testosterone, etc.)	39%
Diabetes or high blood sugar	31%
Hypertension	31%
Carpal tunnel syndrome	28%
Sleep apnoea	27%
Colorectal polyps	24%
Thyroid nodules (goitre)	22%
Heart diseases	15%
None	6%
YEARS SINCE DIAGNOSIS	
Mean, years (SD)	11.4 (8.0)
≤3	21%
4-8	30%
9-15	25%
≥15	24%
PRIOR SURGERY	

Yes	91%
<i>Complete resection, n</i>	61
Yes	12%
No	75%
Not sure	13%
No	9%
PRIOR RADIOTHERAPY TREATMENT	
Yes	36%
No	64%