

Supplementary Table 4

Results for minimum inhibitory concentrations and susceptibility for all isolates. The cell is left blank if no test was performed. N number of isolates tested, % percentage of tested samples that were susceptible, *no single mode value – both values given separated by semicolon (;)

Isolate	Amphotericin MIC [N, mode (range)] N, susceptible (%)	Econazole MIC [N, mode (range)] N, susceptible (%)	Itraconazole MIC [N, mode (range)] N, susceptible (%)	Natamycin MIC [N, mode (range)] N, susceptible (%)	Posaconazole MIC [N, mode (range)] N, susceptible (%)	Voriconazole MIC [N, mode (range)] N, susceptible (%)
<i>Acremonium</i> spp	8, 4 (1-4) 8, (13)	3, 2 (2) 5, (60)	1, 2 (2) 6, (0)	8, 4 (2-8) 8, (100)	1, 1 (1) 1, (100)	8, 1 (0.25-2) 8, (100)
<i>Acrophialophora</i> spp	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 4 (4) 1, (100)		1, 0.06 (0.06) 1, (100)
<i>Alternaria</i> spp	10, 0.5 (0.03-1) 10, (100)	3, 1 (1->64) 4, (50)	7, 0.5 (0.125-1) 7, (100)	10, 2 (0.5-4) 10, (100)	3, 0.25 (0.06-0.25) 3, (100)	10, 8 (0.25-16) 10, (50)
<i>Arthrinium phaeospermum</i>		1, 0.25 (0.25) 1, (100)		1, 0.25 (0.25) 1, (100)		1, 2 (2) 1, (100)
<i>Arthrographis kalrae</i>	1, 1 (1) 1, (100)		1, 1 (1) 1, (100)	1, 2 (2) 1, (100)	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 1, (100)
<i>Aspergillus clavatus</i>	1, 0.125 (0.125) 1, (100)	1, 32 (32) 1, (0)	1, >16 (>16) 1, (0)	1, 4 (4) 1, (100)		1, 2 (2) 1, (100)
<i>Aspergillus flavus</i> complex	14, 1 (0.25-2) 14, (93)	7, 1 (0.5-2) 7, (100)	11, 0.125 (0.06-0.5) 11, (100)	14, 32 (16-64) 14, (7)	7, 0.125 (0.06-0.25) 7, (100)	14, 0.5 (0.25-1) 14, 100
<i>Aspergillus fumigatus</i> complex	33, 0.25 (0.125-1) 33, (100)	16, 2 (1-32) 16, (94)	25, 0.25 (0.06-1) 25, (96)	30, 4 (1-8) 30, (100)	9, 0.06 (0.06-0.25) 9, (100)	33, 0.5 (0.25-2) 33, (100)
<i>Aspergillus nidulans</i> complex	1, 1 (1) 1, (100)	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 4 (4) 1, (100)		1, 0.25 (0.25) 1, (100)
<i>Aspergillus niger</i> complex	6, 0.06 (0.06-0.25) 6, (100)	1, 1 (1) 1, (100)	4, 0.5 (0.5) 4, (100)	5, 4 (2-4) 5, (100)	1, 0.125 (0.125) 1, (100)	6, 0.5,1 (0.5-1) 6, (100)

Isolate	Amphotericin MIC [N, mode (range)] N, susceptible (%)	Econazole MIC [N, mode (range)] N, susceptible (%)	Itraconazole MIC [N, mode (range)] N, susceptible (%)	Natamycin MIC [N, mode (range)] N, susceptible (%)	Posaconazole MIC [N, mode (range)] N, susceptible (%)	Voriconazole MIC [N, mode (range)] N, susceptible (%)
<i>Aspergillus tamarii</i>	1, 0.25 (0.25) 1, (100)	1, 0.5 (0.5) 1, (100)	1, 0.125 (0.125) 1, (100)	1, 64 (64) 1, (0)		1, 0.25 (0.25) 1, (100)
<i>Aspergillus versicolor</i> complex	2, 0.5;2 (0.5-2)* 2, (50)	2, 0.25;2 (0.25-2)* 2, (100)	2, 0.125;0.5 (0.125-0.5)* 2, (100)	2, 2;4, (2-4)* 2, (100)		2, 0.25;0.5 (0.25-0.5)* 2, (100)
<i>Aureobasidium pullulans</i>	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 0.125 (0.125) 1, (100)	1, 4 (4) 1, (100)		1, 0.25 (0.25) 1, (100)
Basidiomycete	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 4 (4) 1, (100)		1, 0.25 (0.25) 1, (100)
<i>Beauveria</i> spp	2, 2,4 (2-4) 2, (0)	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 2, (50)	2, 4;8 (4-8)* 2, (100)	1, 0.5 (0.5) 1, (100)	2, 0.25,1 (0.25-1) 2, (100)
<i>Bipolaris hawaiiensis</i>	1, 1 (1) 1, (100)			1, 2 (2) 1, (100)		1, 1 (1) 1, (100)
<i>Bjerkandera adusta</i>	2, 0.125 (0.125) 2, (100)	1, 0.25 (0.25) 1, (100)	2, 0.25;0.5 (0.25-0.5)* 2, (100)	2, 1,2 (1-2) 2, (100)	1, 2 (2) 1, (0)	2, <0.03;0.06 (<0.03-0.06) 2, (100)
<i>Candida albicans</i>	26, 0.5 (0.25-1) 26, (100)	13, <0.125 (<0.125-0.25) 13, (100)	2, <0.03 (<0.03) 2, (100)	22, 4 (2-32) 22, (95)	1, <0.03 (<0.03) 1, (100)	24, <0.03 (<0.03) 24, (100)
<i>Candida dubliniensis</i>	1, 0.25 (0.25) 1, (100)			1, 8 (8) 1, (100)		1, <0.03 (<0.03) 1, (100)
<i>Candida haemulonii</i>	4, 1 (0.5->16) 4, (75)	3, 0.5 (0.5-2) 3, (100)	1, 0.25 (0.25) 1, (100)			3, 0.06 (<0.03-0.25) 3, (100)
<i>Candida orthopsilosis</i>	3, 0.5 (0.5) 3, (100)	3, 2 (0.25-2) 3, (100)		4, 4 (4) 4, (100)		3, <0.03 (<0.03-0.125) 3, (100)
<i>Candida palmiophila</i>	1, 0.25 (0.25) 1, (100)			1, 2 (2) 1, (100)		1, 0.125 (0.125) 1, (100)

Isolate	Amphotericin MIC [N, mode (range)] N, susceptible (%)	Econazole MIC [N, mode (range)] N, susceptible (%)	Itraconazole MIC [N, mode (range)] N, susceptible (%)	Natamycin MIC [N, mode (range)] N, susceptible (%)	Posaconazole MIC [N, mode (range)] N, susceptible (%)	Voriconazole MIC [N, mode (range)] N, susceptible (%)
Candida parapsilosis	54, 0.5 (0.25-1) 54, (100)	36, 1 (<0.125-4) 36, (100)	8, <0.03 (<0.03-0.06) 8, (100)	32, 4 (4-16) 32, (100)	4, <0.03 (<0.03-0.125) 4, (100)	56, <0.03 (<0.03-0.06) 56, (100)
Candida tropicalis	1, 1 (1) 1, (100)	1, 1 (1) 1, (100)		1, 2 (2) 1, (100)		1, 0.06 (0.06) 1, (100)
Cephalosporium spp	1, 1 (1) 1, (100)	1, 0.125 (0.125) 1, (100)	1, 0.5 (0.5) 1, (100)	1, 4 (4) 1, (100)	1, 0.06 (0.06) 1, (100)	1, 0.125 (0.125) 1, (100)
Ceriporia lacerata	1, 0.5 (0.5) 1, (100)		1, 0.25 (0.25) 1, (100)	1, 2 (2) 1, (100)	1, 0.06 (0.06) 1, (100)	1, 0.5 (0.5) 1, (100)
Chaetomium spp	2, 0.5,1 (0.5-1) 2, (100)	2, 0.25,1 (0.25-1) 2, (100)	2, 0.5 (0.5) 2, (100)	2, 2 (2) 2, (100)		2, 0.5 (0.5) 2, (100)
Cladophialophora boppii	1, 1 (1) 1, (100)			1, 4 (4) 1, (100)		1, 0.5 (0.5) 1, (100)
Cladosporium sphaerospermum	1, 2 (2) 1, (0)	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 2 (2) 1, (100)		1, 0.25 (0.25) 1, (100)
Cladosporium spp	1, 1 (1) 1, (100)					1, 1 (1) 1, (100)
Clavispora lusitaniae (Candida lusitaniae)	2, 0.25;0.5 (0.25-0.5) 2, (100)	2, <0.125 (<0.125) 2, (100)		1, 4 (4) 1, (100)		2, <0.03 (<0.03-1) 2, (100)
Clonostachys rosea	1, 1 (1) 1, (100)	1, 8 (8) 1, (0)	1, 2 (2) 1, (0)	1, 4 (4) 1, (100)		1, 8 (8) 1, (0)
Corynascus sepedonium	1, 0.125 (0.125) 1, (100)	1, 1 (1) 1, (100)		1, 2 (2) 1, (100)		1, 0.25 (0.25) 1, (100)
Cryptococcus diffluens	3, 0.5 (0.5) 3, (100)	1, 2 (2) 1, (100)	1, 0.5 (0.5) 1, (100)	1, 4 (4) 1, (100)	1, 0.5 (0.5) 1, (0)	3, 0.25 (0.125-0.25) 3, (100)

Isolate	Amphotericin MIC [N, mode (range)] N, susceptible (%)	Econazole MIC [N, mode (range)] N, susceptible (%)	Itraconazole MIC [N, mode (range)] N, susceptible (%)	Natamycin MIC [N, mode (range)] N, susceptible (%)	Posaconazole MIC [N, mode (range)] N, susceptible (%)	Voriconazole MIC [N, mode (range)] N, susceptible (%)
Cryptococcus uniguttulatus	1, 2 (2) 1, (0)	1, 4 (4) 1, (100)		1, 4 (4) 1, (100)		1, 1 (1) 1, (100)
Curvularia spp	2, 0.25,0.5 (0.25-0.5) 2, (100)	1, 2 (2) 1, (100)	2, 0.25 (0.25) 2, (100)	2, 1,4 (1-4) 2, (100)	1, 0.125 (0.125) 1, (100)	2, 0.25,1 (0.25-1) 2, (100)
Cutaneotrichosporon curvatus	2, 0.25,0.5 (0.25-0.5) 2, (100)	2, 0.5 (0.5) 2, (100)		1, 2 (2) 1, (100)		2, <0.03;0.125 (<0.03-0.125)* 2, (100)
Cutaneotrichosporon cutaneum	1, 0.5 (0.5) 1, (100)	1, 1 (1) 1, (100)		1, 4 (4) 1, (100)		1, 0.06 (0.06) 1, (100)
Debaryomyces hansenii (Candida famata)	2, 1 (1) 2, (100)	2, 0.25;2 (0.25-2)* 2, (100)	1, 0.25 (0.25) 1, (100)	2, 0.5,1 (0.5-1) 2, (100)		2, <0.03;0.06 (<0.03-0.06)* 2, (100)
Exophiala oligosperma	1, 0.5 (0.5) 1, (100)	1, 1 (1) 1, (100)	1, 0.065 (0.065) 1, (100)	1, 2 (2) 1, (100)		1, 0.125 (0.125) 1, (100)
Fusarium dimerum	1, 1 (1) 1, (100)			1, 2 (2) 1, (100)	1, >16 (>16) 1, (0)	1, 8 (8) 1, (0)
Fusarium dimerum complex	7, 1 (0.25-2) 7, (86)	2, >64 (>64) 4, (0)	3, >16 (>16) 6, (0)	7, 2 (2-4) 7, (100)	1, >16 (>16) 2, (0)	7, 8 (1-16) 7, (14)
Fusarium moniliforme complex	1, 1 (1) 1, (100)			1, 4 (4) 1, (100)		
Fusarium oxysporum complex	12, 1 (1-2) 13, (62)	8, 2 (2-64) 10, (40)	3, >16 (>16) 12, (0)	13, 8 (1-8) 13, (100)		13, 4 (2-16) 13, (92.3)
Fusarium solani complex	11, 1 (0.25-2) 11, (82)	5, 4 (2->64) 7, (43)	5, >16 (16->16) 10, (0)	10, 4 (4-8) 10, (100)	2, >16 (>16) 3, (0)	10, 4 (2-4) 10, (50)

Isolate	Amphotericin MIC [N, mode (range)] N, susceptible (%)	Econazole MIC [N, mode (range)] N, susceptible (%)	Itraconazole MIC [N, mode (range)] N, susceptible (%)	Natamycin MIC [N, mode (range)] N, susceptible (%)	Posaconazole MIC [N, mode (range)] N, susceptible (%)	Voriconazole MIC [N, mode (range)] N, susceptible (%)
Fusarium spp	180, 1 (0.06-4) 180, (66)	92, 4 (0.25->64) 130, (40)	22, >16 (0.5->16) 151, (1)	178, 4 (0.5-16) 178, (100)	22, >16 (1->16) 43, (9)	175, 4 (0.5->16) 179, (60.3)
Gibberella fujikuroi	1, 1 (1) 1, (100)			1, 4 (4) 1, (100)		1, 4 (4) 1, (100)
Irpex lacteus	1, 8 (8) 1, (0)		1, 0.5 (0.5) 1, (100)	1, 1 (1) 1, (100)	1, 0.125 (0.125) 1, (100)	1, 0.5 (0.5) 1, (100)
Lomentospora prolificans	2, 8;>16 (8->16)* 2, (0)	2, >64, (>64) 2, (0)	2, >16, (>160) 2, (0)	2, 4 (4) 2, (100)		2, 16;>16 ((16->16))* 2, (0)
Metarrhizium spp	1, 2 (2) 1, (0)	1, 2 (2) 1, (100)	1, (0)	1, 8 (8) 1, (100)		1, 1 (1) 1, (100)
Meyerozyma guilliermondii (Candida guilliermondii)	29, 0.25 (0.125-1) 29, (100)	19, 2 (0.25-8) 19, (95)	5, 0.25 (0.25-1) 5, (80)	15, 4 (1-4) 15, (100)	5, 0.125 (0.06-0.5) 5, (80)	28, 0.125 (<0.03-0.5) 28, (100)
Naganishia diffluens	1, 1 (1) 1, (100)	1, 1 (1) 1, (100)		1, 2 (2) 1, (100)		1, 0.25 (0.25) 1, (100)
Nakaseomyces glabrata (candida glabrata)	2, 0.5.1 (0.5-1) 2, (100)	1, 0.25 (0.25) 1, (100)	1, 0.5 (0.5) 1, (100)	2, 4 (4) 2, (100)		1, 0.25 (0.25) 1, (100)
Neosartorya hiratsukae	3, 0.25 (0.25-0.5) 3, (100)	1, 0.5 (0.5) 1, (100)	1, 0.125 (0.125) 1, (100)	3, 4 (4-8) 3, (100)		3, 0.5 (0.5-4) 3, (100)
Ochroconis tshawytschae	1, 0.5 (0.5) 1, (100)			1, 1 (1) 1, (100)	1, 0.125 (0.125) 1, (100)	1, 1 (1) 1, (100)
Penicillium chrysogenum	2, 0.25;0.5 (0.25-0.5)* 2, (100)	2, 0.5 (0.5) 2, (100)	2, 0.25;0.5 (0.25-0.25)* 2, (100)	2, 2,4 (2-4)* 2, (100)		2, 1;2 (1-2)* 2, (100)
Penicillium spp	4, 0.25 (0.25-2) 4, (75)	1, 1 (1) 2, (100)	2, 0.06;0.125 (0.06-0.125)* 3, (67)	4, 1 (0.5-32) 4, (75)	1, 0.03 (0.03) 2, (50)	4, 0.25 (0.25-16) 4, (75)

Isolate	Amphotericin MIC [N, mode (range)] N, susceptible (%)	Econazole MIC [N, mode (range)] N, susceptible (%)	Itraconazole MIC [N, mode (range)] N, susceptible (%)	Natamycin MIC [N, mode (range)] N, susceptible (%)	Posaconazole MIC [N, mode (range)] N, susceptible (%)	Voriconazole MIC [N, mode (range)] N, susceptible (%)
Peniphora lycii	1, 0.5 (0.5) 1, (100)	1, 0.5 (0.5) 1, (100)	1, 0.125 (0.125) 1, (100)	1, 2 (2) 1, (100)		1, 8 (8) 1, (0)
Phaeoacremonium kraidenii	1, 0.25 (0.25) 1, (100)			1, 2 (2) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 0.25 (0.25) 1, (100)
Phanerochaete sordida	1, 0.5 (0.5) 1, (100)	1, 1 (1) 1, (100)	1, 0.5 (0.5) 1, (100)	1, 2 (2) 1, (100)	1, 0.25 (0.25) 1, (100)	1, 0.25 (0.25) 1, (100)
Phoma spp	6, 1 (0.03-2) 6, (83)	3, 0.5 (0.5) 3, (100)	6, 0.25 (0.25-1) 6, (100)	6, 2 (0.25-2) 6, (100)	3, 0.125 (<0.03-1) 3, (100)	5, 0.25 (<0.03-1) 5, (100)
Purpureocillium lilacinum	2, 8,16 (8-16) 7, (0)	4, 0.5,1 (0.5-1) 3, (100)	3, 1 (1-2) 4, (50)	1, 8 (8) 7, (14)	1, 0.25 (0.25) 1, (100)	7, 0.25 (0.25-0.5) 7, (100)
Purpureocillium spp	1, 2 (2) 1, (0)		1, 0.5 (0.5) 1, (100)			1, 0.25 (0.25) 1, (100)
Pyrenochaeta unguis-hominis	1, 0.5 (0.5) 1, (100)					1, >16 (>16) 1, (0)
Rhodotorula mucilaginosa	5, 0.5 (0.25-0.5) 5, (100)	4, 4 (0.5-4) 4, (100)		2, 4 (4) 2, (100)		5, 4 (0.5-16) 5, (60)
Saccharomyces cerevisiae				1, 4 (4) 1, (100)		
Sarocladium kiliense	6, 2 (2-4) 6, (0)	4, >64 (0.25->64) 6, (17)	1, >16 (>16) 3, (0)	6, 2 (2-4) 6, (100)		6, 1 (0.125-2) 6, (100)
Sarocladium strictum	2, 1,4 (1-4) 2, 50	2, (0)	2, (0)	2, 2;4 (2-4)* 2, (100)		2, 1;2 (1-2)* 2, (100)
Scedosporium apiospermum complex	5, 1;2 (1-4)* 5, (40)	2, 1;2 (1-2)* 2, (100)	3, 4 (0.5-4) 3, (33)	4, 4 (2-8) 4, (100)		5, 0.5 (0.5-1) 5, (100)

Isolate	Amphotericin MIC [N, mode (range)] N, susceptible (%)	Econazole MIC [N, mode (range)] N, susceptible (%)	Itraconazole MIC [N, mode (range)] N, susceptible (%)	Natamycin MIC [N, mode (range)] N, susceptible (%)	Posaconazole MIC [N, mode (range)] N, susceptible (%)	Voriconazole MIC [N, mode (range)] N, susceptible (%)
<i>Scedosporium de hooghii</i>	2, 2 (2) 2, (0)	1, 0.25 (0.25) 1, (100)	2, 0.5 (0.5) 2, (100)	2, 2 (2) 2, (100)		2, 0.25 (0.25) 2, (100)
<i>Tilletiopsis</i> spp	2, 0.125;2 (0.125-2)* 2, (50)	1, 2 (2) 2, (50)	1, 0.25 (0.25) 2, (50)	2, 4 (4) 2, (100)		2, 0.06;0.5 (0.06-0.5)* 2, (100)
<i>Trichosporon asahii</i>	1, 2 (2) 1, (0)					1, <0.03 (<0.03) 1, (100)
<i>Trichosporon jirovecii</i>	1, 0.5 (0.5) 1, (100)	1, (100)		1, 4 (4) 1, (100)		1, <0.03 (<0.03) 1, (100)
<i>Trichosporon lactis</i>	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 1, (100)				1, 0.03 (0.03) 1, (100)
Unidentified mould	2, 0.06;0.125 (0.06-0.125)* 2, (100)	1, 0.5 (0.5) 1, (100)	1, 0.25 (0.25) 1, (100)	2, 4 (4) 2, (100)		2, 0.125;0.5 (0.125-0.5)* 2, (100)
<i>Verticillium</i> spp	1, 2 (2) 1, (0)	1, 2 (2) 1, (100)	1, >16 (>16) 1, (0)	1, 4 (4) 1, (100)		1, 2 (2) 1, (100)
<i>Wickerhamomyces anomalus</i>	5, 0.25 (0.25) 5, (100)	3, 1 (1-2) 3, (100)	1, 0.5 (0.5) 1, (100)	4, 2,4 (2-4) 4, (100)	1, 0.5 (0.5) 1, (0)	5, 0.125 (0.125-0.25) 5, (100)
<i>Yarrowia lipolytica</i> (<i>Candida lipolytica</i>)	6, 1 (0.5-4) 6, (83)	3, 0.5 (0.25-2) 3, (100)	2, 0.25;0.5 (0.25-0.5)* 2, 100	4, 2 (2-8) 4, (100)	2, 0.25 (0.25) 2, (0)	6, 0.06 (<0.03-0.125) 6, (100)