



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2023 – 08:29 PM JST

PDB ID : 8I80  
Title : Crystal structure of Cph001-D189N  
Deposited on : 2023-02-02  
Resolution : 2.30 Å (reported)

**This wwPDB validation report is for manuscript review**

This is a Full wwPDB X-ray Structure Validation Report.

This report is produced by the wwPDB biocuration pipeline after annotation of the structure.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.32.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.32.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	<div> <div>6%</div> <div>87%</div> <div>9%</div> <div>.</div> </div>
1	B	286	<div> <div>3%</div> <div>91%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	B	501	-	-	X	-
3	ACT	B	502	-	-	X	-

For Manuscript Review

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4452 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Viomycin kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	285	Total	C	N	O	S	0	0	0
			2127	1326	389	404	8			
1	A	278	Total	C	N	O	S	0	0	0
			2082	1299	382	392	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	189	ASN	ASP	engineered mutation	UNP D2B3F1
A	189	ASN	ASP	engineered mutation	UNP D2B3F1

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			7	4	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			7	4	3		
2	B	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		
2	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	103	Total 103	O 103	0	0
4	A	85	Total 85	O 85	0	0

For Manuscript Review


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Viomycin kinase



- Molecule 1: Viomycin kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.30Å 86.94Å 87.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.28 – 2.30 28.26 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.1 (28.28-2.30) 97.2 (28.26-2.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.33 (at 2.29Å)	Xtriage
Refinement program	REFMAC 8.0.004	Depositor
R, $R_{free}$	0.194 , 0.245 0.201 , 0.247	Depositor DCC
$R_{free}$ test set	1390 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.5	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 43.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4452	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/2115	0.57	0/2873
1	B	0.32	0/2161	0.62	0/2939
All	All	0.31	0/4276	0.60	0/5812

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2082	0	2095	18	0
1	B	2127	0	2133	21	0
2	A	14	0	20	0	0
2	B	21	0	30	8	0
3	A	8	0	6	0	0
3	B	12	0	9	3	0
4	A	85	0	0	0	0
4	B	103	0	0	1	0
All	All	4452	0	4293	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:77:CYS:HB3	2:B:501:PEG:C1	2.03	0.89
1:B:77:CYS:HB3	2:B:501:PEG:H12	1.56	0.86
1:B:56:ARG:HD3	3:B:502:ACT:H1	1.69	0.73
1:B:16:PRO:HB2	1:A:203:LEU:HD12	1.69	0.73
1:B:77:CYS:HB3	2:B:501:PEG:H11	1.74	0.68
1:A:56:ARG:HH11	1:A:56:ARG:HG3	1.61	0.66
1:B:38:ASP:OD1	1:A:205:ARG:NH1	2.31	0.64
1:A:56:ARG:HH11	1:A:56:ARG:CG	2.12	0.61
1:B:56:ARG:CD	3:B:502:ACT:H1	2.39	0.52
1:B:249:ARG:HG2	4:B:651:HOH:O	2.10	0.52
1:A:54:PRO:HA	1:A:78:GLU:OE2	2.10	0.51
1:B:75:PRO:HA	1:B:92:LEU:HD23	1.93	0.51
1:A:145:TRP:HB3	1:A:175:LEU:HG	1.93	0.51
1:A:237:ASP:O	1:A:241:THR:HG23	2.12	0.50
1:B:90:LEU:HD12	2:B:501:PEG:H12	1.94	0.49
1:B:50:ALA:HB1	1:B:86:GLU:HG2	1.94	0.48
1:A:105:LEU:HD13	1:A:235:PHE:CD1	2.49	0.47
1:B:89:PHE:O	2:B:501:PEG:H21	2.14	0.47
1:A:52:ARG:NH1	1:A:55:ARG:HH12	2.12	0.47
1:B:167:GLN:O	1:B:170:ARG:HG2	2.14	0.46
1:B:28:GLY:O	1:B:45:ARG:NH1	2.48	0.46
1:B:31:HIS:HE1	1:B:52:ARG:HH12	1.63	0.46
1:B:53:LEU:N	1:B:54:PRO:CD	2.79	0.46
1:B:211:ASP:H	2:B:506:PEG:H21	1.82	0.45
1:A:129:GLY:HA3	1:A:183:GLU:O	2.17	0.44
1:A:48:ALA:O	1:A:52:ARG:HD3	2.17	0.44
1:A:160:MET:HE1	1:A:165:CYS:HA	1.99	0.44
1:B:56:ARG:HB2	3:B:502:ACT:H1	2.00	0.43
1:A:56:ARG:O	1:A:60:MET:HG2	2.18	0.43
1:A:4:LEU:HD13	1:A:20:PRO:O	2.18	0.43
1:B:77:CYS:CB	2:B:501:PEG:H11	2.46	0.42
1:A:160:MET:HE3	1:A:285:TYR:CD1	2.55	0.42
1:B:89:PHE:CD1	2:B:501:PEG:H22	2.53	0.42
1:A:148:PHE:O	1:A:152:VAL:HG23	2.20	0.42
1:B:26:ARG:HB2	1:B:33:VAL:HG22	2.02	0.41
1:A:53:LEU:N	1:A:54:PRO:CD	2.84	0.41
1:A:114:VAL:HG13	1:A:197:TRP:CH2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/286 (96%)	268 (98%)	6 (2%)	0	100	100
1	B	283/286 (99%)	279 (99%)	4 (1%)	0	100	100
All	All	557/572 (97%)	547 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/216 (98%)	209 (99%)	2 (1%)	78	89
1	B	215/216 (100%)	213 (99%)	2 (1%)	78	89
All	All	426/432 (99%)	422 (99%)	4 (1%)	78	89

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	3	ILE
1	B	127	SER
1	A	56	ARG
1	A	199	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	29	GLN
1	B	31	HIS
1	B	32	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	B	504	-	3,3,3	1.06	0	3,3,3	0.69	0
2	PEG	B	501	-	6,6,6	0.28	0	5,5,5	0.34	0
3	ACT	A	304	-	3,3,3	1.18	0	3,3,3	0.67	0
3	ACT	B	502	-	3,3,3	0.91	0	3,3,3	1.00	0
2	PEG	B	505	-	6,6,6	0.23	0	5,5,5	0.14	0
3	ACT	B	503	-	3,3,3	1.15	0	3,3,3	0.58	0
2	PEG	B	506	-	6,6,6	0.23	0	5,5,5	0.20	0
3	ACT	A	303	-	3,3,3	0.94	0	3,3,3	0.90	0
2	PEG	A	302	-	6,6,6	0.18	0	5,5,5	0.10	0
2	PEG	A	301	-	6,6,6	0.18	0	5,5,5	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	B	501	-	-	2/4/4/4	-
2	PEG	B	505	-	-	3/4/4/4	-
2	PEG	B	506	-	-	1/4/4/4	-
2	PEG	A	302	-	-	2/4/4/4	-
2	PEG	A	301	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	PEG	O1-C1-C2-O2
2	A	302	PEG	O1-C1-C2-O2
2	B	505	PEG	C1-C2-O2-C3
2	B	501	PEG	O1-C1-C2-O2
2	B	505	PEG	O1-C1-C2-O2
2	B	505	PEG	O2-C3-C4-O4
2	A	302	PEG	O2-C3-C4-O4
2	A	301	PEG	C4-C3-O2-C2
2	B	501	PEG	C4-C3-O2-C2
2	B	506	PEG	O2-C3-C4-O4

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	PEG	7	0
3	B	502	ACT	3	0
2	B	506	PEG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

For Manuscript Review

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	278/286 (97%)	0.41	18 (6%) 18 24	19, 38, 69, 94	0
1	B	285/286 (99%)	0.04	8 (2%) 53 60	15, 28, 56, 78	0
All	All	563/572 (98%)	0.22	26 (4%) 32 39	15, 33, 64, 94	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	PHE	6.6
1	A	29	GLN	4.4
1	B	21	GLU	4.2
1	A	46	THR	3.6
1	A	52	ARG	3.6
1	A	47	ALA	3.5
1	A	51	ALA	3.2
1	A	248	ARG	3.1
1	A	272	ASP	3.1
1	B	3	ILE	3.0
1	B	200	ASP	3.0
1	A	200	ASP	2.8
1	B	5	GLN	2.8
1	B	106	GLU	2.7
1	A	166	ARG	2.7
1	A	48	ALA	2.7
1	B	142	GLN	2.6
1	A	249	ARG	2.5
1	A	70	CYS	2.4
1	A	176	ASP	2.4
1	A	273	GLY	2.3
1	B	162	ASP	2.3
1	A	49	ALA	2.1
1	B	22	GLY	2.1

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	109	LYS	2.0
1	A	132	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ACT	A	303	4/4	0.70	0.19	45,47,49,54	0
3	ACT	B	503	4/4	0.73	0.30	56,57,57,58	0
2	PEG	B	505	7/7	0.76	0.35	46,48,57,58	0
2	PEG	A	302	7/7	0.76	0.28	64,70,73,74	0
2	PEG	B	501	7/7	0.78	0.23	33,40,47,50	0
3	ACT	A	304	4/4	0.80	0.16	44,44,44,48	0
3	ACT	B	502	4/4	0.86	0.17	39,39,40,42	0
2	PEG	A	301	7/7	0.87	0.23	42,50,53,54	0
3	ACT	B	504	4/4	0.90	0.11	53,55,55,56	0
2	PEG	B	506	7/7	0.91	0.18	25,37,51,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.