

Tables

Table 1 Clinical characteristics of subjects

Characteristics	
n	48
Men (n (%))	31 (64.6)
Age (years)	69.6±14.1
Height (cm)	160.3±10.1
Weight (kg)	60.8±12.5
NYHA Class (n (%))	
I	3(6.3)
II	42(87.5)
III	3(6.3)
NTproBNP(pg/ml)	1028.5
median	[20 - 5,110]
[minimum - maximum]	
Echocardiography	
Ejection fraction (%) (n=42)	55.9±15.1
Estimated end systolic volume	119.2±49.7
(ml) (n=28)	
Estimated end diastolic volume	65.3±36.2
(ml)(n=28)	

Values are mean ± SD, NYHA: New York Heart Association, NTproBNP: Nterminal pro-brain natriuretic peptide

Table 2 subgroup analysis

Age	n (%)	ICC	95% CI	SEM
≥ 75 years old	19(39.6)	0.85	0.72-0.94	2.564
< 75 years old	29(60.4)	0.84	0.73-0.92	2.8
Cardiac index				
≥ 2.2L/min/m ²	10(35.7)	0.84	0.61-0.96	2.548
< 2.2 L/min/m ²	18(64.3)	0.81	0.65-0.92	3.232
NYHA				
I - II	45(93.8)	0.82	0.73-0.89	2.662
III	3(6.3)	0.94	0.74-1.00	3.167

ICC: intraclass correlation coefficient, 95%CI: 95%

Confidence Interval, SEM: standard error of the

measurement, CI: cardiac index, NYHA: New York Heart

Association

Table 3 Number of tests required to obtain a clear LFCT (n=46)

	n (%)
by 1st measurement	35 (76.1)
by 2nd measurement	41 (89.1)

LFCT: lung to finger circulation time