

Livestock movement informs the risk of disease spread in traditional production systems in East Africa

Divine Ekwem¹, Thomas A. Morrison^{1*}, Richard Reeve¹, Jessica Enright², Joram Buza³, Gabriel Shirima³, James K. Mwajombe⁴, Tiziana Lembo^{1**}, J. Grant C. Hopcraft^{1**}

*Corresponding author

**These authors contributed equally

¹Boyd Orr Centre for Population and Ecosystem Health, Institute of Biodiversity, Animal Health & Comparative Medicine, College of Medical, Veterinary and Life Sciences, University of Glasgow, Glasgow, United Kingdom

²School of Computing Science, University of Glasgow, Glasgow, United Kingdom

³Nelson Mandela African Institution of Science and Technology, Arusha, Tanzania

⁴Department of Livestock, Serengeti District Council, Ministry of Livestock and Fisheries, United Republic of Tanzania

Supplementary Materials

Supplementary Methods.

Herd selection for Global Positioning System collaring study of cattle

We selected herds using a four-stage selection process:

1. Selection of resource areas: Six replicate grazing areas were randomly selected from a list of the largest 18 shared grazing areas in Serengeti District. All selected grazing areas were <1 km from at least one major watering point (Fig. 1, main text).
2. Selection of villages: For each selected grazing area, all villages that grazed in the area (n = 26) were included in the study.
3. Selection of herds: In each village, one large and one small herd were selected randomly for Global Positioning System (GPS) collar deployment from the list of all bomas, on condition that each boma was >1 km apart. The number of cattle in each herd was classified by size based on community perceptions as follows: large herds (≥ 90 cattle) and small herds (<90). Within villages, pairwise distances between bomas ranged from 1-5 km.
4. Selection of cattle for collaring: One healthy cow per herd was considered sufficient to track collective group movements¹.

GPS data retrieval and processing

Each sampled boma with collared cattle was visited every four weeks to retrieve data from the GPS devices. Owners were asked weekly if collared animals had remained with the herd to ensure the GPS data were representative of the herd's movement. Occasionally, collared cattle were sold, in which case another collar was deployed on a comparable individual within the same herd.

Collar data were pre-processed by excluding unrealistically high movements speeds (>20 km/hour)^{2,3}. These errors occurred infrequently and likely related to GPS satellite coverage, or possibly transport of collars within a moving vehicle.

Supplementary Figures



Figure S1. Pictures showing the components of Global Positioning System collars used in the study to track movements of individual cattle in a herd. Complete units were mounted on cattle using locally-produced collars.




Figure S2. A schematic used to define contact of two herds A and B (black and black and white cattle, respectively). The blue circle is the spatial scale, which in this case has a buffer size of 200 meters in diameter (double arrow). The buffer size indicates the proximity of a contact, which in this case was defined as two GPS fixes from two different herds with location less than 200 metres apart within one hour. The figure also shows that, although the distance between the two collared cattle in their herds was 200 meters, other cattle from both herds were likely to be in the path (less than 200 m) of collared cattle A and B.




Figure S3. Distribution of the overall livestock herd mobility across herd sizes during the study period: (a) total distance moved daily and (b) maximum distance displaced from the home boma. Boxplots represent median and interquartile range (IQR). Herds were classified as: large (>90), medium (30-90) and small (<30).




Figure S4. The proportion of contacts that occurred for each hour in the day. The spatial window (contact distance) was set at 50m, 200m and 500m. The temporal window (duration of contact) ranged from one hour to one week.




Figure S5. Pairwise contact rates across all spatiotemporal definitions of contacts decline with distance (km) between pairs of cattle's bomas. The contact rate between a pair of collared cattle was measured as the proportion of time they were in contact given a contact definition. The black fitted line is the regression line. The dashed reference line shows the distance beyond which collared cattle had no contact based on all contact definitions.




Figure S6. Mean standardised contact rates of cattle increase with spatiotemporal windows (i.e. buffer size and lagged time). Error bars are the standard error of the mean. The spatial window (contact distance or buffer size) was set at 50, 100, 200, 500 and 1000 m. The temporal window (lagged time interval) ranged from 1 to 168 hours (i.e. one hour to one week). The reference line at 200 meters indicates the assumed close contact distance. Herds from home bomas located within 5, 10 and 15 km comprise an area of 19.6, 78.5, 176.7 km², respectively.

Supplementary Tables

Table S1. Details of the binomial regression models (i.e. contact spatial window of 50 metres and temporal windows of 1 hour to 1 week), which examined factors that affect the probability of herd contacts relative to resource areas that was measured across a range of spatiotemporal proximities. Models were tested individually and continuous variables were scaled to mean zero and one standard deviation (SD).

Model	Fixed effects of final model	Model summary
Spatiotemporal proximity of 50 metres in one hour	Distance to boma and Herd size.	Most contacts between herds occurred away from their bomas and among small herd types. Odds ratio increased by 1.5 times (CI: 1.40 - 1.61) per one standard deviation (SD) increase in distance to bomas. The probability of encounter between herds decreased by 2.09 times with one SD increase in herd size.
Spatiotemporal proximity of 50 metres in one day	Distance to boma, Rainfall, Distance to the grazing, dipping points, and interaction terms of Distance to boma and Herd size.	A higher proportion of contacts occurred away from the bomas depending on herd size. Larger herds had higher contacts away from bomas, while the reverse was the case for smaller herd sizes as contact probability decreased by 0.81 (CI: 0.68 - 0.96) with one SD increase in herd size. Higher proportion of contacts occurred close to grazing areas (OR: 0.84; CI: 0.71- 0.99) and around dipping points (OR: 0.84; CI: 0.72 - 0.99).
Spatiotemporal proximity of 50 meters in one week	Distance to boma, Rainfall, Distance to watering and dipping points, Herd size and interaction terms of Rainfall and Distance to grazing.	The probability of contact was 2.58 (CI: 2.31 - 2.88) times higher with one SD increase in distance to bomas. Herd encounters were lowest with increased distance to watering and dipping locations. Although contacts were highest when rainfall level was lowest, higher contacts were observed away from grazing areas with increased rainfall.

Table S2. Details of the binomial regression modes (i.e. contact spatial window of 200 metres and temporal windows of 1 hour to 1 week), which examined factors that affect the probability of herd contacts relative to resource areas that was measured across a range of spatiotemporal proximities. Models were tested individually and continuous variables were scaled to mean zero and one standard deviation (SD).

Model	Fixed effects of final model	Model summary
Spatiotemporal proximity of 200 metres in one hour	Distance to boma, Herd size, Rainfall, Distance to watering point, and the interaction terms of Rainfall and Distance to watering point	Most contacts between herds occurred away from their bomas. Odds ratio increased by 2.83 times (CI: 2.36 - 3.40) per one standard deviation (SD) increase in distance. The probability of encounter between herds decreased by 0.88 with one SD increase in herd size. Contacts reduced significantly with increased distance to watering points, which decreased further with higher precipitation levels.
Spatiotemporal proximity of 200 metres in one day	Distance to boma, Herd size, Rainfall, Distance to grazing area and dipping point.	A higher proportion of contacts occurred away from the bomas. Odds ratio increased by 2.63 times (CI: 2.40 - 2.88) per one standard deviation (SD) increase in distance. Probability of encounters was higher for small herds compared to large. Contacts reduced significantly with increased distance to watering and dipping points .
Spatiotemporal proximity of 200 meters in one week	Distance to boma, Herd size, Rainfall, and Distance to grazing, watering and dipping points and their interaction terms.	A higher proportion of contacts occurred away from the bomas, which depended on herd size. Larger herds had higher contacts away from bomas, while the reverse was the case for smaller herd sizes as contact probability decreased by 0.80 (CI: 0.78 - 0.82) with one SD increase in herd size. Higher proportion of contacts occurred away from watering locations but close to dipping points and around grazing areas, mostly when precipitation levels were low.

Table S3. Details of the binomial regression models (i.e. contact spatial window of 500 metres and temporal windows of 1 hour to 1 week), which examined factors that affect the probability of herd contacts relative to resource areas that was measured across a range of spatiotemporal proximities. Models were tested individually and continuous variables were scaled to mean zero and one standard deviation (SD).

Model	Fixed effects of final model	Model summary
Spatiotemporal proximity of 500 metres in one hour	Distance to boma, Herd size, Rainfall, and Distance to grazing, watering and dipping points and their interaction terms	Most contacts between herds occurred away from their bomas. Odds ratio increased by 1.99 times (CI: 1.79 - 2.21) per one standard deviation (SD) increase in distance. Probability of encounters was higher for small herds compared to large. Contacts reduced significantly with increased rainfall, and increase distance to dipping points and watering points, but higher encounters were observed away from grazing locations which increased further with higher precipitation levels.
Spatiotemporal proximity of 500 metres in one day	Distance to boma, Rainfall, Distance to grazing, watering and dipping points and their interaction terms	The probability of contact was 2.61 times (CI: 2.47 - 2.75) per one standard deviation (SD) increase in distance to bomas, especially for large herd types. Herd encounters reduced with increasing distance to grazing location (OR: 0.86; CI: 0.82 - 0.90) and dipping (OR: 0.47; CI: 0.41 - 0.52), but increased with increasing distance to watering points, especially during high precipitation levels.
Spatiotemporal proximity of 500 meters in one week	Distance to boma, Rainfall, Herd size, Distance to grazing, watering and dipping points, and their interaction terms	The probability of contact was 2.89 times (CI: 2.78 - 3.00) per one standard deviation (SD) increase in distance to bomas, especially for large herd types. Herd encounters reduced with increasing distance to grazing location (OR: 0.92; CI: 0.90 - 0.95) and dipping (OR: 0.39; CI: 0.36 - 0.42), but increased with increasing distance to watering points, especially during high precipitation levels. Probability of encounters was higher for small herds compared to large.

References

1. Moritz, M., Galehouse, Z., Hao, Q. & Garabed, R. B. Can One Animal Represent an Entire Herd? Modeling Pastoral Mobility Using GPS/GIS Technology. *J. Hum. Ecol.* **40**, 623-630 (2012).
2. Gamo, R., Rumble, M., Lindzey, F. & Stefanich, M. GPS Radio Collar 3D Performance as Influenced by Forest Structure and Topography. *Biotelemetry* **15**, 464-473 (2001).
3. Gerlach, F. L. & Jasumback, A. E. Global Positioning System Canopy Effects Study. *US Department of Agriculture Forest Service, Technology and Development Center: Missoula, MT, USA.* (1989).