

6-gene machine learning-found signature for the differentiation of pancreatic ductal carcinoma and chronic pancreatitis patients on peripheral blood mononuclear cells RNA-Seq data

Francisco Carrillo-Perez¹, Octavio Caba^{2,3,4*},
Cristina Jiménez-Luna^{2,3,4}, Francisco Ortuño^{5,6},
Daniel Castillo-Secilla⁷, Luis Javier Herrera⁵, Jose Prados^{2,3,4},
Ignacio Rojas⁵

¹Stanford Center for Biomedical Informatics Research (BMIR), Stanford University, School of Medicine, Stanford, 94305-547, CA, USA.

²Institute of Biopathology and Regenerative Medicine (IBIMER), University of Granada, 18016, Granada, Spain.

³Biosanitary Institute of Granada (ibs.GRANADA)), ibs.GRANADA, 18014, Granada, Spain.

⁴Department of Anatomy and Embryology, Faculty of Medicine, University of Granada, 18071, Granada, Spain.


⁵Department of Computer Engineering, Automation and Robotics, University of Granada, 18014, Granada, Spain.

⁶Computational Medicine Platform, Andalusian Public Foundation Progress and Health-FPS, 41013, Sevilla, Spain.

⁷CoE Data Intelligence, Fujitsu Technology Solutions S.A, 28224, Madrid, Spain.

*Corresponding author(s). E-mail(s): ocaba@ugr.es;

Contributing authors: fcperrez@stanford.edu; crisjilu@ugr.es;
fortuno@ugr.es; daniel.castillosecilla@fujitsu.com; jherrera@ugr.es;
jcprados@ugr.es; irojas@ugr.es;



Supplementary Figure 1 Difference in expression levels for the 15-gene signature found using machine learning techniques on the GSE133684 dataset