

Article title: Morphological and molecular characterization of two populations *ex situ* regenerated from a germplasm bank accession of the wild potato *Solanum chacoense* Bitter

Journal name: Genetic Resources and Crop Evolution

Author names: POULSEN HORNUM, Anabela; CAMADRO, Elsa Lucila

Author for correspondence: CAMADRO, Elsa Lucila

Affiliation:

Unidad Integrada: Facultad de Ciencias Agrarias (FCA), Universidad Nacional de Mar del Plata (UNMdP)- Estación Experimental Agropecuaria (EEA) "Domingo R. Pasquale", Instituto Nacional de Tecnología Agropecuaria (INTA), Ruta Nacional 226 km 73.5, 7620 Balcarce, Buenos Aires, Argentina.

Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Godoy Cruz 2290, C1425FQB CABA, Argentina

E-mail adress:

camadro.elsa@inta.gob.ar






Fig.S1 Phenogram based on 11 quantitative (vegetative and reproductive) morphological characters evaluated in: A) 25 plants of the original accession CI 898 of *Solanum chacoense* Bitter (OA-S1), B) 19 plants of the population regenerated by the method proposed by Camadro (2012) (PR-MC) and 15 plants of the population regenerated by the method currently used in germplasm banks (FAO 2013) (PR-MGB).

*Note: plant= genotype. CCC Cophenetic Correlation Coefficient. Dotted line: arbitrary cut at Euclidean Distance