

Mapping ultra-processed foods (UPFs) in India: A formative research to adapt and inform a UPF consumption screening tool for India

Suparna Ghosh-Jerath

SGhosh-Jerath@georgeinstitute.org.in

The George Institute for Global Health, New Delhi, India <https://orcid.org/0000-0002-2229-4455>

Meenu Singh

Public Health Foundation of India, Gurgaon, Haryana, India

Gaurika Kumar

Public Health Foundation of India, Gurgaon, Haryana, India

Inderdeep Kaur Bhambra

Public Health Foundation of India, Gurgaon, Haryana, India

Sahiba Kohli

The George Institute for Global Health, New Delhi, India

Neha Khandpur

Wageningen University, the Netherlands; University of Sao Paulo, Brazil; Harvard T. H. Chan School of Public Health, US

Fernanda H. M. Leite

Center for Epidemiological Research in Nutrition and Health, Faculty of Public Health, University of Sao Paulo, Sao Paulo, Brazil

K. Srinath Reddy

Public Health Foundation of India, Gurgaon, Haryana, India

Research Article

Keywords: Ultra-processed foods, Nova food classification, Nova-UPF screener, UPF sub-categories, Tool adaptation, Online market survey, India

Posted Date: July 12th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3139180/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at BMC Public Health on August 14th, 2024. See the published version at <https://doi.org/10.1186/s12889-024-19624-1>.

Abstract

Background

Increased consumption of ultra-processed foods (UPFs) which are usually high in salt, sugar, fats, preservatives and additives are associated with diet-related non-communicable diseases. In India, there are no standard criteria for identifying UPFs that are similar to Nova food classification (based on extent and purpose of industrial processing). Scientific literature in India classifies foods as unhealthy based on the presence of excessive amounts of specific nutrients which makes it difficult to distinguish UPFs from other commercially available processed foods. The purpose of this paper is to map UPFs in the Indian food market and assess their consumption by Indian population.

Methods

A literature review and online grocery market survey were conducted to map the types of UPFs in the Indian food market, their availability and consumption. In online grocery market survey, the ingredient list of the foods provided on their label were scrutinized for the presence of ultra-processed ingredients. All UPFs identified were free-listed, followed by saliency analysis to understand preferred UPFs by Indian consumers. Findings were then triangulated to develop Indian UPF sub-categories to inform a UPF screening tool.

Results

UPFs lack uniform definition in India; hence proxy names such as *junk-foods*, *fast-foods*, *ready-to-eat foods*, *instant-foods*, *processed-foods*, *packaged-foods*, *high-fat-sugar-and-salt foods* are used for denoting UPFs. Our data showed that the current literature from India does not clearly differentiate between processed culinary ingredients (Nova food group 2), processed foods (Nova food group 3) and UPFs (Nova food group 4). Saliency analysis showed sugar-sweetened beverages (e.g. sodas and cold-drinks) are the most preferred UPFs while frozen ready-to-cook/eat foods (e.g. chicken nuggets and frozen kebabs) are least preferred. Online grocery market survey also revealed that traditional recipes are transitioning into UPFs. Based on our data, a screening tool for UPF consumption consisting of twenty-four UPF sub-categories were listed and divided into three sections: drinks and beverages, ready-to-eat and ready-to-cook foods was developed.

Conclusion

India needs to adapt Nova food classification and systematically define Indian food categories based on level of processing, followed by an assessment of the extent of UPF consumption in India. Clear policy guidelines and regulations around UPF consumption should be devised to address the impact of UPF consumption on NCDs.

Background

Non-communicable diseases (NCDs), one of the leading causes of premature morbidity and mortality across the globe, kills about 41 million people each year i.e. equivalent to over 7 out of 10 deaths worldwide (1). Deaths due to NCDs have been on the rise in India from 37.9% in 1990 to 61.8% in 2016 (2, 3), which is largely attributed to one of the major risk factors i.e. overweight/obesity (4). The recent national-level data shows an increasing prevalence of overweight and obesity among all segments of the Indian population (5, 6) Additionally, due to their thin fat phenotype, Indian infants and children, who comprise almost ¼ of the total population, are also predisposed to obesity (7, 8). These risk factors are further amplified by changing food environments and behavioural variables such as tobacco, alcohol, drug use and low

physical activity (9). Exposure to unhealthy food environments in genetically predisposed children further increases their risk of developing obesity and diet-related non-communicable diseases (DR-NCDs) in the long term (10).

The rapidly changing food environment is characterized by diets transitioning from minimally-processed staple foods (such as pulses and coarse cereals) high in vitamins, minerals and fibre to refined, processed and ultra-processed foods (UPFs) (11, 12). The Indian population is exposed to a wide variety of ultra-processed food products which are hyper-palatable, packaged, convenient, affordable and have a longer shelf life such as sugar-sweetened beverages, chips, commercial bakery products such as biscuits and bread, and ready-to-cook meals. These food products have increased in the food supply and are widely advertised and targeted towards vulnerable sections of the population including children and youth (13, 14). In addition, consumer demand for convenience foods has also increased due to the increased incomes, the number of nuclear families, single-member households and working women (15, 16). Further, these commercial UPFs have largely penetrated the rural boundaries of the country and are likely making their way into households of diverse geographic and socio-economic attributes (17, 18).

To identify UPFs from a wide range of food products available, a food classification system called the “Nova food classification system” based on the purpose and the level of processing was developed by Monteiro et. al. 2009 (19). The Nova system categorizes foods into four categories: (1) unprocessed/ minimally processed foods, (2) processed culinary ingredients, (3) processed foods and (4) ultra-processed foods (19). A series of industrial processes, chemical preservatives, additives, artificial colours and/ or flavours are used to produce the highly palatable and attractive UPFs (20). Several of these food products are high in saturated fats or trans-fats, added sugars, and salt and low in dietary fibre, various micronutrients and other bioactive compounds (21, 22, 23, 24, 25, 26, 27, 28, 29).

Overconsumption of UPFs has been associated with higher body mass index (BMI), obesity and DR-NCDs including type-2 diabetes, hypertension, cardiovascular diseases, and certain types of cancers (30, 31, 32). Given their potential role in contributing to the NCD burden in India, it is crucial to understand the range of UPFs available in the Indian market and their consumption. Therefore, this paper aimed at (i) mapping the types of UPFs in the Indian food markets, (ii) assessing the composition of these food products to confirm that they qualify as UPFs (iii) assessing their consumption among Indian consumers, (iv) listing the UPFs in order of their preference among the Indian consumers. This exercise provided vital information for developing India-specific UPF sub-categories to inform a tool that would be used for screening UPF consumption in the Indian population.

Methods

The present study was based on a secondary review of available literature complemented by an online grocery market survey that was conducted between April 2021 and February 2022. The literature review was conducted in phase I to map the various types of UPFs reported in different studies assessing their consumption and availability among various population groups and regions in India. It also aimed at collating proxy nomenclatures used for UPFs (any food classification such as *junk food*, or *convenience food* that is used to identify foods and beverages with characteristics of UPFs as defined by the Nova food classification system) in peer-reviewed research on food intake in India. Articles for review were searched from three electronic databases (Medline, NCBI and Google Scholar). Based on the proxy names for UPFs used in the policy documents (33, 34, 35, 36, 37, 38) a literature search was conducted by using a combination of the following key terms while incorporating the Boolean logic: *junk food**, *fast food**, *modern food**, *westernized food**, *ultra-processed food**, *UPF**, *convenience food**, *ready-to-eat food**, *ready-to-eat snack**, *ready-to-cook food**, *RTE instant food**, *frozen food**, *canned food**, *tinned food**, *processed food**, *packaged food**, *high fat*, *sugar and salt food** and *HFSS**. Search limiters such as India for location and publication dates between January 2012 and December 2022 were used.

The inclusion criteria were, studies: (i) that included well-defined UPFs or their proxy nomenclature with examples, (ii) conducted in rural and/or urban areas of India, (iii) published in the English language, (iv) published between 2012 and 2022 (based on the high growth in UPFs sales during this decade) (39, 40). Publications that did not define the food

category studied such as processed food, UPF, HFSS food or citing any examples of foods under the category were excluded from the review. This study did not include review articles on the consumption or availability of UPFs in India. Data from articles reviewed were collated into MS-Excel to record key variables such as proxy terms for UPFs, their examples, location of the study (national/specific state), geographical area (urban/rural), sample size, sampling method/study design, age group (years) consuming the food category and data collection tool (Additional file 2).

An online grocery market survey for extracting detailed information on the UPFs identified in the mapping exercise was also conducted which complemented the literature review. The objective was to review the specific food labels (particularly the ingredient lists) and confirm if the products qualified as UPFs. For this online survey, three researchers reviewed the most popular Indian online grocery stores like Big Basket, Grofers, and Amazon which are considered as largest grocery retail brands (41). During the survey, the preliminary free list of foods and beverages considered as UPFs prepared from the literature review were cross-checked for their composition and processing techniques based on their food labels. This activity confirmed that the food product under question qualified as UPF. All the food products were scrutinized for the steps of processing and use of ingredients such as sugars (fructose, invert sugar, maltodextrin, dextrose, lactose, high fructose corn syrup, fruit juice concentrates), modified oils (hydrogenated fats, interesterified fats), extracted proteins (hydrolysed proteins, soy protein isolate, gluten, casein, whey protein, mechanically separated meat), additives (flavours, flavour enhancers, colours, emulsifiers, emulsifying salts, artificial sweeteners, thickeners, acidity regulators and foaming, anti-foaming, bulking, carbonating, gelling, glazing, leavening and anticaking agents,) and class II preservatives (such as benzoic acid, sorbic acid, sodium and calcium propionate, sodium diacetate) which classified them as UPFs (19, 42).

Using the data from the literature reviewed, a saliency analysis was conducted to determine the commonly purchased, preferred and/or consumed (collectively referred to as 'preferred' in this paper) UPFs in India. Indicators including purchasing trends, consumer preferences and frequency of consumption of UPFs amongst Indian consumers were extracted. Saliency is a statistical accounting of items for rank and frequency of mention, across all respondents within a given domain (for example, a colour chosen most often from a free list of ten colours by a study population, is referred to as most salient) (43). For this, the UPFs identified were sub-categorized and sorted out as 'most to the least preferred' UPFs by consumers (Additional file 1). Accordingly, a saliency score was calculated for each sub-category of UPFs using a set of steps and formulas (43), as illustrated in Fig. 1. The UPF sub-categories were then classified (as per consumer preference) according to the composite salience score cut-offs (defined after dividing the distribution of the composite salience scores obtained for each sub-category into thirds) as follows: (i) ≥ 0.61 as commonly preferred, (ii) $0.61 - 0.51$ as moderately preferred, and (iii) < 0.51 as rarely preferred UPFs.

Results

The literature search and selection process is illustrated in Fig. 2. A total of twenty-three research articles that matched the inclusion criteria were included in the final review. An overview of the articles with information on variables such as the proxy nomenclature for UPFs used, examples of food products under the proxy nomenclature, location, geography (rural and/or urban) where the study was conducted, age groups of the study participants, sample size, sampling methodology and data collection method is provided in Additional File 2. These studies were conducted in both rural and/or urban areas of different regions of the country among a diverse population aged between 9 to 69 years (Table 1).

Table 1
Demographic characteristics of the participants from the reviewed studies (n = 23)

Region	Number of studies	Geographical area				Age group (years)
		Rural	Urban	Both	Not Defined	
East	2	0	1	1	0	12–18
West	3	0	3	0	0	10–24
North	5	1	4	0	0	9–20
South	9	1	5	0	3	11–60
PAN India	4	0	1	2	1	10–69
TOTAL	23	2	14	3	4	9–69

Several proxy nomenclatures and classifications such as fast foods, junk foods, instant foods, processed foods etc. used for UPFs in the Indian context were identified in peer-reviewed literature. Table 2 details the proxy nomenclature used along with the food items listed under those categories. Several processed foods (Nova group 3) and culinary ingredients (Nova group 2) were clubbed under these proxy nomenclatures of UPF.

Table 2
Nomenclature and examples of foods considered as UPFs and their proxies in Indian studies (n = 23)

UPF and its proxy nomenclature	Foods and beverages identified as UPF	Non-UPFs items clubbed with UPF	References
Ultra-processed foods	Chips, cake, ready-to-serve beverages, beverage concentrates, bakery products and confectionery		44
HFSS	<i>Namkeen, papad</i> , packaged chips, soya or mustard sauce, processed or packaged cheese, butter	<i>Papad, namkeen</i> (if freshly prepared)	45
Processed foods	<p>Sweet snacks: chocolate, honey, biscuits, cookies, jams, peanut butter, chocolate spread, <i>rusk</i> (excludes prepared sweets and cakes)</p> <p>Salty snacks: crackers, potato chips, banana chips, and other salty snacks</p> <p>Soft drinks: carbonated drinks, juices, milk-based drinks, squashes and powdered drinks</p> <p>Milk: liquid milk and milk powder</p> <p>Dairy products: butter, <i>ghee</i> (clarified butter) and cheese</p> <p>Edible oils: edible oils and <i>vanaspati</i> (partially hydrogenated vegetable oil)</p> <p>Processed wheat: <i>atta</i> (wheat flour), bread, vermicelli and pasta</p> <p>Other processed foods: soup, ready-to-eat meals, ready-to-cook mix, frozen food, breakfast cereals, noodles, ketchup, table sauces</p>	<p>Honey</p> <p>Liquid milk</p> <p>Butter, <i>ghee</i></p> <p>Edible oils</p> <p><i>Atta</i> (Processed wheat), vermicelli, pasta</p> <p>Breakfast cereals (if not coated/extruded)</p>	46
	<p><i>Masala puri, churmuri, panipuri</i>, noodles, <i>gobi manchurian</i>, fried rice, coffee, tea, cola beverages, fruit beverages, <i>kharasev, kodubale, chakli, bajji, bonda, vada</i>, chips, puffs, cake, bread, <i>rusk</i>, bun, puffed rice</p>	<p><i>Masala puri, churmuri, panipuri</i>, fried rice (if freshly prepared), coffee, tea, <i>kharasev, kodubale, chakli, bajji, bonda, vada</i>, (if freshly prepared), puffed rice</p>	17
Packaged foods	<p>Edible oils, ready meals, sauces, dressings and condiments, soup, sweet spreads, baby foods, dairy</p> <p>Confectionery, ice cream and frozen, desserts, savoury, snacks, sweet biscuits, snack bars and fruit snacks, baked goods, breakfast cereals, processed fruit and vegetables, processed meat and seafood</p> <p>Rice, pasta and noodles</p>	<p>Edible oils</p> <p>Plain breakfast cereals, processed fruits and vegetables</p> <p>Rice, pasta, and noodles</p>	40
Junk foods	Burger, pizza, pasta, instant noodles	-	6
	Chips, chocolates, bakery products (pastries, cream rolls, patties), soft drinks, sweetened beverages (squashes, sweet juices), sweets (<i>ladoo, jalebi, barfi</i>), ice cream	Sweets (<i>ladoo, jalebi, barfi</i>) and <i>samosa</i> (if freshly prepared)	18

UPF and its proxy nomenclature	Foods and beverages identified as UPF	Non-UPFs items clubbed with UPF	References
	<p>South Indian: <i>Idli, dosa, vada, uttapam</i></p> <p>Bakery items: Biscuits, bread, cake, pastry, patties</p> <p>Italian food: Pizza, pasta, macaroni</p> <p>Chinese: <i>Maggi, manchurian</i>, noodles, spring roll</p> <p>Sweet dish: Chocolate, <i>halwa</i>, ice cream, <i>mithai</i></p> <p>Fried food: bread <i>pakora</i>, burger, <i>chaat, kachori, kulcha chana</i>, potato chips, <i>samosa, tikki</i></p> <p>Beverages: tea, coffee, carbonated drinks, juices</p>	<p><i>Idli, dosa, vada, uttapam</i> (if freshly prepared)</p> <p>Bread <i>pakora, chaat, kachori, kulcha chana</i>, potato chips, <i>samosa, tikki</i></p> <p>Tea, coffee</p>	47
	Ice-creams, sweets, chocolates, fried foods, bakery foods, pizza, fruits, carbonated beverages	Fried and bakery foods (if freshly prepared)	48
	Pizza, burger, puffs, pastry, cake, biscuits, french fries, noodles, doughnuts, Pepsi, Coca-Cola, Slice, 7-UP, Bovonto, Sprite	-	49
	Cakes, chocolates, ice-creams, sweets, chips, <i>Kurkure</i> , street fast foods, Indian deep-fried foods, noodles	Sweets, Indian deep-fried foods, street fast foods (if freshly prepared)	50
Indian fast foods	<i>Chat, pakora, samosa</i> , patties, <i>panipuri, puri-kachori, chole bhature</i> , stuffed <i>paratha, idli, dosa, vada, upma, poha, jalebi</i> , sweets, non-veg items, roasted veg and non-veg items	<i>Chat, pakora, samosa, patties, panipuri, puri-kachori, chole bhature, stuffed paratha, idli, dosa, bada, upma, poha, jalebi</i> , sweets, non-veg items, roasted veg and non-veg items	51
Fast foods	Carbonated drinks, fruit juice, milk and shakes	Fruit Juices (if freshly prepared), milk	52
	Pizza, burger, chocolate, ice cream, cookies/cake, <i>chowmien</i> , pasta and <i>Maggi</i> .	Pasta	53
	Chocolates/pastries/sweets, <i>papads</i> and pickles, biscuits or other bakery items like bread, toast, buns, cold drinks/soft drinks, potato chips/ <i>namkeens</i> /deep-fried snacks, ice cream/milkshakes, <i>paneer</i> , cheese/butter, pizza/burger/frankie or any other fast food including Chinese food	Sweets, <i>papads</i> and pickles (if freshly prepared), <i>namkeens</i> /deep-fried snacks, <i>paneer</i> , butter	54
	Pizza, burger, <i>samosa</i> , chocolates	<i>Samosa</i> (if freshly prepared)	55
	Fizzy drinks, tea, coffee, mocktails, slushy, milkshakes, pizza, french fries, burger, sub, rolls, cakes/pastries	Tea, coffee	56
	Pastries, pizza, french fries, cheese items, Chinese food, soft drinks, coffee/tea, sweetened fruit drinks, energy drinks	Coffee/tea	57
Fast foods, Junk foods, Instant foods, street foods	Fast foods: Burgers, pizzas, fries, hamburgers, patties, and nuggets, Indian foods like <i>pakora, samosa, namkeen</i> etc.	<i>Pakora, samosa, namkeen</i> etc (if freshly prepared)	58
		Corn flakes	

UPF and its proxy nomenclature	Foods and beverages identified as UPF	Non-UPFs items clubbed with UPF	References
	<p>Junk foods: Chips, chocolate, ice cream, soft drinks, and sandwich</p> <p>Instant foods: Noodles, corn flakes, and soup powder</p> <p>Street foods: <i>Chaat, gol gappa, samosa, tikki, noodles, chowmein, and vada pav</i></p>	<i>Chaat, gol gappa, samosa, tikki, (if freshly prepared)</i>	
Convenience foods	Ready-to-eat (RTE) <i>dosa</i> , RTE <i>idli</i> , RTE <i>chapathi</i> and RTE <i>parathas</i> , RTE <i>upma</i>	-	59
Modern foods	Bread, fruits and juices, cereals, eggs and oats	Fruits and juices (freshly prepared), cereals, eggs, oats	59
Western fast foods	Pizza, burger, french fries, doughnuts, hot dog, noodles	-	51
Food, snacks and beverage	Breakfast cereal, bread/toast, biscuits/cookies, cake/pastries, french fries, packaged potato chips, pizza, burger, ice cream, ice candy, chocolates, vegetable roll/wrap, chicken roll/wrap/nuggets, egg roll, popcorn, noodles, tea (black tea/milk tea/Irish tea), coffee (cold coffee/black coffee), health drinks (<i>Bournvita/Horlicks</i>), soft drink (Sprite, Coke, Pepsi), energy drink (Red Bull, Gatorade)	Breakfast cereal (plain), tea (black tea/milk tea/Irish tea), coffee (cold coffee/black coffee)	60
Instant food products	<i>Rasam</i> mix, lemon rice mix, <i>puliyogare</i> , brinjal <i>kulambu</i> mix, pasta, <i>sambhar</i> rice and masala, <i>pulao</i> mix and <i>paneer</i> butter masala	-	61
Soft drinks	Pepsi, Fanta, Slice, Maaza, Coca-Cola	-	62

Table 2 demonstrates a lack of standard and uniform definitions for UPFs in India. Nova food classification categories were merged and addressed with proxy nomenclatures of UPFs such as junk, modern, westernized, convenience, ready-to-eat/cook, HFSS, packaged and processed foods in included studies. Thus, with this limited data with no clear-cut definition for UPFs, the actual estimation of the consumption of UPFs in the Indian context is difficult.

The online grocery market survey supplemented the information derived from the literature review and complemented information on the preliminary free list of UPFs identified in the Indian food environment. Packaged Indian traditional foods and snacks such as bottled and packaged pickles, *namkeens* (cereal and pulse-based extruded snacks), *papads*, frozen non-vegetarian meals and snacks, and frozen ready-to-eat meals (like *rajmah curry* and rice, *biryani*, *dal makhni*, etc.) were specifically checked for ultra-processing, and presence of ingredients and additives that qualified them as UPFs. Table 3 presents the ingredients found in these traditional snacks that confirmed their UPF classification. Ready-to-eat meals were excluded from the list of UPFs as their composition did not include ingredients that qualify them as UPFs. Label readings of 375 packaged food items were completed and 81 foods were identified as UPFs. Extensive label reading provided more clarity on the ingredients of the identified UPFs and their correct classification.

Table 3
List of traditional Indian snacks/meals and ingredients that qualify them as UPFs

Traditional Indian snacks/meals	Ingredients that qualify for UPFs
Pickles	Acidity Regulator (INS 330, 260), Preservative (INS 211), Capsicum Extract, Paprika Extract
<i>Namkeens</i>	Maltodextrin, Dextrose, Milk Solids, Acidity Regulators (INS 330 & 296), Flavor Enhancers (INS 627 and INS 631), Citric Acid (INS 330), Paprika Extract (INS 160 C), Tomato Powder, Tamarind Powder, Date Powder, Anticaking Agent (INS 551), Spinach Powder, Edible Starch, Potato Solids, Natural [E100 and E141(ii)] and Synthetic food colour (E133 / FD & C Blue No.1) and Added flavour (Nature identical flavouring), Leavening Agent (INS 500 (I), Synthetic Food Colours (INS 133), Compounded Asafoetida [Carrier (INS 414), Soy Lecithin (E322), Antioxidant (E319)
Frozen non-vegetarian foods/snacks	Emulsifiers [INS 451 & INS 450], Sugar, Antioxidant [INS 250], Milk Solids, Modified Starch, Milk Protein, Stabiliser (sodium triphosphate), Onion extract, Spice extracts, Preservative (sodium nitrite), Soya Protein, Texturizer (INS 451), Preservative (INS 250), Color Fixative (INS 250), Raising Agent (INS 500), Maltodextrin, Wheat Fibre
<i>Papads</i>	Raising agent [INS 500 (ii)], Preservatives (INS 200), Raising agent [INS 500 (i) and INS 500 (ii)], Sodium bicarbonate [E 500 (ii)]

Several food items were also found to be in transition from processed foods to UPFs. This was because some brands of these foods belonged to the processed foods category while other brands fitted into the category of UPFs as per the Nova food classification system. Such foods include ready-to-eat (RTE) breakfast cereals (e.g. *poha*, *upma*, etc.), RTE Indian curries (e.g. *paneer makhani*, *butter chicken*, etc.), RTE meals (such as *biryani*, etc.), Indian RTE bread (e.g. *thepla*, *paratha*, etc.), ready-to-cook mixes (e.g. *idli* mix, *dal vada* mix, etc.) and many more. Traditional packaged foods like *poha*, instant *idli* or *dosa* or *sambhar* or *dhokla* mixes, etc. qualified as UPFs after screening their labels based on their level of processing and the presence of additives.

Following this, the specific UPFs identified were sub-categorised and consumer preferences were assessed using saliency scores. Table 5 shows the order of preference towards different UPF sub-categories based on the saliency scores. The last column in the table indicates 'commonly', 'moderately' or 'rarely' preferred UPFs by consumers in India. The commonly preferred UPFs were sugar-sweetened beverages (such as cold drinks, diet coke, sodas, and energy drinks) followed by extruded snacks (such as potato chips, cheese balls, puff corns, etc.) and breads. Meanwhile, the three rarely preferred UPFs were margarine and frozen/ packaged vegetarian and non-vegetarian snacks and meals (such as stuffed/plain *parantha*, *naan*, *palak paneer*, *rajma*, cutlets, fish/seafood snacks, salami, sausages).

The outcome of this extensive exercise of literature review, online grocery market survey and saliency analysis was the development of 24 sub-categories of UPFs for India (Table 4).

Table 4
Sub-categories of UPFs (n = 24), divided into three sections

<p>Section A – Drinks and Beverages</p> <p>A1. Aerated/cold/soft drinks; Diet drinks</p> <p>A2. Energy drinks; Sports drinks (like clear drinks with added electrolytes, vitamins, minerals)</p> <p>A3. Packaged and branded, bottled or tetra packs fruit/vegetable-based juices (like mango juice, orange juice, vegetable juice); Packaged and branded drink concentrates (like <i>sherbets</i>, <i>thandai</i>); Packaged and branded powdered drink mixes (like fruit flavoured mixes, iced tea mix)</p> <p>A4. Packaged and branded flavoured milk (like chocolate milk); Packaged and branded milk or malt-based powdered health drinks, Protein powder; Packaged and branded yogurt/curd-based drinks (like flavoured <i>lassi</i>, probiotic drinks); Packaged and branded milk substitutes (like soymilk, almond milk); Ready-to-drink tea/coffee mixes (like masala tea, choco mocha); Dairy whiteners</p>
<p>Section B – Ready-to-eat</p> <p>B1. Packaged and branded bread (like sliced bread, pao, burger buns, pizza base, tortillas)</p> <p>B2. Packaged and branded biscuits, cream biscuits, cookies, cream puffs/rolls</p> <p>B3. Packaged and branded Indian namkeens (like aloo bhujia, mixtures, murukku, flavoured and coated nuts)</p> <p>B4. Packaged and branded chips (plain and flavoured), nachos, puffs</p> <p>B5. Pizza, burgers, french fries, wraps from fast-food chains</p> <p>B6. Packaged and branded extruded coated breakfast cereals (like sugar-coated cornflakes, chocolate breakfast cereals, ragi bites); Packaged, and branded cereal bars (like granola bars, energy bars)</p> <p>B7. Packaged and branded cakes, muffins, waffles, donuts; Packaged and branded Indian sweets (like flavoured <i>sonpapadi</i>, <i>gulab jamun</i>)</p> <p>B8. Packaged, branded and flavoured yogurt, fruit yogurt, <i>kheer</i>/ <i>payasam</i></p> <p>B9. Packaged and branded ice creams, flavoured ice bars, <i>kulfi</i></p> <p>B10. Packaged and branded fruit-based preserves (like jams, marmalades, jellies)</p> <p>B11. Chocolates; Toffees; Lollipops; Chewing gums; Fruit candies; <i>Aam papad</i>/ Fruit leathers/ bars; Flavoured mouth fresheners</p> <p>B12. Packaged and branded ketchup, chutneys/ instant chutney powders/tastemakers (like tamarind chutney, green chutney, lemon rice spice mix); Packaged and branded pickles (veg./ non-veg.), sauces (like pasta-pizza sauce), instant gravies/curries/pastes (like mustard paste, puliyogare paste, ginger-garlic paste)</p> <p>B13. Packaged and branded milk-based spreads, mayonnaise, dips, cheese products (like cheese slices/cubes/spreads), salad dressings, nut spreads (like hazelnut spread, peanut butter), chocolate spreads</p> <p>B14. Margarine; Packaged and branded flavoured butter (like garlic butter), coconut cream/milk</p>
<p>Section C – Ready-to-cook</p> <p>C1. Packaged and branded instant soups, instant noodles/ pasta</p> <p>C2. Packaged and branded instant dishes/snacks (like <i>poha</i>, <i>upma</i>, savoury oats); Packaged and branded ready-to-cook powdered mixes (like <i>idli</i>, <i>dosa</i>, <i>dhokla</i>, <i>aloo bonda</i>, <i>dahi vada</i>)</p> <p>C3. Packaged, branded and frozen ready-to-cook vegetarian snacks (like french fries, paneer snacks, vegetarian burger patty, <i>aloo tikki</i>, veg. <i>samosas/pizza</i>); Packaged and branded frozen <i>parathas</i>, puff pastry/spring roll sheets</p> <p>C4. Packaged, branded and frozen ready-to-cook non-vegetarian snacks (like chicken nuggets, chicken tikka, kebabs, seafood/fish snacks, sausages, salami, non- veg. pizza/momos)</p> <p>C5. Packaged and branded ready-to-cook meals in cups (like <i>rajma chawal</i>, <i>biryani</i>, <i>idli sambar</i>, <i>bisi-belle bath</i>, <i>schezwan rice</i>)</p>

Section A – Drinks and Beverages

C6. Packaged and branded bread mixes; Packaged and branded dessert mixes (like jelly, custard, ice cream, *gulab jamun*, cake, brownie, pancake)

Discussion

The present study documents the findings of a literature review and an online grocery market survey to map UPFs available in Indian food markets and their consumption. No standard nomenclature was followed in the studies for identifying categories of UPFs consumed in India during the literature review period, (2012–2022). Studies have merged several food categories under proxy names to provide a profile of the UPF environment in India. Evidence from other countries such as Argentina, Ecuador and Brazil also indicates the need for a standard, clear-cut definition of UPFs to prevent their misclassification (63, 64). Saliency analysis identified the preferred UPFs among the Indian population with Sugar-sweetened beverages being the most preferred UPFs and frozen non-vegetarian snacks being the least preferred. This could also be attributed to the fact that sugar-sweetened beverages have been present in the Indian market for a long period of time while frozen snacks are relatively a new entry into the Indian market. After analysing the ingredient list of UPFs, it was observed that product formulation of several traditional Indian foods has transitioned from unprocessed or minimally processed ingredients to ultra-processed ingredients. Thus, the new version of traditional Indian foods has transitioned into UPFs with the addition of synthetic ingredients such as artificial colours, flavours, anti-caking agents, etc. The study findings contributed towards the development of 24 sub-categories of UPFs for India based on the principles of the Nova food classification system.

In India, there are several proxy terms used for UPFs and there is a lack of clarity in the classification of foods, with 'fast foods' and 'junk foods' being the most used terms to indicate UPF foods. The presence of unhealthy foods in the Indian food environment has been associated with overweight, obesity and NCDs (30, 65, 66, 67, 68). Therefore, it is crucial to understand the actual consumption of UPFs and its health consequences in India. The Indian UPF market is slowly expanding with increasing sales of ready-to-eat meals, savoury snacks, processed fruits, vegetables, meats and other packaged foods as per the packaged food sales data from 2015-19 (40). However, for exploring these associations in the context of UPFs, a food classification system needs to be adapted in India as in the Indian food policy environment, 'HFSS foods' is still the most commonly used terminology, a large majority of which are UPFs (60, 61, 63, 69, 70, 71). Therefore, the Nova food classification system can serve this purpose and can be accepted by the food regulatory authorities. The Nova food classification system has been well accepted and used to assess dietary patterns in several high and middle-income countries such as Canada, The United Kingdom, The United States, Chile, Sweden, New Zealand, Australia, Mexico, Norway, Spain, Belgium, Colombia, France and Taiwan (19, 72).

The present paper identified only a limited number of Indian studies which were primarily reported from 2 geographical regions. More such surveys on the consumption of UPFs are desirable to identify common regional UPFs. In the Indian context, several UPFs are indigenously produced by local retailers apart from the huge market share of renowned branded UPFs. These locally accessible UPFs have better penetration into the local markets. Regulations are highly desirable to identify these sources of UPFs. An appropriate regulatory environment can protect consumers against the marketing of these potentially unhealthy foods. Consumer education regarding these UPFs is also desirable.

Saliency analysis indicated the preference of Indian consumers towards specific UPFs including sugar-sweetened beverages, extruded snacks, breads, fruit-based preserves, cookies and biscuits, Indian sweet mixes, sauces and pickles, instant noodles/soups/ pasta and savoury puff rolls. Studies from other Low and Middle-Income Countries (LMICs) demonstrated similar trends in preference towards packaged confectioneries, savoury snacks, deep-fried foods, biscuits, candy/ chocolate, savoury snacks, canned red and luncheon meats, pre-fried french fries, mayonnaise, ketchup, fast-food such as sandwiches and pizzas, chips and salty snacks (including tortillas and pretzels), sweets and sweetened beverages

and sausages (including canned) (73, 74). Increasing sales of packaged foods in various countries including several LMICs has been highlighted by the Euromonitor data (40).

Indian food industries are converting traditional recipes into UPFs which are being marketed as convenience foods. Consumers need to be made aware of the negative health effects of consuming these modified traditional foods that are ultra-processed. Additionally, regulatory measures should be implemented to monitor misleading claims, such as labelling these modified traditional foods as 'healthy'. These modified versions of traditional foods contain UPF ingredients, are highly processed and often sold and perceived as "healthy foods". To safeguard consumer health, it is important to appropriately identify UPFs associated with the rising incidence of DR-NCDs in India. Attention is needed to prevent other traditional foods, which are also gradually transitioning into UPFs.

Conclusions

There is a need to define UPFs, specify sub-categories and devise robust guidelines for defining UPFs in the Indian context. A national survey based on an exhaustive categorization of UPFs may provide crucial information on UPF consumption patterns in India. This is likely to inform policies on regulating the Indian UPFs market, undertake consumer education initiatives and create nutrition literacy around UPFs and thus contain their indiscriminate consumption.

Strengths

This review is one of the first attempts to explore the UPFs in the Indian food market and understand their availability and consumption along with the proxy nomenclatures used for UPFs in India. This review highlights gaps in the Indian food regulatory environment on the availability and sales of UPFs, which will direct the attention of food, nutrition, and public health diaspora to make efforts in channelling and strengthening the food regulatory environment towards UPFs.

Study limitations

There is an unavailability of representative data from PAN India on UPF consumption to extrapolate the findings. Most of the studies were documented from the Southern part of India and largely represent the urban population, hence the data is not representative of the rural population. Currently, several UPFs are referred to as junk foods, fast foods, western foods, processed foods, HFSS, etc. Due to the lack of a standard definition of UPFs, culinary ingredients (category 2 of Nova food classification) and processed foods (category 3 of Nova food classification) were clubbed under proxy categories of UPFs and therefore, the findings of various studies were difficult to interpret and compare. We also could not look into the packaging of the foods and beverages explored as this information is not usually declared on the packages.

List Of Abbreviations

UPFs - Ultra-processed foods

NCDs - Non-communicable diseases

DR-NCDs - Diet-related non-communicable diseases

HFSS - High Fat Sugar Salt

LMICs - Low and Middle-Income Countries

RTE - Ready-to-eat

Declarations

Ethics approval and consent to participate

Approval was obtained from the Institutional Ethics Committee at the Public Foundation of India, and the ethics committee of University of Sao Paulo. Informed consent was obtained from all participants.

Consent for publication

Not applicable

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Competing interests

The authors declare that they have no competing interests.

Funding

This work is funded through the Innovative Methods and Metrics for Agriculture and Nutrition Action (IMMANA) programme (Grant IMMANA 3.06), led by the London School of Hygiene & Tropical Medicine (LSHTM). IMMANA is co-funded with UK Aid from the UK government and by the Bill & Melinda Gates Foundation INV-002962 / OPP1211308. Under the grant conditions of the Foundation, a Creative Commons Attribution 4.0 Generic License has already been assigned to the Author Accepted Manuscript version that might arise from this submission.

Author's contribution

The study was conceptualized by KSR, SGJ, NK and FHML. The literature review and market survey were conducted by MS, GK, IKB. The first draft of the manuscript was prepared by MS, GK, IKB, SK and SGJ. The manuscript was critiqued and edited by SGJ, NK, FHML and KSR. SG-J had primary responsibility for final content; and all authors read and approved the final manuscript.

Acknowledgement

The authors would like to thank the experts from the regional and national consultations for their invaluable contribution for developing the UPF sub-categories for India. We are grateful to Dr Suneeta Chandorkar, Dr Pulkit Mathur, Dr Pranati Das, Dr Kanika Varma, Dr Meenakshi Bajaj and Ms Mita Shukla for coordinating and facilitating the regional consultations. We would also like to thank Dr Eram Rao for helping us identifying key experts for the national consultation. We would also appreciate the contribution of Dr Shukrani Shinde for her support for coordinating the expert consultations and online grocery market survey.

Author's Information

SGJ (PhD.) is Program Head, Nutrition at George Institute (TGI), India; MS (PhD.) is subject matter expert at Public Health Foundation of India (PHFI); GK (MSc) and IKB (MSc) are research assistants at PHFI; SK (PhD.) is a consultant at TGI; NK is a Research Scientist in University of São Paulo (USP); FHML is a Researcher at Nupens/USP; KSR (DM) is the former president of PHFI.

References

1. World Health Organization. Non-communicable diseases country profiles 2018. Available from: <https://apps.who.int/iris/handle/10665/274512>. [Cited 18 January 2023].

2. World Health Organization. Non-communicable diseases Progress Monitor 2022. Available from: <https://apps.who.int/iris/handle/10665/353048>. [Cited 18 January 2023]
3. Indian Council of Medical Research (ICMR), Public Health Foundation of India (PHFI) and Institute for Health Metrics and Evaluation (IHME). India: Health of the Nation's States-The Indian State-level Disease Burden Initiative. New Delhi: ICMR, PHFI, and IHME; 2017). Available from: https://phfi.org/downloads/171110_India_Health_of_Nation_states_Report_2017.pdf [Cited 20 January 2023]
4. Alwan A. Global status report on non-communicable diseases 2010. World Health Organization, 2011. Available from: https://apps.who.int/iris/bitstream/handle/10665/44579/9789240686458_eng.pdf. [Cited 8 May 2023]
5. International Institute for Population Sciences (IIPS) and ICF. National Family Health Survey (NFHS-5), India, 2019-21: Mizoram. Mumbai: IIPS. 2021. Available from: https://pdf.usaid.gov/pdf_docs/PA00ZBN4.pdf . [Cited 18 January 2023]
6. Sethi V, Lahiri A, Bhanot A, Kumar A, Chopra M, Mishra R, Alambusha R, Agrawal P, Johnston R and de Wagt A. Adolescents, Diets and Nutrition: Growing well in a Changing World, The Comprehensive National Nutrition Survey, Thematic Reports, (1), 2019.
7. D'Angelo S, Yajnik CS, Kumaran K, Joglekar C, Lubree H, Crozier SR, et al. Body size and body composition: a comparison of children in India and the UK through infancy and early childhood. *J Epidemiology Community Health*. 2015 July 16;69(12):1147.
8. India - Sample Registration System (SRS)-Statistical Report 2020 . Available from: <https://censusindia.gov.in/nada/index.php/catalog/44376>. [Cited 18 January 2023].
9. Dandona L, Dandona R, Kumar GA, Shukla DK, Paul VK, Balakrishnan K, et al. Nations within a nation: variations in epidemiological transition across the states of India, 1990–2016 in the Global Burden of Disease Study. *The Lancet*. 2017 Dec;390 (10111):2437–60.
10. Laar A, Barnes A, Aryeetey R, Tandoh A, Bash K, Mensah K, Zotor F, et al. Implementation of healthy food environment policies to prevent nutrition-related non-communicable diseases in Ghana: National experts' assessment of government action. *Food Policy*, 93, 101907. 2020. <https://doi.org/10.1016/J.FOODPOL.2020.101907>
11. Gaiha R, Kaicker N, Imai K. Has Dietary Transition Slowed in India? An Analysis Based on the 50th, 61st and 66th Rounds of the National Sample Survey. 2013.
12. Baker P, Friel S. Processed foods and the nutrition transition: Evidence from Asia. *Obesity reviews*. 2014 Jul;15(7):564-77.
13. Bassi S, Bahl D, Arora M, Tullu FT, Dudeja S, Gupta R. Food environment in and around schools and colleges of Delhi and National Capital Region (NCR) in India. *BMC Public Health*. 2021 Dec;21(1):1-3.
14. Harris J, Heard A, Schwartz M. Older but still vulnerable: all children need protection from unhealthy food marketing. Yale Rudd Center for Food Policy & Obesity. 2014; Available from: www.uconnruddcenter.org/files/Pdfs/Protecting_Older_Children_3_14.pdf. [Cited 24 December 2022]
15. Monteiro CA, Moubarac JC, Cannon G, Ng SW, Popkin B. Ultra-processed products are becoming dominant in the global food system. *Obesity reviews*. 2013 Nov;14:21-8.
16. Magalhães V, Severo M, Correia D, Torres D, de Miranda RC, Rauber F, Levy R, Rodrigues S, Lopes C. Associated factors to the consumption of ultra-processed foods and its relation with dietary sources in Portugal. *Journal of nutritional science*. 2021;10.
17. Prakash J. Consumption trends of processed foods among rural population selected from South India. *Int J Food Nutr Sci*. 2015;2(6):1-6.
18. Gupta A, Kapil U, Singh G. Consumption of junk foods by school-aged children in rural Himachal Pradesh, India. *Indian journal of public health*. 2018 Jan 1;62(1):65.

19. Monteiro CA, Cannon G, Lawrence M, da Costa Louzada ML, Pereira Machado P. Ultra-processed foods, diet quality, and health using the NOVA classification system. 2019: 48 p.
20. Monteiro CA, Cannon G, Levy RB, Moubarac JC, Louzada ML, Rauber F, Khandpur N, Cediel G, Neri D, Martinez-Steele E, Baraldi LG. Ultra-processed foods: what they are and how to identify them. *Public health nutrition*. 2019 Apr;22(5):936-41.
21. Moubarac JC, Batal M, Louzada ML, Steele EM, Monteiro CA. Consumption of ultra-processed foods predicts diet quality in Canada. *Appetite*. 2017 Jan 1;108:512-20.
22. Rauber F, Louzada ML, Steele EM, Millett C, Monteiro CA, Levy RB. Ultra-processed food consumption and chronic non-communicable diseases-related dietary nutrient profile in the UK (2008–2014). *Nutrients*. 2018 May 9;10(5):587.
23. da Costa Louzada ML, Ricardo CZ, Steele EM, Levy RB, Cannon G, Monteiro CA. The share of ultra-processed foods determines the overall nutritional quality of diets in Brazil. *Public health nutrition*. 2018 Jan;21(1):94-102.
24. Louzada ML, Martins AP, Canella DS, Baraldi LG, Levy RB, Claro RM, Moubarac JC, Cannon G, Monteiro CA. Impact of ultra-processed foods on micronutrient content in the Brazilian diet. *Revista de saude publica*. 2015 Aug 7;49.
25. Louzada ML, Martins AP, Canella DS, Baraldi LG, Levy RB, Claro RM, Moubarac JC, Cannon G, Monteiro CA. Ultra-processed foods and the nutritional dietary profile in Brazil. *Revista de saude publica*. 2015 Jul 10;49.
26. Monteiro CA, Levy RB, Claro RM, de Castro IR, Cannon G. Increasing consumption of ultra-processed foods and likely impact on human health: evidence from Brazil. *Public health nutrition*. 2011 Dec;14(1):5-13.
27. Moubarac JC, Martins AP, Claro RM, Levy RB, Cannon G, Monteiro CA. Consumption of ultra-processed foods and likely impact on human health. Evidence from Canada. *Public health nutrition*. 2013 Dec;16(12):2240-8.
28. Poti JM, Mendez MA, Ng SW, Popkin BM. Is the degree of food processing and convenience linked with the nutritional quality of foods purchased by US households?. *The American journal of clinical nutrition*. 2015 Jun 1;101(6):1251-62.
29. Luiten CM, Steenhuis IH, Eyles H, Mhurchu CN, Waterlander WE. Ultra-processed foods have the worst nutrient profile, yet they are the most available packaged products in a sample of New Zealand supermarkets. *Public health nutrition*. 2016 Feb;19(3):530-8.
30. Beslay M, Srour B, Méjean C, Allès B, Fiolet T, Debras C, Chazelas E, Deschasaux M, Wendeu-Foyet MG, Hercberg S, Galan P. Ultra-processed food intake in association with BMI change and risk of overweight and obesity: A prospective analysis of the French NutriNet-Santé cohort. *PLoS medicine*. 2020 Aug 27;17(8):e1003256.
31. Levy RB, Rauber F, Chang K, Louzada ML, Monteiro CA, Millett C, Vamos EP. Ultra-processed food consumption and type 2 diabetes incidence: a prospective cohort study. *Clinical Nutrition*. 2021 May 1;40(5):3608-14.
32. de Miranda RC, Rauber F, de Moraes MM, Afonso C, Santos C, Rodrigues S, Levy RB. Consumption of ultra-processed foods and non-communicable disease-related nutrient profile in Portuguese adults and elderly (2015–2016): the UPPER project. *British Journal of Nutrition*. 2021 May;125(10):1177-87.
33. Food Safety and Standards (Safe food and balanced diets for School Children) Regulations 2020. Available from: https://www.fssai.gov.in/upload/uploadfiles/files/Gazette_Notification_Safe_Food_Children_07_09_2020.pdf. [Cited 12 May, 2023]
34. Report of Working Group on Addressing Consumption of Foods High in Fat, Salt and Sugar (HFSS) and Promotion of Healthy Snacks in Schools of India. Working Group Constituted by Ministry of Women and Child Development Government of India 2015. Available from: <https://www.nipccd.nic.in/file/reports/hfss.pdf>. [Cited 12 May, 2023]
35. What India Eats, ICMR-NIN 2020. Available from: https://www.millet.res.in/pdf/what_india_eats.pdf. [Cited 12 May, 2023]
36. National Programme for Prevention and Control of Cancer, Diabetes, Cardiovascular Diseases and Stroke (NPCDCS). Directorate General of Health Services, Ministry of Health and Family Welfare, Government of India. Handbook for Counsellors Reducing Risk Factors for Non-communicable Disease. 2017 Sep. Available from:

https://main.mohfw.gov.in/sites/default/files/Handbook%20for%20Counselors%20-%20Reducing%20Risk%20Factors%20for%20NCDs_1.pdf. [Cited 12 May,2023]

37. Recommended Dietary Allowances and Estimated Average Requirements Nutrient Requirements for Indians. A Report of the Expert Group - Indian Council of Medical Research National Institute of Nutrition 2020. .Available from: https://www.nin.res.in/RDA_Full_Report_2020.html. [Cited 12 May, 2023]
38. National Multi-sectoral Action Plan for Prevention and Control of Common NCDs 2017-2022. .Available from: https://main.mohfw.gov.in/sites/default/files/National%20Multisectoral%20Action%20Plan%20%28NMAP%29%20for%20Prevention%20and%20Control%20of%20Common%20NCDs%20%282017-22%29_1.pdf. [Cited 12 May, 2023]
39. Baker P, Machado P, Santos T, Sievert K, Backholer K, Hadjidakou M, Russell C, Huse O, Bell C, Scrinis G, Worsley A. Ultra-processed foods and the nutrition transition: Global, regional and national trends, food systems transformations and political economy drivers. *Obesity Reviews*. 2020 Dec;21(12):e13126.
40. Packaged Food in India. Euromonitor International 2020.
41. India Online Grocery Market Size Report 2021-2028 . Grand view Research.. Available from: <https://www.grandviewresearch.com/industry-analysis/india-online-grocery-market>. [Cited 12 May, 2023]
42. Food Safety and Standards (Food Products Standards and Food Additives) Regulations, 2011. Available from: <https://www.fssai.gov.in/cms/food-safety-and-standards-regulations.php>. [Cited 8 May, 2023]
43. Quinlan M. Considerations for collecting free-lists in the field: examples from ethobotany. *Field methods*. 2005 Aug;17(3):219-34
44. Jain A, Mathur P. Intake of ultra-processed foods among adolescents from low-and middle-income families in Delhi. *Indian Pediatrics*. 2020 Aug;57(8):712-4.
45. National Non-communicable Disease Monitoring Survey (NNMS 2017-18). ICMR-NCDIR. Bengaluru, India. 2021. Available from: chrome-extension://efaidnbmnmmnibpajpcglclefindmkaj/viewer.html?Pdfurl=https%3A%2F%2Fwww.ncdirindia.org%2Fnnms%2Fresources%2Fchapter_4_2_3.pdf&clen=181451&chunk=true. [Cited 18 Jan, 2023]
46. Law C, Green R, Kadiyala S, Shankar B, Knai C, Brown KA, Dangour AD, Cornelsen L. Purchase trends of processed foods and beverages in urban India. *Global food security*. 2019 Dec 1;23:191.
47. Goel S, Kaur T, Gupta M. Increasing proclivity for junk food among overweight adolescent girls in district Kurukshetra, India. *Age (in years)*. 2013;16:17.
48. Bhavani V, Prabhavathy Devi N. Junk and Sink: A Comparative Study on Junk Food Intake among Students of India'. *Shanlax International Journal of Arts, Science and Humanities*. 2020:13-8.
49. Naveenkumar D, Parameshwari S. A Study On Junk Food Consumption and its Effect on Adolescents in Thiruvannamalai District, Tamil Nadu, India. *August 2019 Aug; 8(8): 781-793 [ISSN NO: 1301-2746]*
50. Hiregoudar V, Suresh CM, Singode C, Raghavendra B. Proportion, Patterns, and Determinants of Junk Food Consumption among Adolescent Students. *Annals of Community Health*. 2021 Jul 1;9(2):46-51.
51. Kumari R, Kumari M. Quality aspects of fast foods and their consumption pattern among teenagers of rural-urban region of Sabour block in Bhagalpur district of India. *The Pharma Innovation Journal*. 2020a. 2020;9(4):96-102.
52. Amin T, Choudhary N, Jabeen AN. Study of Fast Food Consumption Pattern in India in Children Aged 16-20 years. . *Intl. J. Food. Ferment. Technol*. 2017 Jun; 7(1): 1-8.
53. Singh M, Mishra S. Fast food consumption pattern and obesity among school going (9-13 year) in Lucknow District. *International Journal of Science and Research*. 2014 Jun;3(6):1672-4.
54. Kotecha PV, Patel SV, Baxi RK, Mazumdar VS, Shobha M, Mehta KG, Mansi D, Ekta M. Dietary pattern of schoolgoing adolescents in urban Baroda, India. *Journal of health, population, and nutrition*. 2013 Dec;31(4):490.
55. Joseph N, Nelliyanil M, Rai S, YP RB, Kotian SM, Ghosh T, Singh M. Fast food consumption pattern and its association with overweight among high school boys in Mangalore city of southern India. *Journal of clinical and diagnostic*

research: JCDR. 2015 May;9(5):LC13.

56. Prabhu NB. Examining fast-food consumption behaviour of students in Manipal University, India. *African Journal of Hospitality, Tourism and Leisure*. 2015;4(2):621-30.
57. Khongrangjem T, Dsouza SM, Prabhu P, Dhang VB, Pari V, Ahirwar SK, Sumit K. A study to assess the knowledge and practice of fast food consumption among Pre-University students in Udipi Taluk, Karnataka, India. *Clinical Epidemiology and Global Health*. 2018 Dec 1;6(4):172-5.
58. Landge JA, Khadkikar GD. Lifestyle and nutritional status of late adolescent in an urban area of Western Maharashtra: cross sectional study. *International Journal of Community Medicine and Public Health*. 2020 Aug;7(8):3027.
59. Iyer ML, Kumar NP. A Study of Breakfast Habits of Urban Indian Consumers. *Research journals of economics and business study*. Res J Economics and Business Studies. 2014 Jun; 3(8): 107-116. [ISSN: 2251-1555].
60. Rathi N, Riddell L, Worsley A. Food consumption patterns of adolescents aged 14–16 years in Kolkata, India. *Nutrition journal*. 2017 Dec;16(1):1-2.
61. Vasan M. Consumers' Preference and Consumption Towards Instant Food Products. *Think India Journal*. 2019 Dec 20;22(14):8333-7.
62. Harrell M, Medina J, Greene-Cramer B, Sharma SV, Arora M. Understanding Eating Behaviors of New Dehli's Youth. *Journal of Applied Research on Children*. 2015;6(2):8.
63. Aguirre A, Borneo MT, El Khori S, Borneo R. Exploring the understanding of the term "ultra-processed foods" by young consumers. *Food Research International*. 2019 Jan 1;115:535-40.
64. Sarmiento-Santos J, Souza MB, Araujo LS, Pion JM, Carvalho RA, Vanin FM. Consumers' Understanding of Ultra-Processed Foods. *Foods*. 2022 May 7;11(9):1359.
65. Mendonça RD, Lopes AC, Pimenta AM, Gea A, Martinez-Gonzalez MA, Bes-Rastrollo M. Ultra-processed food consumption and the incidence of hypertension in a Mediterranean cohort: the Seguimiento Universidad de Navarra Project. *American journal of hypertension*. 2017 Apr 1;30(4):358-66.
66. Srour B, Fezeu LK, Kesse-Guyot E, Allès B, Méjean C, Andrianasolo RM, Chazelas E, Deschasaux M, Hercberg S, Galan P, Monteiro CA. Ultra-processed food intake and risk of cardiovascular disease: prospective cohort study (NutriNet-Santé). *BMJ*. 2019 May 29;365.
67. Srour B, Fezeu LK, Kesse-Guyot E, Alles B, Debras C, Druesne-Pecollo N, Chazelas E, Deschasaux M, Hercberg S, Galan P, Monteiro CA. Ultra-processed food consumption and risk of type 2 diabetes among participants of the NutriNet-Santé prospective cohort. *JAMA internal medicine*. 2020 Feb 1;180(2):283-91.
68. Fiolet T, Srour B, Sellem L, Kesse-Guyot E, Allès B, Méjean C, Deschasaux M, Fassier P, Latino-Martel P, Beslay M, Hercberg S. Consumption of ultra-processed foods and cancer risk: results from NutriNet-Santé prospective cohort. *bmj*. 2018 Feb 14;360.
69. Food Safety and Standards of India - Annual Report 2020-21 . Available from: https://fssai.gov.in/upload/uploadfiles/files/FSSAI_Annual_Report_2020_21_Eng_01_08_2022.pdf. [Cited [cited 7March 2023].
70. Food Safety and Standards of India - Annual Report 2019-20 . Available from: https://fssai.gov.in/upload/uploadfiles/files/FSSAI_Annual_Report_2019_20_English_Hindi.pdf. [Cited 7 March 2023]
71. Food Safety and Standards of India - Annual Report 2018-19 . Available from: https://fssai.gov.in/upload/uploadfiles/files/Annual_Report_2018_19_English_28_10_2020.pdf. [Cited 7 March 2023]
72. Monteiro CA, Cannon G, Moubarac JC, Levy RB, Louzada MLC, Jaime PC. The UN Decade of Nutrition, the NOVA food classification and the trouble with ultra-processing. *Public Health Nutr*. 2017;21(1):5–17.
73. Islam MR, Rahman SM, Rahman MM, Pervin J, Rahman A, Ekström EC. Gender and socio-economic stratification of ultra-processed and deep-fried food consumption among rural adolescents: A cross-sectional study from Bangladesh. *Plos one*. 2022 Jul 28;17(7):e0272275.

74. Pries AM, Filteau S, Ferguson EL. Snack food and beverage consumption and young child nutrition in low-and middle-income countries: A systematic review. Maternal & child nutrition. 2019 Jun;15:e12729.

Footnotes

‡ Thin fat phenotype is the excessive fat accumulation in an individual having normal Body Mass Index (4, 5).

Tables

Table 5 is available in the Supplementary Files section.

Figures




Figure 1

Saliency analysis method for free-listed UPF sub-categories




Figure 2

Flow diagram reporting the screening and selection of studies reporting consumption and availability of UPFs in India

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile1.xlsx](#)
- [AdditionalFile2.xlsx](#)
- [Table5.docx](#)