

Health risk assessment of heavy metals in marine fish caught from the northwest Persian Gulf

Solaleh Torabi

Gonbad Kavous University

Mohammad Gholizadeh (✉ Gholizade_mohammad@yahoo.com)

Gonbad Kavous University

Mahsa Yazarlo

Gorgan University of Agricultural Sciences and Natural Resources

Zahra Riahi

Gonbad Kavous University

Research Article

Keywords: Fish, Heavy metals, Target hazard quotient, Risk assessment, Persian Gulf

Posted Date: August 18th, 2023

DOI: <https://doi.org/10.21203/rs.3.rs-3257359/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Additional Declarations: No competing interests reported.

Version of Record: A version of this preprint was published at Biological Trace Element Research on November 8th, 2023. See the published version at <https://doi.org/10.1007/s12011-023-03946-z>.

Abstract

Due to the presence of metalloids and toxic metals such as cadmium, lead, copper, and zinc in fish tissue, it may endanger health considering the countless benefits of consuming fish, which can harm the human body if consumed in toxic amounts. Therefore, it is vital to monitor the concentration of metals in fish meat to ensure compliance with food safety regulations and, protect the consumer. We considered the levels of Ni, Zn, Cu, Pb, and Cd in 60 marine fish samples (3 species) collected from coastal areas of the northwestern coast of the Persian Gulf and estimated their health risk. Mean concentrations of Ni, Zn, Cu, Pb, and Cd were 1.88 ± 0.07 , 27.16 ± 8.11 , 11.55 ± 4.12 , 14 ± 0.06 , and 0.19 ± 0.03 $\mu\text{g/g}$ wet weight. Estimated average daily intakes (EDIs) for adults and children of Ni, Zn, Cu, Pb, and Cd were 0.89-4.15, 12.89-60.02, 5.47-25.53, 0.54-2.51 and 0.09-0.42 $\mu\text{g/kg}$ bw/day. Considering that there are different levels of certain elements in marine fish in the Persian Gulf, this study provides suggestions for risk management and a scientific basis for food safety assessment.

Introduction

Fish is of interest to consumers due to its polyunsaturated fatty acids, vitamin B, high-quality protein, and other nutrients. However, with the rapid development of industry and urbanization in some estuaries and coastal areas, the problem of environmental pollution has become more serious. These pollutants can accumulate in the muscle tissues of marine organisms and bring potential health risks. In recent years, the association of heavy metals with environmental pollution has caused ecological concerns and environmental severe threats. Marine fish can absorb pollutants such as toxic metals indirectly through food and directly from the water.

For example, Esmaeilbeigi et al. (2023) found the Pb with a mean level of 1.47 ± 0.33 ($\mu\text{g/g}$ ww) in *Atropus atropos* from the Persian Gulf, Iran. Sadeghi et al. (2019) observed the Ni with a mean level of 81.1 $\mu\text{g/g}$ in *Otolithes ruber* from the Oman Sea, Iran. Keshavarzi et al. (2018) described a range level of Cd (1.37 – 3.14 mg/kg) in *Anodontostoma chacunda*, and *Cynoglossus arel* from the Musa estuary Mahshahr harbor, Iran (Persian Gulf). Saadatmand et al. (2022) found the level of Cu (8.62 mg/kg) in *Siganus javus* from the Persian Gulf. All these studies showed the possibility of heavy metal contamination in marine fish.

All ecosystems on earth exhibit metals and metalloids of natural or artificial origin (Foomani et al., 2020; Mirzaei et al., 2020). Both natural and artificial factors can introduce heavy metals into aquatic ecosystems (Gholizadeh et al., 2020). Due to their high toxicity, bio-accumulation, long-term stability, and biomagnification, anthropogenically produced metals in marine environments severely threaten the food chain (Esmaeilbeigi et al., 2023). Over the past ten years, significant progress has been made in the identification of metal pollution sources, in the discovery of metal sinks, and in the decoding of pollution input chronologies. Significant progress has also been made in determining the biotoxicity of various natural and inorganic substances that humans have released into the environment.

Aquatic ecosystems, such as seas and coastlines, are vulnerable to various environmental pollutants, including those released by natural sources and those caused by human activity, such as heavy metals, organic and oil compounds, and herbicides. The most critical human-caused sources of heavy metals are irrigation with wastewater from cities and businesses, solid waste, hydrocarbon fuels, chemical fertilizers, and

pesticides (Doğan-Sağlamtimur *et al.*, 2010; Gholizadeh and Patimar, 2018). Due to their toxicity, trace elements persist in the environment, contaminate food chains, and result in various health issues (Maleki *et al.*, 2020; Parang and Esmaeilbeigi, 2022).

While necessary for the development and growth of organisms, these materials are toxic and dangerous when used in excess. Living things are seriously threatened when exposed to trace elements in the environment regularly (Sadeghi *et al.*, 2019). Fish species, mainly, are among the most significant marine species regarding their ecological and economic importance (Shahifar *et al.*, 2020; Kazemi *et al.*, 2022). Fish consumption has reportedly increased worldwide since 1990 (IFOSY, 2018). To evaluate the potential health risk to consumers and the preservation of the marine environment, it is crucial to measure the concentration of trace elements (such as Hg, Pb, Cd, and As). The accumulation of heavy metals in the bodies of aquatic animals living in polluted marine environments has been the subject of several investigations (Keshavarzi *et al.*, 2018; Banan *et al.*, 2022). Due to their resilience and bio-accumulation, reduced bio-degradability, and potential for biomagnification in the food chain, heavy metals are two assemblages of impurities in the sea environment (Parang and Esmaeilbeigi, 2022).

Some heavy metals are essential to life as trace elements, but the biotoxic effects of many of them on human biochemistry are incredibly problematic. Therefore, it is necessary to thoroughly understand the factors, such as concentrations and oxidation states, that render them harmful and how biotoxicity occurs. Natural and manufactured sources, particularly mining and industrial operations and vehicle exhausts (for lead), are responsible for releasing these metals into the environment. Any metallic element with a moderately high density and is toxic or poisonous even at low concentrations is called a "heavy metal" (Raychaudhuri *et al.*, 2021).

The following substances are classified as heavy metals: lead (Pb), cadmium (Cd), zinc (Zn), arsenic (As), silver (Ag), copper (Cu), chromium (Cr), mercury (Hg), iron (Fe), and platinum group elements (Lee and Stuebing, 1990). While Zn and Cu are among the essential metals for properly operating biological systems, Pb and Hg are generally regarded as the most critical pollution indicators. Through effective biomagnification in the food chain, mercury (Hg) accumulates in both aquatic systems and humans (Yoshino *et al.*, 2020). The term "pollutant" refers to any environmental substance that has unfavorable effects, degrades environmental welfare, decreases human well-being, and may finally result in death. Any substance in the environment with adverse effects that impair the environment's welfare lowers human well-being and may eventually because death called a "pollutant." Since they cannot be degraded or destroyed, heavy metals are persistent environmental contaminants that are naturally present in the earth's crust. They are slightly ingested through food and the air, and over time, they bio-accumulate in the body (Kumar *et al.*, 2020). Most of the time, there are no significant environmental or public health risks from heavy metals filtered through soil and rocks into aquatic systems. At least some anthropogenic sources, like mining, agriculture, and aquaculture, cause metal burdens to rise above natural levels (Ding *et al.*, 2022).

Some heavy metals, including Zn, Fe, Mg, and Ca, have been described as essential to human biology and daily health. Others, however, have been shown to have no known biological significance for human physiology and biochemistry and can even be toxic at low concentrations (such as Pb, Cd, As, and methylated forms of Hg). Dietary intakes must be maintained within administrative limits, even for those of biological

importance, because excesses can cause poison or toxicity (Allowances, 1989). Concern over the quality of food is growing in several global regions. The assurance has prompted studies on the toxicological effects of these elements in food that they are present. Due to their toxicity and accumulation by marine organisms, heavy metals are considered the most critical type of pollution in the aquatic environment (Butnariu, 2022). Metal contamination in water systems typically takes the form of dissolved and suspended particles, which eventually settle and are absorbed by living things. Being at the top of the food chain and able to accumulate high concentrations of a few metals, fish is one of the crucial aquatic organisms in the food chain (Chan *et al.*, 2021). Depending on the level of development, age, and other physiological factors, metal distribution varies between fish species (Yoshino *et al.*, 2020).

The Persian Gulf is regarded as one of the world's most significant closed aquatic ecosystems. It has been impacted by the growing population and industrialization of its bordering countries, and it has discovered a worrying situation due to the pollutants entering the water bodies in the Persian Gulf and the aquatic organisms that live there (Hassanshahian *et al.*, 2020). *Psettodes erumei*, *Sphyraena jello*, and *Sillago sihama* are a few indicator species that can be used to gauge the extent of pollution. The species, found in the northern parts of the Persian Gulf and valued for their marketability and consumer appeal, contribute significantly to the southern Iranian fishing industry. Specialists in nutrition, medicine, and environmental sciences face a significant challenge in monitoring heavy metals because they do not decompose naturally through chemical or biological processes, unlike organic compounds (Keshavarzi *et al.*, 2018; Hassanshahian *et al.*, 2020). Based on this, the current study's objective is to measure heavy metals such as nickel, zinc, cadmium, copper, and lead in three commercial fish species of the Persian Gulf: *Psettodes erumei*, *Sphyraena jello*, and *Sillago sihama*. After determining the health risks associated with the metals, the study will then determine the maximum amounts that can be consumed.

Materials and methods

In the spring of 2022, trawl nets were used to collect samples from the four coastal communities of Asaluyeh, Dayyer, Deylam, and Bandar Bushehr in the Bushehr province (Fig. 1). Sixty aquatic Specimens (20 of each species), *Psettodes erumei*, *Sphyraena jello* and *Sillago sihama*, were studied after placing them in acid-washed plastics and frozen until analysis. In polyethylene bags, samples were returned to the Persian Gulf University lab and kept frozen (-20°C) until analysis.

Table 1
Characteristics of sampling stations in Bushehr province

Site	Geographic position	Characteristics
Deylam	30°03'30 N, 50°08'20 E	Deylam port with a sandy beach with various coastal activities (tourism and fishing activities)
Genaveh	29°35'19 N, 50°29'43 E	
Bushehr	28°55'06 N, 50°48'30 E	The province's seat has high population density, a high rate of municipal sewage and the traffic of commercial and fishing vessels, no agricultural pollution, tourism activities, several swimming beaches and is adjacent to the Bushehr nuclear power plant.
Mond	28°00'08 N, 51°22'21 E	
Dayyer	27°49'54 N, 51°56'13 E	Iran's largest fishing port, the leading supplier of fish and shrimp, the traffic area of commercial vessels and municipal sewage and the nearest city to Asaluyeh and a relatively suitable index for comparison with the relatively clean area and the highly contaminated area of Asaluyeh
Asaluyeh	27°28'01 N, 52°36'35 E	The most industrialized and polluted part of the province in the presence of enormous gas and petrochemical plants, a platform of massive vessels carrying chemical and petrochemical materials, gas extraction, and desalination plants related to the mentioned industries

In the tissues of the species, the pollution caused by high concentrations of zinc, lead, nickel, copper, and cadmium was examined. In the spring of 2022, in the province of Bushehr, sampling was done randomly among the fish caught and prepared for the market from a particular area of the pier. The samples were cleaned with distilled and city water, and the total length and weight were determined using a ruler and a scale, respectively, with an accuracy of 0.1 cm and 0.01g. Following the biopsy, the muscle tissue was isolated, and cleaned with distilled water.

A 50 ml Erlenmeyer flask was filled with 3 g fish muscle tissue wet weight. The samples were given a dose of 4 mg of nitric acid (65 percent), and they were then placed under the hood at room temperature for at least an hour to begin the initial digestion process. The samples were mixed with 1.5 ml of perchloric acid (70%) and then placed on a hot plate (sand bath) at 140°C for six hours to finish the digestion process. The samples were digested and then left to cool at room temperature. The samples were then diluted with deionized water to a volume of 25 ml, the solutions were filtered through Whatman filter paper (40 microns), and the samples were stored in a polyethylene container in a frigid environment (4°C) until analysis (Nollet, 2004). The level of heavy metals in fish muscle was determined using an ICP-OES instrument made by Liberty RL.

Based on the national per capita consumption (average daily fish consumption for adults of 31.92 grams), the following equations estimate the CR, EDI, and THQ as well as the carcinogenic risk potential of heavy metals introduced into the human body as a result of fish consumption was calculated daily per person (IFOSY, 2018).

$$EDI = \frac{C \times FIRD}{BW}$$

In this context, EDI stands for estimated daily intake of metals by the body, C for identified metal concentration in consumed food, and FIRD for determined daily food intake rate in grams. The target hazard quotient (THQ), used to express non-toxic effects, is the ratio between the amount of exposure metals' amount and their reference dose. If this rate is less than one, no risk can be seen.

Consumers' health will be at risk if this ratio equals or exceeds one (IFOSY, 2018). In this case, EFr is the Exposure Frequency (365 days per year), ED_{tot} is the Exposure Duration, total (72 years), FIR is the food intake rate (about 31.92 grams per person per day for fish), C is the concentration of Heavy metal in the studied food (mg/g), RfDo is Reference Dose, oral (mg/kg/day), BW_a Weight Body, adult (70 kg) and ATN is Averaging time, noncarcinogens (365 days/year times the number of years of exposure is about 72 years) (IFOSY, 2018).

$$THQ = \frac{EFr \times ED_{tot} \times FIR \times C}{(RfDo \times BW_a \times ATN) \times 10^{-3}}$$

The pollutant carcinogenic risk index (CR) is obtained from the following relationship. In this regard, EDI is the body's estimated daily intake of metals, and CSF is the cancer slope factor (Tepanosyan *et al.*, 2017).

$$CR = EDI \times CSF$$

A cancer risk index indicates a range of anticipated potential risks for agents that cause cancer between 10^{-6} (lifetime risk of developing cancer 1 in 1,000,000) and 10^{-4} (lifetime risk of developing cancer 1 in 10,000). As a result, chemicals that contain risk factors (less than 10^{-6}) are not regarded as chemicals of concern (Tepanosyan *et al.*, 2017). Based on the U.S. Environmental Protection Agency (USEPA) established standard defaults, the relevant calculations were completed. Nickel compounds are categorized as class I carcinogenic elements by the International Agency for Research on Cancer (IARC).

Statistical analysis

The kolmogoraph-smirnov normality test was used to check the normality of the data Levene-test was used to evaluate the homogeneity of variance. One way analysis of variance (ANOVA) test with a significance level of 5% was used to analyze data between elements in the sampling location for the statistical analysis of the concentration of heavy metals in fish muscle tissue (El-Moselhy *et al.*, 2014). The software Excel 2013 was also used to create the graphs and tables. The resulting data were compared to relevant national and international standards.

Results

Ten specimens of each species were examined in the current study. The average total length (cm) of *Sillago sihama* (20.57 ± 2.50), *Sphyræna jello* (34.71 ± 5.96), and *Psettodes erumei* (44.91 ± 15.46) were based on biometric data. Table 2 lists the biometric findings and traits of the investigated species, including diet and habitat.

Table 2
Characteristics of the studied species (standard deviation $\pm$ mean)

Species	Number of species	The total weight (g)	Total length (cm)	Diet	Habitat
<i>P. erumei</i>	20	00.72 $\pm$ 908. 508	44.15 $\pm$ 91.46	Carnivorous	Benthic
<i>S. jello</i>	20	14.03 $\pm$ 224.33	71.96 $\pm$ 34.5	Carnivorous	Pelagic
<i>S. sihama</i>	20	21.07 $\pm$ 82.13	57.50 $\pm$ 20.2	Planktivorous	Demersal

Table 3 displays the typical levels of the studied species' muscle tissue's heavy metal content (copper, nickel, lead, zinc, and cadmium). Based on the obtained results, the highest concentration of metals was observed respectively in the muscle tissue of *P. erumei* > *S. sihama* > *S. jello*.

Table 3
The average concentration of heavy metals in the muscle tissue of the studied species based on milligrams per gram of wet weight (mean $\pm$ standard deviation)

Species	Cd	Pb	Cu	Zn	Ni
<i>P. erumei</i>	18.06 $\pm$ 0.0	84.48 $\pm$ 0.0	60.34 $\pm$ 2.1	32.69 $\pm$ 2.3	80.3 $\pm$ 0.0
<i>S. jello</i>	4.02 $\pm$ 0.0	2.01 $\pm$ 0.0	90.04 $\pm$ 0.0	10.08 $\pm$ 6.10	54.58 $\pm$ 1.0
<i>S. sihama</i>	5.02 $\pm$ 0.0	9.01 $\pm$ 0.0	53.12 $\pm$ 0.0	60.44 $\pm$ 30.8	30.14 $\pm$ 0.0

The estimated daily intake (EDI) of the elements under investigation was calculated for the current study and is shown in Table 4. According to the findings, the rate of estimated daily intake (EDI) and estimated weekly intake (EWI) for nickel, lead, and cadmium metals in all studied species are higher than the limit of provisional tolerable daily intake (PTDI) and provisional tolerable weekly intake (PTWI) in an adult. Estimated daily intake (EDI) and estimated weekly intake (EWI) for copper and zinc were both below the limit of the provisional tolerable weekly intake (PTWI). In addition, the daily and weekly intake rate of heavy metals in *Psettodes erumei* was as follows: copper > zinc > nickel > lead > cadmium. In *Sphyræna jello* fish, daily and weekly intake of heavy metals is in order: copper > nickel > zinc > cadmium > lead. The daily and weekly intake rates of heavy metals for *Sillago sihama* were copper > zinc > nickel > cadmium > lead.

Table 4
Estimation of daily and weekly intake of heavy metals due to consumption of studied fishes by consumers (adults and children)

Species	metals	EWI ^e (adults- children)	EDI ^d (adults- children)	PTDI ^c	PTWI ^b	PTWI ^a
<i>P. erumei</i>	Ni	12.41–2.66	0.38–1.77	0.012	2.45	0.035
	Cu	35.83–7.68	5.1–12.10	0.5	245	3.5
	Zn	40.19–8.61	5.1 -74.23	1	490	7
	Pb	13.2–03.79	1.0-86.40	0.0035	1.75	0.025
	Cd	2.0 -82.60	0.0-40.09	0.001	0.49	0.007
<i>S. jello</i>	Ni	23.5 -88.12	3.0 -41.73	0.012	2.45	0.035
	Cu	94.20 -42.23	13.2 -49.89	0.5	245	3.5
	Zn	13.2 -87.97	1.0 -98.42	1	490	7
	Pb	0.0 -25.05	0.0–4.01	0.0035	1.75	0.025
	Cd	0.54 – 0.12	0.0–8.02	0.001	0.49	0.007
<i>S. sihama</i>	Ni	4.64–0.99	0.0 -66.14	0.012	2.45	0.035
	Cu	101.44-473.38	14.49–67.63	0.5	245	3.5
	Zn	8.1-13.74	1.16–0.25	1	490	7
	Pb	0.23 – 0.05	0.03 – 0.01	0.0035	1.75	0.025
	Cd	0.81 – 0.17	0.12 – 0.02	0.001	0.49	0.007
a = Provisional tolerable weekly intake (PTWI), in µg/week/kg of body weight						
b = Provisional tolerable weekly intake (PTWI) for an adult with an average body weight (70 kg), in µg/g/week for a 70 kg person						
c = Provisional tolerable daily intake (PTDI), in µg/day for a 70 kg person						
d = estimated daily intake (EDI), in µg/day for a 70-kg person and a 15-kg child						
e = estimated weekly intake (EWI), in µg/day for a 70 kg adult and a 15 kg child						

Based on the concentration of heavy metals, the calculation of the maximum amount of fish that can be eaten daily (kg/day) and the maximum amount of fish that can be eaten monthly (serving/month) is shown for adults and children of the studied species (Table 5). *Perumei* were related to zinc metal (9.07 kg/day), *S. jello* (17.21 kg/day), and *S. sihama* (18.67 kg/day) to lead metal, which is the highest permissible limit of fish consumption for two age groups of adults and children. In addition, *S. sihama* and both adult and child populations had higher permissible lead consumption rates (2503.14 servings per month) than other metals.

Table 5
The amount permissible limit (kg per day) and the permissible rate of consumption
(number of servings per month) of heavy metals in studied fishes

Species	CR _{mm} (number of servings per month)		CR _{lim} (kg per day)		
	children	adults	Children	adults	metals
<i>P. erumei</i>	50.16	243.08	0.37	1.75	Ni
	260.55	1215.90	1.94	9.07	Cu
	30.97	144.52	00.23	1.08	Zn
	9.56	44.59	0.07	0.33	Pb
	11.05	51.58	0.08	0.38	Cd
<i>S. jello</i>	26.07	121.6	0.19	0.91	Ni
	98.87	461.39	0.74	3.44	Cu
	89.73	418.74	0.67	3.12	Zn
	494.62	2308.23	3.69	17.21	Pb
	57.47	268.19	0.43	2.00	Cd
<i>S. sihama</i>	134.10	625.79	4.67	1.00	Ni
	19.72	92.03	0.15	0.69	Cu
	153.03	714.16	1.14	5.33	Zn
	536.39	2503.14	4.00	18.67	Pb
	38.31	178.80	0.29	1.33	Cd

Table 6 provides the target hazard quotient (THQ) and total target hazard quotient (TTHQ) of nickel, zinc, copper, lead, and cadmium in the studied fishes for consumers (children and adults). The following vital presumptions were considered when calculating the risk index: a) the amount of imported metal equals the amount absorbed in the body. b) Pollutants are unaffected by cooking. c) Iranians live an average of 72 years; d) adults weigh an average of 70 kg, while children weigh an average of 15 kg. The risk index is less than one for each of the three species of metals and fish studied, which means there is no risk to consumer health. Additionally, the combined risk of the heavy metals used in *S. jello* and *S. sihama* is less than one, indicating no health risk to the consumers.

Table 6

Target hazard quotient (THQ) of non-cancerous diseases and hazard index (HI) of studied heavy metals, in adults (a) and children (c), based on consumption of aquatic species muscle

Species	THQ)target hazard quotient)										Hazard index (HI)	
	Cd		Pb		Cu		Zn		Ni			
	C	a	c	a	c	a	c	a	c	a	c	a
<i>P. erumei</i>	0.4	0.09	0.47	0.1	0.14	0.031	0.02	0.004	0.09	0.02	1.12	0.24
<i>S. jello</i>	0.08	0.02	0.009	0.002	0.05	0.01	0.05	0.01	0.17	0.04	0.35	0.08
<i>S. sihama</i>	0.12	0.03	0.008	0.002	0.03	0.006	0.23	0.05	0.03	0.01	0.41	0.09

Table 7

Comparison of the concentration of studied metals ($\mu\text{g/g}$) in the muscle of fishes with international standards ($\mu\text{g/g}$).

Species	Metals	Mean	Minimum	Maximum	USEPA	FDA	WHO
<i>P. erumei</i>	Ni	80.3 $\pm$ 0.0	0.39	1.83	0.5	-	60–80
	Cu	60.34 $\pm$ 2.1	1.32	4.84	120	-	10
	Zn	32.69 $\pm$ 2.0	1.44	3.11	120	-	100
	Pb	84.48 $\pm$ 0.0	0.34	1.64	0.5	5	0.4
	Cd	18.06 $\pm$ 0.0	0.13	0.28	2	2	0.2
<i>S. jello</i>	Ni	54.58 $\pm$ 1.0	0.2	4.18	0.5	-	60–80
	Cu	90.04 $\pm$ 0.0	0.86	0.94	120	-	10
	Zn	10.08 $\pm$ 6.10	2.26	11.74	120	-	100
	Pb	0.2.01 $\pm$ 0.0	0.013	0.023	0.5	5	0.4
	Cd	04.02 $\pm$ 0.0	0.02	0.05	2	2	0.2
<i>S. sihama</i>	Ni	30.14 $\pm$ 0.0	0.19	0.51	0.5	-	60–80
	Cu	53.12 $\pm$ 0.0	0.4	0.66	120	-	10
	Zn	60.44 $\pm$ 30.8	19.9	40.1	120	-	100
	Pb	09.01 $\pm$ 0.0	0.07	0.1	0.5	5	0.4
	Cd	05.02 $\pm$ 0.0	0.04	0.08	2	2	0.2

Discussion

Economic activities like refining crude oil, making aluminum and zinc, building and repairing ships, and loading and unloading various minerals and chemicals, have all increased recently (Gholizadeh and Patimar,

2018). These have destroyed the delicate water ecosystem and increased pollution, which has developed along Iran's southern coast. The harmful effects affecting marine ecosystems, including aquatic life and humans, make identifying, measuring, and tracking heavy metal elements essential (Bozorgpour *et al.*, 2017). From six main ports in Bushehr province, Persian Gulf, 60 pieces of fish were examined in the spring of 2022. *P. erumei*, *S. sihama*, and *S. jello* fish were tested for their ability to absorb the five heavy metals (lead, cadmium, nickel, copper, and zinc) required by the World Health Organization and the Food and Agriculture Organization of the United Nations.

There are many different ways that heavy metals end up in aquatic environments, including atmospheric precipitation, waste material discharge, unintentional leakage, water discharge from ship balance tanks, discharge of industrial, agricultural, and domestic sewage, and soil erosion (Bashir *et al.*, 2020) and fish may take in some heavy metals from the environment via the food chain or water (Zaynab *et al.*, 2020). The results of the ANOVA analysis revealed a significant difference in the number of metals present in the fish muscle tissue in the sampling area ($p < 0.05$). Applying different management and environmental conditions, sewage discharge, and aquaculture activities in the investigated regions may cause a significant difference in the concentration of heavy elements in other species and areas. Dural *et al.* (2007) demonstrated that there are differences in the concentration of heavy metals in the bodies of aquatic organisms in the various areas (the Persian Gulf, the Egyptian Gulf, the Scandinavian Gulf, and the California wetlands) as a result of multiple environmental factors, including temperature, light, and human activity. The current study used the spring season to examine 60 pieces of fish. Shahri and Velayatzadeh (2017), the length and the total weight of 48 fish were used to study the impact of seasons on the buildup of metals in the muscle of *Acanthopagrus latus* and *Platycephalus indicus* in the Oman Sea. In the present study, the concentration of metals was observed in the muscle of *P. erumei* > *S. sihama* > *S. jello*.

P. erumei (9.07 kg per day), *S. jello* (17.21 kg per day), and *S. sihama* (18.67 kg per day) have the highest daily limits for zinc consumption for two age groups of fish, adults and children. Additionally, compared to other metals, the acceptable lead consumption rate for *S. sihama* (2503.14 monthly servings) was higher. Cu and Zn are crucial for most animals' growth, cell metabolism, and survival. As a result, their necessity can be attributed to the generally high levels of these metals (Khansari *et al.*, 2005). A crucial component of the human glucose tolerance factor, Cr is a fundamental micronutrient trace metal with a pattern similar to Zn and Ni (Karahana, 2023). However, its excess may cause diabetes or contribute to diabetes. Cd may not be a significant metal for organisms, but it is highly toxic to wildlife and humans. The risk index is less than one for all metals and the three studied fish species, indicating no risk to consumer health. There is no risk to consumers' health because the combined risk index for heavy metals in *S. jello* and *S. sihama* is less than one. Most investigated organisms are benthic, with a 5 to 80 meters depth range, and can feed close to the bed. Being carnivores, they frequently eat fish and crustaceans for food (Norouzi, 2020); therefore, if the amount of heavy metal accumulation in the fish was more significant than that in the muscle of *S. jello* fish, then it is one of the species that is placed above the food web. The bioaccumulation of heavy metals occurs in the muscle of the fishes. According to an assessment of the risk of heavy metal accumulation in green tiger shrimp (*Penaeus semisulcaus*) in the waters of Bushehr provinces, the amount of metal accumulation was 0.40 and 25.43 µg/g, respectively. The calculated risk index for adults and children was less than 1, indicating no risk to the health of shrimp consumers in terms of heavy metals (Hossain *et al.*, 2022).

The body's ability to control the concentrations of some metals and the fact that body size and biochemical components have little to no bearing on variability may help to explain why there is no relationship between some metal concentrations and length (Hossain *et al.*, 2022). The findings indicated a correlation between the concentration of copper, zinc, and cadmium elements and total fish length in the study area; as the fish's total length increases, so does the concentration of the elements, as mentioned earlier in fish muscle tissue. Dadolahi *et al.*, (2008) found that the amount of lead and cadmium metals in the muscle and gill tissue of *Arabibarbus grypus* have a direct linear relationship with total length and weight. Shirvani *et al.*, (2016) demonstrated that there is a statistically significant relationship between the concentration of metals and body size by examining the number of heavy metals in the muscle and liver tissues of two species of *Terapon jarbua* and *Sillago sihama* on the coast of the southern basin of Qeshm Island, in other words, the intensity of pollution increases with an increase in fish length and body weight.

The combined effects of the rainy season's (July to September) dilution of water salinity and spawning, which affect metal levels by lowering concentrations, may be to blame for the loss of metals in soft tissue. Under specific environmental or physiological conditions, increasing the levels of some organisms in various tissues may have a detrimental effect on the normal operations of various organs and tissues (Seebacher and Franklin, 2012). Esmaeilbeigi *et al.* (2023) mentioned that although metallothionein's metal binding is steady, it is also dynamic due to quick metal exchanges within and between proteins. The latter property enables metal exchange within and between metallothionein molecules and between metallothionein and donor or acceptor ligands. Fish organs generally respond poorly to heavy metal poisoning. Due to its crucial function in human nutrition and the need to ensure its safety for consumption, fish muscle tissue was examined in this study. Although it is clear that muscles are not a biologically active site for the transfer and accumulation of metals, there is a severe risk to human health in polluted aquatic habitats where the concentration of metals in fish muscles may be higher than is safe for human consumption. The World Health Organization (WHO) and the US Food and Drug Administration (FDA) set acceptable standards for heavy metals, and the values of the results of this study were compared with those standards (Table 7). The comparison of the results revealed that the lead concentration in the muscle tissue of *P. erumei* is higher than the reference standards. Other metal concentrations in fish are below the World Health Organization's permissible limit. The potential hazard quotient (THQ) was used to calculate the risk of dietary intake of heavy metals from seafood consumption (Table 6).

Tahsini *et al.* (2018) assessed the concentration of heavy metals in the fish breeding ponds of Sanandaj city and the risk of their ingestion in rainbow trout. Additionally, Tahsini *et al.* (2018) determined that adults should consume 888.7 g of food daily, while children should consume 177.7 g. The findings demonstrated that fish muscle tissue has an average concentration of iron, nickel, zinc, copper, and magnesium metals that is lower than the permitted limit according to international standards. The existence of numerous industries along the coasts, the discharge of industrial and urban wastewater, which contains all kinds of heavy metals, and causes an increase in the concentration of this metal in this area are the reasons why the maximum concentration of nickel metal in fish muscle tissue is higher than the USEPA standard (Barakat, 2011). Additionally, the THQ index results for the studied region's fish muscle tissue were calculated to be less than one. The THQ is less than one in the risk assessment, which means there will be no adverse health effects for the consumer and no threat to the consumer's natives. Fish with a considerable length had a higher cancer

slope index in the metals under investigation. Other findings demonstrated that the amount of CR in element exposure carries a negligible risk of carcinogenesis.

Conclusion

The concentration of nickel, zinc, copper, lead, and cadmium metals in the muscle of fish with different length classes show a notable difference and a positive correlation, according to the study's findings and measurement of metals in the studied fishes in Bushehr Province. With increasing body length, the concentration of these metals also rises. The concentration of nickel metal was higher than the permissible limit compared to the standard of the World Health Organization, which could be problematic for consumers. Fish consumption poses no risks to consumers, according to the findings of research into THQ levels in adults. Finally, it is suggested to offer management solutions to improve the current conditions after considering the results and realizing the Bushehr coastal area's relative pollution and ecological significance.

Declarations

Data Availability Statement: All the data are within the article.

Competing Interests: None

Funding: This study is supported financially by Gonbad Kavous University (project number: 205413), Gonbad-e Kavus, Golestan province, Iran.

Ethical approval: In the current study, fish were caught by fisherman, not specifically for the project.

Consent to participate: This is review study so there is no need for consent to participate.

Consent to Publish: The participant has consented to the submission of the case report to the journal.

Author Contributions: All authors contributed to the study conception and design. **S. T:** Conceptualization, Methodology, Writing - Original Draft, Review & Editing, Visualization; **M. Gh:** Supervision, Writing - Original Draft, Review & Editing, Project administration, Funding acquisition; **M. Y:** Investigation, Resources, Data Curation; **Z. R:** Writing - Original Draft

References

1. Allowances, RD, 1989. Recommended dietary allowances. National Research Council-National Academy Press: Washington, DC, USA.
2. Bashir, I., Lone, F.A., Bhat, R.A., Mir, S.A., Dar, Z.A. and Dar, S.A., 2020. Concerns and threats of contamination on aquatic ecosystems. *Bioremediation and biotechnology: sustainable approaches to pollution degradation*, pp.1-26.
3. Barakat, M.A., 2011. New trends in removing heavy metals from industrial wastewater. *Arabian journal of chemistry*, 4(4), pp.361-377.

4. Bozorgpour, R., Omaraee, B. and Asadi, M.V.Z., 2017. Study and Analysis of Obstacles and Challenges Facing Ship-Repair Industry in Iran. *Open Journal of Marine Science*, 7(4), pp.485-493.
5. Butnariu, M., 2022. Heavy metals as pollutants in the aquatic Black Sea ecosystem. In *Bacterial Fish Diseases* (pp. 31-57). Academic Press.
6. Chan, W.S., Routh, J., Luo, C., Dario, M., Miao, Y., Luo, D. and Wei, L., 2021. Metal accumulations in aquatic organisms and health risks in an acid mine-affected site in South China. *Environmental geochemistry and health*, 43, pp.4415-4440.
7. Dadolahi, S., Nabavi, S., khiror, N. 2008. The relationship of some biometric specifications with the accumulation of heavy metals in muscle tissue and freshwater fish gills in the Arvand River. *Iranian Fisheries Scientific Journal*, 17(4): 27-34(In Persian).
8. Ding, C., Chen, J., Zhu, F., Chai, L., Lin, Z., Zhang, K. and Shi, Y., 2022. Biological Toxicity of Heavy Metal (loid) s in Natural Environments: From Microbes to Humans. *Frontiers in Environmental Science*, p.681.
9. Doğan-Sağlamtimur, N. and Kumbur, H., 2010. Metals (Hg, Pb, Cu, and Zn) bioaccumulation in sediment, fish, and human scalp hair: a case study from the city of Mersin along the southern coast of Turkey. *Biological trace element research*, 136, pp.55-70.
10. Dural, M., Goksu, M. Z. L., Ozak, A. A. 2007. Investigation of heavy metal levels in economically important fish species from the Tuzla lagoon. *Food Chemistry*, 102(1): 415-421.
11. El-Moselhy, K. M., Othman, A. I., Abd El-Azem, H., & El-Metwally, M. E. A. 2014. Bioaccumulation of heavy metals in some tissues of fish in the Red Sea, Egypt. *Egyptian journal of basic and applied sciences*, 1(2), 97-105.
12. Esmaeilbeigi, M., Kazemi, A., Gholizadeh, M., & Rezaeiye, R. D. 2023. Microplastics and heavy metals contamination in *Atropus atropus* and associated health risk assessment in the northwest of the Persian Gulf, Iran. *Regional Studies in Marine Science*, 57, 102750.
13. Foomani, A., Gholizadeh, M., Harsij, M. and Salavatian, M., 2020. River health assessment using macroinvertebrates and water quality parameters: A case of the Shanbeh-Bazar River, Anzali Wetland, Iran. *Iranian Journal of Fisheries Sciences*, 19(5), pp.2274-2292.
14. Gholizadeh, M., and Patimar, R. 2018. Ecological risk assessment of heavy metals in surface sediments from the Gorgan Bay, Caspian Sea. *Marine pollution bulletin*, 1: 137: 662-7.
15. Gholizadeh, M., Mohammadzadeh, B. and Kazemi, A., 2022. Human health risk assessment via the consumption of *platycephalus indicus* in the Persian Gulf, Iran. *Pollution*, 8(3), pp.740-750.
16. Hassanshahian, M., Amirinejad, N. and Askarinejad Behzadi, M., 2020. Crude oil pollution and biodegradation at the Persian Gulf: A comprehensive and review study. *Journal of Environmental Health Science and Engineering*, 18, pp.1415-1435.
17. Hossain, M.B., Bhuiyan, N.Z., Kasem, A., Hossain, M., Sultana, S., Nur, A.A.U., Yu, J., Albeshr, M.F. and Arai, T., 2022. Heavy Metals in Four Marine Fish and Shrimp Species from a Subtropical Coastal Area: Accumulation and Consumer Health Risk Assessment. *Biology*, 11(12), p.1780.
18. IFOSY, 2018. Fish consumption per capita in Iran Iranian Fisheries Organization Statistical Yearbook. 22-23.

19. Karahan, F., 2023. Evaluation of trace element and heavy metal levels of some ethnobotanically important medicinal plants used as remedies in Southern Turkey in terms of human health risk. *Biological Trace Element Research*, 201(1), pp.493-513.
20. Kazemi, A., Esmailbeigi, M., Sahebi, Z. and Shooshtari, S.J., 2022. Hydrochemical evaluation of groundwater quality and human health risk assessment of trace elements in the largest mining district of South Khorasan, Eastern Iran. *Environmental Science and Pollution Research*, 29(54), pp.81804-81829.
21. Keshavarzi, B., Hassanaghaei, M., Moore, F., Rastegari Mehr, M., Soltanian, S., Lahijanzadeh, A.R., Sorooshian, A., 2018. Heavy metal contamination and health risk assessment in three commercial fish species in the Persian Gulf. *Marine Pollution Bulletin*, 129, 245–252.
<http://dx.doi.org/10.1016/j.marpolbul.2018.02.032>
22. Khansari, F.E., Ghazi-Khansari, M., Abdollahi, M. 2005. Heavy metals content of canned tuna fish. *Food chemistry*, 93(2):293-6.
23. Kumar, A., Kumar, A., MMS, C.P., Chaturvedi, A.K., Shabnam, A.A., Subrahmanyam, G., Mondal, R., Gupta, D.K., Malyan, S.K., Kumar, S.S. and A. Khan, S., 2020. Lead toxicity: health hazards, influence on food chain, and sustainable remediation approaches. *International journal of environmental research and public health*, 17(7), p.2179.
24. Lee, Y. H., and Stuebing, R. B. 1990. Heavy metal contamination in the River Toad, *Bufo juxtasper* (Inger), near a copper mine in East Malaysia. *Bulletin of Environmental Contamination and Toxicology*, (USA). 1:45(2).
25. Maleki, P., Rahman, P., Jafariyan, H., Salmanmahiny, A., Ghorbani, R., Gholizadeh, M. and Harsij, M., 2020. The risk assessment of water pollution in the Gorgan Bay catchment using the WRASTIC index. *Environmental Nanotechnology, Monitoring & Management*, 14, p.100393.
26. Mirzaei, M., Marofi, S., Solgi, E., Abbasi, M., Karimi, R. and Riyahi Bakhtyari, H.R., 2020. Ecological and health risks of soil and grape heavy metals in long-term fertilized vineyards (Chaharmahal and Bakhtiari province of Iran). *Environmental geochemistry and health*, 42, pp.27-43.
27. Nollet, L. M. L. Hand book of food analysis. CRC Press. USA. 2004.
28. Norouzi, M., 2020. Evaluating the accumulation and consumption hazard risk of heavy metals in the fish muscles of species living in the waters of the Persian Gulf, Iran. *Pollution*, 6(4), pp.849-862.
29. Parang, H., Esmailbeigi, M., 2022. Total mercury concentration in the muscle of four mostly consumed fish and associated human health risks for fish-ermen and non-fishermen families in the Anzali Wetland, Southern Caspian Sea. *Regional Studies in Marine Science*, 52, 102270.
<http://dx.doi.org/10.1016/j.rsma.2022.102270>.
30. Raychaudhuri, S.S., Pramanick, P., Talukder, P. and Basak, A., 2021. Polyamines, metallothioneins, and phytochelatins—Natural defense of plants to mitigate heavy metals. *Studies in Natural Products Chemistry*, 69, pp.227-261.
31. Saadatmand, M., Dadolahi-Sohrab, A., Tavani, M. B., Khazaei, S. H., and Saadatmand, F. 2022. Monitoring heavy metal contamination on the Iranian coasts of the Persian Gulf using biological indicators: risk assessment for the consumers. *Environmental Monitoring and Assessment*, 194(2), 83.
32. Sadeghi, P., Loghmani, M. and Afza, E., 2019. Trace element concentrations, ecological and health risk assessment in sediment and marine fish *Otolithes ruber* in Oman Sea, Iran. *Marine pollution bulletin*, 140,

pp.248-254.

33. Seebacher, F. and Franklin, C.E., 2012. Determining environmental causes of biological effects: the need for a mechanistic physiological dimension in conservation biology. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1596), pp.1607-1614.
34. Shahifar, R., Patimar, R., Fazli, H., Raeisi, H., Gholizadeh, M. and Jafaryan, H., 2020. Growth and mortality parameters of Caspian kutum, *Rutilus kutum*, in southern Caspian Sea. *International Journal of Aquatic Biology*, 8(1), pp.56-65.
35. Shahri, E., Velayatzadeh, M. 2017. The Effect of Cold and Warm Seasons on Accumulation of Nickel, Cadmium and Lead in Muscle of *Acanthopagrus latus* and *Platycephalus indicus* from Oman Sea (Chabahar). *Iranian Journal of Marine Science and Technology Research*, 12(1): 10-21(In Persian).
36. Shirvani, M. 2016. Measurement of lead and cadmium metals in fish tissues (*Sillago sihama*) and (*Terapon jarbua*) in the southern part of Qeshm Island. *Journal of Marine Science and Technology*, 11(1); 1-9(In Persian).
37. Tahsini, H., Alizadeh, M., Goilian, H. 2018. Evaluation of the concentration of heavy metals and its consumption risk in rainbow trout in fish ponds (Case study: Kamyaran and Sanandaj). *Journal of Environmental Health Engineering*, 6(2): 187-196 (In Persian).
38. Tepanosyan, G., Maghakyan, N., Sahakyan, L., and Saghatelyan, A. 2017. Heavy Metals Pollution Levels and Children Health Risk Assessment of Yerevan Kindergartens Soils. *Ecotoxicology and Environmental Safety*, 142, 257–265. doi:10.1016/j.ecoenv.2017.04.013
39. Yoshino, K., Mori, K., Kanaya, G., Kojima, S., Henmi, Y., Matsuyama, A. and Yamamoto, M., 2020. Food sources are more important than biomagnification on mercury bioaccumulation in marine fishes. *Environmental pollution*, 262, p.113982.
40. Zaynab, M., Al-Yahyai, R., Ameen, A., Sharif, Y., Ali, L., Fatima, M., Khan, K.A. and Li, S., 2022. Health and environmental effects of heavy metals. *Journal of King Saud University-Science*, 34(1), p.101653.

Figures




Figure 1

Location map of the studied area (map of Iran, location of sampling stations in Bushehr province)