

Table 2. Demographic and clinical characteristics of the subset of patients included in the visual field analysis.

Variable	Entire cohort	NTG	HTG
Eyes/patients (n)	229/139	144/85	85/54
Baseline Age (years)	63.7 (± 10.4)	63.2 (± 10.7)	64.5 (± 10.0)
Sex, female/male (n)	86/53	54/31	32/22
Laterality, right/left (n)	118/111	74/70	44/41
CCT, mean ($\pm$ SD), micron	539 (± 42)	535 (± 39)	545 (± 46)
Baseline MD, median (IQR), dB	-3.97 (-1.86 to -8.04)	-4.94 (-1.96 to -9.54)	-2.80 (-1.09 to -6.18)
MD rate, mean ($\pm$ SD), dB/year	-0.59 (± 0.33)	-0.61 (± 0.35)	-0.57 (± 0.29)
*Baseline IOP, median (IQR), mmHg	15 (13 – 18)	14 (12 – 16)	18 (15 – 21)
*Mean IOP, mean ($\pm$ SD), mmHg	14.9 (± 2.8)	13.6 (± 1.9)	17.0 (± 2.7)
*Peak IOP, median (IQR), mmHg	19 (16 – 21)	17 (15 – 19)	22 (20 – 25)
*SD IOP, median (IQR), mmHg	2.2 (1.7 – 2.9)	2.0 (1.7 – 2.4)	2.6 (1.9 – 3.6)
Pre-treatment IOP, median (IQR), mmHg	18 (15 – 22)	16 (14 – 18)	24 (22 – 28)
*IOP reduction, median (IQR), %	20.7 (8.3– 31.4)	17.1 (4.5– 23.7)	31.9 (20.1– 41.7)
Length of follow-up, median (IQR), years	5.7 (4.6 – 6.9)	5.0 (4.2 – 6.4)	6.4 (5.4 – 8.6)
Number of visual fields, median (IQR)	10 (8 – 12)	10 (9 – 12)	9 (8 – 11)

*The baseline and final IOP in the series corresponded to the date of the first and last visual field included in the analyses.

CCT: central corneal thickness; dB: decibel; IOP: intraocular pressure; IQR: interquartile range; HTG: high-tension glaucoma; MD: mean deviation; NTG: normal-tension glaucoma; SD: standard deviation