

Supplementary Table 1. Longitudinal changes in problematic behaviors before and after treatment (A) overall and by treatment group (B, C, and D).

(A) OVERALL			
Problematic Behavior	Before	After	p value
Anxiety	82,3	69,2	<0,001
Cutting	77,7	25,4	<0,001
Prior hospitalization	42,3	1,5	<0,001
Familiar disfunction	72,4	48,8	<0,001
Eating disorder	16,2	2,3	<0,001
Problematic alcohol/substance use	20,8	10	0,001
(B) Group A			
Problematic Behavior	Before	After	p value
Anxiety	84,1	68,2	0,02
Cutting	81,8	15,9	<0,001
Prior hospitalization	52,3	4,5	<0,001
Familiar disfunction	68,2	59,1	0,063
Eating disorder	15,9	2,3	0,016
Problematic alcohol/substance use	11,4	0	<0,05
(C) Group B			
Problematic Behavior	Before	After	p value
Anxiety	80,4	62,7	0,006
Cutting	78,4	13,7	<0,001
Prior hospitalization	33,3	0	<0,001
Familiar disfunction	80	28	<0,001
Eating disorder	21,6	2	0,001
Problematic alcohol/substance use	17,6	7,8	0,09
(D) Group C			
Problematic Behavior	Before	After	p value
Anxiety	82,9	80	0,5
Cutting	71,4	54,3	0,016
Prior hospitalization	42,9	0	<0,001
Familiar disfunction	66,7	66,7	0,75
Eating disorder	8,6	2,9	0,25
Problematic alcohol/substance use	37,1	25,7	0,063