

3.0	Study Design	Describe the study design.	STORMS	Observational (Case-Control, Cohort, Cross-sectional survey, etc.) or Experimental (Randomized controlled trial, Non-randomized controlled trial, etc.). For a brief description of common study designs, see: DOI: 10.11613/BM.2014.022 If applicable, describe any blinding (e.g., single or double-blinding) used in the course of the study.	Yes	INTRODUCTION section, paragraph 4 Blinding does not apply to this study.
3.1	Participants	State what the population of interest is and the method by which participants are sampled from that population. Include relevant information on the physiological state of the subjects or stage in the life history of disease under study when participants were sampled.	STORMS	Examples of the population of interest could be: adults with no chronic health conditions, adults with type II diabetes, newborns, etc. This is the total population to whom the study is hoped to be generalizable. The sampling method describes how potential participants were selected from that population. If the participants are from a substudy of a larger study, provide a brief description of that study and cite that study. Clearly state how cases and controls are defined. An example of relevant physiological state might be pre/post menopausal for a vaginal microbiome study; examples of stage in the life history of disease could be whether	Yes	It Is fully described under the METHODS section, under the "Subject recruitment" subsection.

				specimens were collected during active or dormant disease or before or after treatment.		
3.2	Geographic location	State the geographic region(s) where participants were sampled from.	mlxS: geographic location (country and/or sea, region)	Geographic coordinates can be reported to prevent potential ambiguities if necessary.	Yes	Study participants were collected from two medical institutes in Japan as described in the METHODS section under the "Subject recruitment" subsection.

3.3	Relevant Dates	State the start and end dates for recruitment, follow-up, and data collection.	STORMS	Recruitment is the period in which participants are recruited for the study. In longitudinal studies, follow-up is the date range in which participants are asked to complete a specific assessment. Finally, data collection is the total period in which data is being collected from participants including during initial recruitment through all follow-ups.	Yes	Sampling points are provided in a Supplementary Figure (e.g., “Month 0” as the initial time point, and the following points are indicated as months passed.)
3.4	Eligibility criteria	List any criteria for inclusion and exclusion of recruited participants.	Modified STROBE	Among potential recruited participants, how were some chosen and others not? This could include criteria such as sex, diet, age, health status, or BMI. If there is a primary and validation sample, describe inclusion/exclusion criteria for each.	Yes	It is fully described under the METHODS section, under the “Subject recruitment” subsection.
3.5	Antibiotics Usage	List what is known about antibiotics usage before or during sample collection.	STORMS	If participants were excluded due to current or recent antibiotics usage, state this here. Other factors (e.g. proton pump inhibitors, probiotics, etc.) that may influence the microbiome should also be described as well.	NA	

3.6	Analytic sample size	Explain how the final analytic sample size was calculated, including the number of cases and controls if relevant, and reasons for dropout at each stage of the study. This should include the number of individuals in whom microbiome sequencing was attempted and the number in whom microbiome sequencing was successful.	STORMS	Consider use of a flow diagram (see template at https://stormsmicrobiome.org/figures) . Also state sample size in abstract. If power analysis was used to calculate sample size, describe those calculations.	Yes	The analytic sample size is written in the RESULTS section depending on the samples relevant for each case vs. control cross-sectional analysis.
3.7	Longitudinal Studies	For longitudinal studies, state how many follow-ups were conducted, describe sample size at follow-up by group or condition, and discuss any loss to follow-up.	STORMS	If there is loss to follow-up, discuss the likelihood that drop-out is associated with exposures, treatments, or outcomes of interest.	Yes	The number of samples longitudinally collected from each participant is written in the RESULTS section.
3.8	Matching	For matched studies, give matching criteria.	Modified STROBE	"Matched" refers to matching between comparable study participants as cases and controls or exposed / unexposed. Indicate whether participants were individual or frequency matched and in what ratio were they matched (e.g. 1 case to 1 control).	NA	
3.9	Ethics	State the name of the institutional review board that approved the study and protocols, protocol number	STORMS		Yes	It is stated in the

		and date of approval, and procedures for obtaining informed consent from participants.				“Regulatory compliance” subsection in the METHODS section.
4.0	Laboratory methods	State the laboratory/center where laboratory work was done.	STORMS	Provide a reference to complete lab protocols if previously published elsewhere such as on protocols.io. Note any modifications of lab protocols and the reason for protocol modifications.	Yes	The detailed lab protocols reference is cited.
4.1	Specimen collection	State the body site(s) sampled from and how specimens were collected.	MIxS: sample collection device or method; host body site	Use terms from the Uber-anatomy Ontology (https://www.ebi.ac.uk/ols/ontologies/uberon) to describe body sites in a standardized format.	Yes	Only fecal samples were collected as specimens.
4.2	Shipping	Describe how samples were stored and shipped to the laboratory.	STORMS	Include length of time from collection to receipt by the lab and if temperature control was used during shipping.	Yes	The METHODS section is under the “Fecal sample collection and DNA extraction” subsection.
4.3	Storage	Describe how the laboratory stored samples, including time between collection and storage and any preservation buffers or refrigeration used.	STORMS	State where each procedure or lot of samples was done if not all in the same place. Include reagent/lot/catalogue #s for storage buffers.	Yes	The METHODS section is under the “Fecal sample

						collection and DNA extraction" subsection.
4.4	DNA extraction	Provide DNA extraction method, including kit and version if relevant.	MIxS: nucleic acid extraction	If any DNA quantification methods were used prior to DNA amplification or at the pooling step of library preparation, state so here.	Yes	The METHODS section is under the "Fecal sample collection and DNA extraction" subsection.
4.5	Human DNA sequence depletion or microbial DNA enrichment	Describe whether human DNA sequence depletion or enrichment of microbial or viral DNA was performed.	STORMS		Yes	METHODS section, under the "Quality control" subsection.
4.6	Primer selection	Provide primer selection and DNA amplification methods as well as variable region sequenced (if applicable).	MIxS: pcr primers		Yes	METHODS section, under the "Quality control" subsection.
4.7	Positive Controls	Describe any positive controls (mock communities) if used.	STORMS	If used, should be deposited under guidance provided in the 8. X items.	NA	
4.8	Negative Controls	Describe any negative controls if used.	STORMS	If used, should be deposited under guidance provided in the 8.X items.	NA	

4.9	Contaminant mitigation and identification	Provide any laboratory or computational methods used to control for or identify microbiome contamination from the environment, reagents, or laboratory.	STORMS	Includes filtering of reagents and other steps to minimize contamination. It is relevant to state whether the specimens of interest have low microbial load, which makes contamination especially relevant.	Yes	METHODS section, under the “Quality control” subsection.
4.10	Replication	Describe any biological or technical replicates included in the sequencing, including which steps were replicated between them.	STORMS	Replication may be biological (redundant biological specimens) or technical (aliquots taken at different stages of analysis) and used in extraction, sequencing, preprocessing, and/or data analysis.	NA	
4.11	Sequencing strategy	Major divisions of strategy, such as shotgun or amplicon sequencing.	MIxS: sequencing method	For amplicon sequencing (for example, 16S variable region), state the region selected. State the model of sequencer used.	Yes	Shotgun sequencing
4.12	Sequencing methods	State whether experimental quantification was used (QMP/cell count based, spike-in based) or whether relative abundance methods were applied.	STORMS	These include read length, sequencing depth per sample (average and minimum), whether reads are paired, and other parameters.	Yes	Relative abundance methods were applied in METHODS under the “Taxonomic profiling” and “Functional profiling” subsections.
4.13	Batch effects	Detail any blocking or randomization used in study design to avoid confounding batches with exposures or outcomes. Discuss any likely sources of batch effects, if known.	STORMS	Sources of batch effects include sample collection, storage, library preparation, and sequencing and are commonly unavoidable in all but the smallest of studies.	No	A possible batch effect is the samplings from two different

						hospitals. Although not shown in the manuscript, we validated that this did not significantly affect the overall microbial structures.
4.14	Metatranscriptomics	Detail whether any mRNA enrichment was performed and whether/how retrotranscription was performed prior to sequencing. Provide size range of isolated transcripts. Describe whether the sequencing library was stranded or not. Provide details on sequencing methods and platforms.	STORMS	Provide details on any internal standards which may have been used as well as parameters and versions of any software or databases used.	NA	
4.15	Metaproteomics	Detail which protease was used for digestion. Provide details on proteomic methods and platforms (e.g. LC-MS/MS, instrument type, column type, mass range, resolution, scan speed, maximum injection time, isolation window, normalised collision energy, and resolution).	STORMS	Provide details on any internal standards which may have been used as well as parameters and versions of any software or databases used.	NA	
4.16	Metabolomics	Specify the analytic method used (such as nuclear magnetic resonance spectroscopy or mass spectrometry). For mass spectrometry, detail which fractions were obtained (polar and/or non-polar) and how these were analyzed. Provide details on metabolomics methods and platforms (e.g. derivatization, instrument type, injection type, column type and instrument settings).	STORMS	Provide details on any internal standards which may have been used as well as parameters and versions of any software or databases used.	Yes	METHODS section, under the “Metabolome analysis” subsection.

5.0	Data sources/ measurement	For each non-microbiome variable, including the health condition, intervention, or other variable of interest, state how it was defined, how it was measured or collected, and any transformations applied to the variable prior to analysis.	MIxS: host disease status	State any sources of potential bias in measurements, for example multiple interviewers or measurement instruments, and whether these potential biases were assessed or accounted for in study design. Use terms from a standardized ontology such as the Experimental Factor Ontology (https://www.ebi.ac.uk/efo/) to describe variables of interest in a standardized format.	Yes	The METHODS section is under the “Measurement/Evaluation of CRC (colorectal cancer) risk factors” subsection.
6.0	Research design for causal inference	Discuss any potential for confounding by variables that may influence both the outcome and exposure of interest. State any variables controlled for and the rationale for controlling for them.	STORMS	For causal inference, this item refers to describing the assumptions that would be required to draw causal inferences from observational data. See Vujkovic-Cvijin, I., Sklar, J., Jiang, L. et al. Host variables confound gut microbiota studies of human disease. <i>Nature</i> 587, 448–454 (2020). https://doi.org/10.1038/s41586-020-2881-9 for more details on confounding in observational microbiome studies. For example, hypothesized confounders may be controlled for by multivariable adjustment. Consider using a directed acyclic graph (DAG) to describe your causal model and justify any variables controlled for.	Yes	Confounding factors were tested using PERMANOVA, and the results are provided in a supplementary figure.

				DAGs can be made using www.dagitty.net .		
6.1	Selection bias	Discuss the potential for selection or survival bias.	STORMS	Selection bias can occur when some members of the target study population are more likely to be included in the study/final analytic sample than others. Some examples include survival bias (where part of the target study population is more likely to die before they can be studied), convenience sampling (where members of the target study population are not selected at random), and loss to follow-up (when the probability of dropping out is related to one of the things being studied).	No	We collected samples from as many FAP patients as possible for our effort. We did not select samples but used all samples for microbiome analysis.
7.0	Bioinformatic and Statistical Methods	Describe any transformations to quantitative variables used in analyses (e.g. use of percentages instead of counts, normalization, rarefaction, categorization).	STORMS	If a variable is analyzed using different transformations, state rationale for the transformation and for each analysis which version of the variable is used.	Yes	METHODS under “Taxonomic profiling” and “Functional profiling” subsections.

				In case of any complex or multistep transformations, give enumerated instructions for reproducing those transformations.		
7.1	Quality Control	Describe any methods to identify or filter low quality reads or samples.	MIxS: sequence quality check	If samples were excluded based on quality or read depth, list the criteria used, the number of samples excluded, and the final sample size after quality control.	Yes	METHODS section, under the “Quality control” subsection.
7.2	Sequence analysis	Describe any taxonomic, functional profiling, or other sequence analysis performed.	MIxS: feature prediction; similarity search method		Yes	METHODS under “Taxonomic profiling” and “Functional profiling” subsections.
7.3	Statistical methods	Describe all statistical methods.	Modified STROBE	Describe any statistical tests used, exploratory data analysis performed, dimension reduction methods/unsupervised analysis, alpha/beta metrics, and/or methods for adjusting for measurement bias. If multiple statistical methods are possible, discuss why the methods used were selected. If a multiple hypothesis testing correction method was used, describe the type of correction used.	Yes	METHODS section, under the “Statistical tests” subsection.

				State which taxonomic levels are analyzed.		
7.4	Longitudinal analysis	If the study is longitudinal, include a section that explicitly states what analysis methods were used (if any) to account for grouping of measurements by individual or patterns over time.	STORMS		Yes	RESULTS section under "Drastic microbial shift correlated with increased CRC risk on longitudinal analysis on FAP cases".
7.5	Subgroup analysis	Describe any methods used to examine subgroups and interactions.	STROBE		Yes	INTRODUCTI ON section, paragraph 4.
7.6	Missing data	Explain how missing data were addressed.	STROBE	"Missing data" refers to participant measurements such as covariates, exposures, outcomes, or time points that should have been collected but were not, not to zeros in taxonomic abundance tables or data points not applicable to that observation.	Yes	Missing data is marked as "-" in a supplementar y table.
7.7	Sensitivity analyses	Describe any sensitivity analyses.	STROBE		NA	

7.8	Findings	State criteria used to select findings for reporting.	STORMS	For example, false discovery rate with total number of tests, effect size threshold, significance threshold, microbes of interest.	Yes	METHOD section, under the "Statistical analysis" subsection.
7.9	Software	Cite all software (including read mapping software) and databases (including any used for taxonomic reference or annotating amplicons, if applicable) used. Include version numbers.	Modified STREGA	Installed packages, add-ons or libraries should be stated and cited in addition to the software used. All parameters employed that differ from the default of that software/version should be provided. This is in addition to, not a replacement for, publishing of code as outlined in the section Reproducible Research.	Yes	METHODS section, under each of the metagenomic analyses subsections.
8.0	Reproducible research	Make a statement about whether and how others can reproduce the reported analysis.	STORMS	Any protected information that has been excluded or provided under controlled access should be listed along with any relevant data access procedures. "On request from authors" is not sufficiently detailed; formal data access procedures and conditions should be defined. If data are unavailable, state so clearly. Consider using a specialized rubric for reproducible research (such as: https://mbio.asm.org/content/9/3/e00525-18.short).	Yes	METHODS section, under "Data and code availability".

				Consider preregistering the study protocol (such as on osf.io or https://plos.org/open-science/preregistration/).		
8.1	Raw data access	State where raw data may be accessed including demultiplexing information.	STORMS	Robust, long-term databases such as those hosted by NCBI and EBI are preferred. If using a private repository, provide rationale.	Yes	METHODS section, under “Data and code availability”.
8.2	Processed data access	State where processed data may be accessed.	STORMS	Unfiltered data should be provided. Robust, long-term databases such as those hosted by NCBI and EBI-EMBL are preferred. Repositories like zenodo (https://zenodo.org/) or publisso (https://www.publisso.de/en/working-for-you/doi-service/) can be used to provide a DOI and long-term storage for processed datasets, even those which cannot be published openly.	Yes	Microbiome profiles and metabolite profiles are provided in supplementary tables.

8.3	Participant data access	State where individual participant data such as demographics and other covariates may be accessed, and how they can be matched to the microbiome data.	STORMS	If re-categorized, transformed, or otherwise derived variables were used in the analysis, these variables or code for deriving them should be provided. Examples of how participant data can be matched to microbiome data are: using the same set of anonymized identifiers, or using different anonymized identifiers but providing a map. Provided data should be sufficient to independently replicate the current analysis.	Yes	It is provided in a supplementary table.
8.4	Source code access	State where code may be accessed.	STORMS	If a standard or formalized workflow was employed, reference it here.	NA	
8.5	Full results	Provide full results of all analyses, in computer-readable format, in supplementary materials.	STORMS	For example, any fold-changes, p-values, or FDR values calculated, provided as a spreadsheet. Use a machine-readable, plain-text format such as csv or tsv.	Yes	It is provided in supplementary tables.

Results

9.0	Descriptive data	Give characteristics of study participants (e.g. dietary, demographic, clinical, social) and information on exposures and potential confounders.	STROBE	Typically reported in a table included in the paper or as a supplementary table. Indicate number of participants with missing data for each variable of interest. This includes environmental and lifestyle factors that may affect the relationship between the microbiome and the condition of interest. Participant diet and medication use should be summarized, if known. At minimum, the age and sex of all participants should be summarized.	Yes	RESULTS section, paragraphs 1 and 2, supplementary tables.
10.0	Microbiome data	Report descriptive findings for microbiome analyses with all applicable outcomes and covariates.	STORMS	This includes measures of diversity as well as relative abundances. These descriptive findings should be reported both for the sample overall and for individual groups.	Yes	Relative abundances are visualized for top 10 most abundant genera. Measurement of diversity result is provided in supplementary figures.
10.1	Taxonomy	Identify taxonomy using standardized taxon classifications that are sufficient to uniquely identify taxa.	STORMS	If not using full taxonomic hierarchy, make sure it is clear whether names stated are species, genera, family, etc.	Yes	Taxonomy names and IDs are stated as defined by

				Italicize genus/species pairs. Consult journal guidelines or standardized references on taxonomic nomenclature. For instance, https://wwwnc.cdc.gov/eid/page/scientific-nomenclature		mOTU version 2.6.1.
10.2	Differential abundance	Report results of differential abundance analysis by the variable of interest and (if applicable) by time, clearly indicating the direction of change and total number of taxa tested.	STORMS	If there are more than two groups, include omnibus (multigroup) test results if applicable to the research question. If applicable, reported effect sizes should include a measure of uncertainty such as the confidence interval.	Yes	Differentially abundant species and genes are visualized in boxplots with statistical significance.
10.3	Other data types	Report other data analyzed--e.g. metabolic function, functional potential, MAG assembly, and RNAseq.	STORMS		Yes	METHOD section, under the “Functional profiling” subsection.
10.4	Other statistical analysis	Report any statistical data analysis not covered above.	STORMS	This could include subgroup analysis, sensitivity analyses, and cluster analysis. Visualizations should be easily interpretable and colorblind-friendly. The caption and/or main text should provide a detailed description of visualizations for visually-impaired readers.	Yes	The METHOD section under “Microbial community structure analysis” fully described other statistical analyses such as the

						Dirichlet multinomial mixture model.
--	--	--	--	--	--	--------------------------------------

Discussion

11.0	Key results	Summarise key results with reference to study objectives	STROBE			
12.0	Interpretation	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence.	STROBE	Define or clarify any subjective terms such as "dominant," "dysbiosis," and similar words used in interpretation of results. When interpreting the findings, consider how the interpretation of the findings may be summarized or quoted for the general public such as in press releases or news articles. If causal language is used in the interpretation (such as "alters," "affects," "results in," "causes," or "impacts"), assumptions made for causal inference should be explicitly stated as part of 6.0 and 13.0. Distinguish between function potential (ie inferred from metagenomics) and observed activity (ie metatranscriptomic, metabolomic, proteomic) if discussing microbial function.	Yes	DISCUSSION section, paragraphs 2-5.

13.0	Limitations	Discuss limitations of the study, taking into account sources of potential bias or imprecision.	STROBE	Also consider limitations resulting from the methods (especially novel methods), the study design, and the sample size.	Yes	DISCUSSION section, paragraph 5
13.1	Bias	Discuss any potential for bias to influence study findings.	STORMS	May include sampling method, representativeness of study participants, or potential confounding.	No	We collected samples from as many FAP patients as possible for our effort. We did not select samples but used all samples for microbiome analysis.
13.2	Generalizability	Discuss the generalisability (external validity) of the study results	STROBE	To what populations or other settings do you expect the conclusions to generalize?	Yes	DISCUSSION section, paragraph 6.
14.0	Ongoing/future work	Describe potential future research or ongoing research based on the study's findings.	STORMS		Yes	DISCUSSION section, paragraph 6.

Other information

15.0	Funding	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	STROBE		Yes	Acknowledgements section
15.1	Acknowledgements	Include acknowledgements of those who contributed to the research but did not meet criteria for authorship.	STORMS	For general guidelines on authorship, see http://www.icmje.org and https://www.elsevier.com/authors/journal-authors/policies-and-ethics/credit-author-statement	Yes	Acknowledgements section

15.2	Conflicts of Interest	Include a conflicts of interest statement.	STORMS		Yes	Reporting Summary
16.0	Supplements	Indicate where supplements may be accessed and what materials they contain.	STORMS		Yes	The video file is included in the INTRODUCTION section, paragraph 3.
17.0	Supplementary data	Provide supplementary data files of results with for all taxa and all outcome variables analyzed. Indicate the taxonomic level of all taxa.	STORMS	Depending on the analysis performed, examples of the supplemental results included could be mean relative abundance, differential abundance, raw p-value, multiple hypothesis testing-adjusted p-values, and standard error. All discussed taxa should include the taxonomic level (e.g. class, order, genus).	Yes	

