

Supplementary material

The prognostic impact of treatment centralization in patients with testicular germ cell tumors: Analysis of Hospital-Based Cancer Registry data in Japan

Shuhei Suzuki¹, Yoshiyuki Nagumo¹, Shuya Kandori^{1,*}, Kousuke Kojo¹, Satoshi Nitta¹,
Ichiro Chihara¹, Masanobu Shiga¹, Atsushi Ikeda¹, Takashi Kawahara¹, Akio Hoshi¹,
Hiromitsu Negoro¹, Mathis J Bryan,² Ayako Okuyama^{3,4}, Takahiro Higashi³, Hiroyuki
Nishiyama¹

Authors' academic affiliations:

1) Department of Urology, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki
305-8575, Japan

2) International Medical Center, University of Tsukuba Affiliated Hospital, 2-1-1,
Amakubo, Tsukuba, Ibaraki 305-8576, Japan

3) Center for Cancer Registries, Center for Cancer Control and Information Service,
National Cancer Center, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

4) Graduate School of Nursing, St Luke's International University, 10-1 Akashicho,
Chuo-ku, Tokyo 104-0044, Japan

1 ***Corresponding author:** Dr. Shuya Kandori, Department of Urology, University of
2 Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan.
3 Tel.: +81-29-853-3223, Fax: +81-29-853-8854. Email: shuya79@md.tsukuba.ac.jp
4

1 Table S1 Characteristics of patients with TGCT


2

	Seminoma	Non-seminoma	P-value
No. of patients	1262	505	
Age group			< 0.001
20s	146 (11.6%)	175 (34.7%)	
30s	446 (35.3%)	184 (36.4%)	
40s	404 (32.0%)	91 (18.0%)	
50s-	266 (21.1%)	55 (10.9%)	
Summary stage			< 0.001
StageI	1053 (83.4%)	294 (58.2%)	
StageII	120 (9.5%)	64 (12.7%)	
StageIII	89 (7.1%)	147 (29.1%)	
Institutional care volume			0.52
High-volume	684 (54.2%)	283 (56.0%)	
Low-volume	578 (45.8%)	222 (44.0%)	

3

4

1 **Fig.S1 The overall survival of patients with seminomas (A) and non-seminomas (B),**
2 **stratified by summary stage**



3
4

1 **Fig. S2 The overall survival of stage II-III seminoma (A) and non-seminoma (B)**
2 **patients, stratified by institutional care volume**

