

Table 1. Demographic Characteristics and Transplant Data of Recipients Divided into Patients with Non-Melanoma Skin Cancer (NMSC) and without NMSC.

Characteristics	All (n = 100)	With NMSC (n=39)	Without NMSC (n = 61)	P Value
Age (years). (mean $\pm$ SD)	57 $\pm$ 10.8	60 $\pm$ 7.4	55.8 $\pm$ 10.8	0.01
Gender. M; n (%)	60 (63)	28 (72)	35 (57.4)	0.11
Race (C/NC)	83/17	36/3	47/14	0.10
Smoker; n (%)	23(23)	9 (23.1)	14 (23)	1
Associated Comorbidities, n (%)				
Hypertension	83 (83)	33(84.6)	50 (82)	1.0
Diabetes mellitus	44 (44)	13 (33.3)	32 (52.5)	0.09
Transplant				
Transplantation time (years)	9 $\pm$ 5.6	10 $\pm$ 6.5	8 $\pm$ 4.8	0.04
Deceased donor; n (%)	81 (81)	30 (77)	51 (83.6)	0.44
DCE; n (%)	44 (44)	19 (48.7)	25 (41)	0.53
Living donor; n (%)	18 (18)	8 (20.5)	10 (16.4)	0.60
Type of NMSC, n (%)				
CEC	23 (23)	23 (59)	-	-
CBC	16 (16)	16 (41)	-	-
Actinic keratosis	33 (33)	23 (59)	10 (16.4)	0.0001
Family history of NMSC	15 (15)	11 (28)	4 (6.6)	0.002
Number of lesions, n (%)				
1	24 (24)	24 (61.5)	-	-
More 1	15 (15)	15 (38.5)	-	-
Fitzpatrick, n (%)				
Up to III	82 (82)	35 (89.7)	47 (77)	0.11
IV or higher	18 (18)	4 (10.3)	14 (23)	0.12
Immunosuppressive Drugs, n (%)				
With Azathioprine	25 (25)	15 (38.4)	28 (45.9)	0.53
With CNIs	43 (43)	12 (30.7)	14 (22.9)	0.40
With MTORi	26 (26)	11 (28)	14 (23)	0.70
Induction. [n (%)]				
Thymoglobulin	16 (16)	15 (38.4)	7 (11.5)	0.002
Basiliximab	68 (68)	22 (56.4)	46 (75.4)	0.07

NMSC: Non-Melanoma Skin Cancer; C: Caucasian; NC: Non-Caucasian; ECD: Extended Criteria Donor; SCC: Squamous Cell Carcinoma; BCC: Basal Cell Carcinoma; CNIs: Calcineurin Inhibitors; MTORi: mTOR Inhibitors. Significant P values are highlighted in bold.