

Table 5. Factors Associated with the Exclusive Breast-feeding Practice of study participants in Yeka Sub-city, Addis Ababa, Ethiopia, 2022.

Characteristics	Options	EBF Practice		COR (95%CI)	AOR (95%CI)
		Not practiced	Practiced		
Educational status of husband	No	26	29	1	1
	Yes	223	398	1.600(0.919-2.785)	1.704(1.003-3.251) **
Educational status of the mother	No	17	30	1	1
	Yes	233	402	0.978(0.528-1.811)	0.752(0.366-1.544)
EBF counseling during ANC (n=669)	No	81	133	1	1
	Yes	148	307	2.879(1.851, 4.476)	2.764(1.522,4.998)*
PNC counseling	No	84	108	1	1
	Yes	167	324	1.509(1.073-2.122) *	1.471(1.036-2.089) **
Husband support on EBF	No	54	65	1	1
	Yes	197	367	1.548(1.037-2.309) *	1.353(0.883-2.074)
Birth order	First	102	219	1	1
	Second	94	136	0.674(0.474-0.959) *	0.717(0.501-1.026)
	Three and above	54	77	0.664(0.436-1.011)	0.649(0.424-0.994) **
Mode of delivery	SVD	188	303	1	1
	C/S	62	129	1.291(0.907-1.838)	1.273(0.886-1.827)

Significant at multivariate analysis=**

AOR=Adjusted odds Ratio

COR=Crude odds Ratio