

Additional file 5

Differences of symptom groups in the sub-scores of counseling and treatment service preferences

(I) Comparing symptom groups	(J) Comparing symptom groups	Mean difference (I–J)	<i>p</i>	95% CI	
				LL	UL
Professional and personal confidants					
Without	PPD	.051	.134	-.012	.118
PPD or PAD	PAD	-.089	.283	-.241	.083
	Comorbid ^a	.169	.002*	.071	.272
PPD	Without ^b	-.051	.134	-.120	.014
	PAD	-.140	.111	-.306	.035
	Comorbid ^a	.119	.043*	.002	.240
PAD	Without ^b	.089	.283	-.092	.247
	PPD	.140	.111	-.045	.313
	Comorbid ^a	.258	.008*	.063	.445
Comorbid	Without ^b	-.169	.002*	-.270	-.073
PPD and PAD	PPD	-.119	.043*	-.233	-.008
	PAD	-.258	.008*	-.438	-.070
Psychotherapeutic services					
Without	PPD	.112	.022*	.015	.208
PPD or PAD	PAD	-.193	.126	-.439	.059
	Comorbid ^a	-.073	.343	-.220	.080
PPD	Without ^b	-.112	.022*	-.208	-.017
	PAD	-.306	.020*	-.562	-.039
	Comorbid ^a	-.185	.037*	-.355	.007
PAD	Without ^b	.193	.126	-.059	.438
	PPD	.306	.020*	.038	.563
	Comorbid ^a	.121	.416	-.170	.403
Comorbid	Without ^b	.073	.343	-.080	.220
PPD and PAD	PPD	.185	.037*	.006	.357
	PAD	-.121	.416	-.417	.179

Note. Bootstrap results are based on 5,000 bootstrap samples. CI = bias-corrected and accelerated bootstrap interval, LL = lower limit, UL = upper limit.

^a PPD and PAD, ^b PPD or PAD.

* *p* < .05, two-tailed.