

Supplementary materials to

Amine modification of calcium phosphates by low-pressure plasma for bone regeneration

Joe Kodama¹, Anjar Anggraini Harumningtyas^{2,3}, Tomoko Ito², Miroslav Michlicek^{4,5}, Satoshi Sugimoto², Yuichiro Ukon¹, Junichi Kushioka¹, Rintaro Okada¹, Takashi Kamatani¹, Kunihiro Hashimoto¹, Daisuke Tateiwa¹, Hiroyuki Tsukazaki¹, Shinichi Nakagawa¹, Shota Takenaka¹, Takahiro Makino¹, Yusuke Sakai¹, David Necas⁴, Lenka Zajickova^{4,5}, Satoshi Hamaguchi^{2,*}, Takashi Kaito^{1,*}

¹ Department of Orthopaedic Surgery, Osaka University Graduate School of Medicine, Osaka, 565-0871, Japan

² Center for Atomic and Molecular Technologies, Graduate School of Engineering, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871 Japan

³ Center for Accelerator Science and Technology, National Nuclear Energy Agency of Indonesia (BATAN), Jalan Babarsari Kotak Pos 6101 ykbb Yogyakarta 55281, Indonesia

⁴ Plasma Technologies, CEITEC – Central European Institute of Technology, Masaryk University, Kamenice 5, Brno 62500, Czech Republic

⁵ Department of Physical Electronics, Faculty of Science, Masaryk University, Kotlářská 2, Brno 61137, Czech Republic

*Corresponding authors

Takashi Kaito, MD, PhD.

Department of Orthopaedic Surgery, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan.

Tel: +81-6-6879-3552

Fax: +81-6-6879-3559

E-mail: takashikaito@ort.med.osaka-u.ac.jp

Satoshi Hamaguchi, PhD.


Center for Atomic and Molecular Technologies (CAMT),
Graduate School of Engineering, Osaka University,
2-1 Yamadaoka, Suita, Osaka 565-0871, Japan.

TEL: +81-6-6879-7913 (06-6879-7913)

FAX: +81-6-6879-7916 (06-6879-7916)

E-mail: hamaguch@ppl.eng.osaka-u.ac.jp

Supplementary Figure S1. Plasma treatment enhanced the hydrophilicity, cell infiltration in vitro, and denser tissue generation was observed at 3 weeks in vivo.



Supplementary Movie S2 is provided as a separate file.

Supplementary Figure S3. Photo of the implantation of porous disks into rat calvarial bone defects. Plasma treated disks showed better blood infiltration.

