

**SOCIAL AND BIOLOGICAL IMPACT OF OIL PALM (*Elaeis guineensis* Jacq.)
PLANTATIONS IN THE EASTERN BRAZILIAN AMAZON**

Biological and Conservation

Adriana P. Manhães ^{a*}, Flávia Rocha ^b, Tatiana Souza ^a, Karoline Marques ^{a,c}, Leandro Juen ^c, Luciano Montag ^c, Bruno Coutinho ^a

^a Conservation International, Rio de Janeiro, Rio de Janeiro, Brazil

^b Laboratory of Studies on Biodiversity and Sustainability, Federal Rural University of Rio de Janeiro, Seropédica, Rio de Janeiro, Brazil

^c Ecology and Conservation Laboratory, Federal University of Pará, Belém, Pará, Brazil


* correspondence author

E-mail address: adriana.manhaes@arvorescer.com

Full present address: Federal University of Rio de Janeiro, Av. Carlos Chagas Filho, #373, 21941-902, Rio de Janeiro, Rio de Janeiro, Brazil

Online Resource 1

Location of the Agropalma company where the seven taxonomic groups of biodiversity were sampled, municipality of Tailândia, Pará (Brazil). Plots sampling the palm oil plantation streams are represented by circles while forest streams are represented by triangles.



Source: Juen et al. (2016).

References

Juen L, Cunha EJ, Carvalho FG, et al (2016) Effects of Oil Palm Plantations on the Habitat Structure and Biota of Streams in Eastern Amazon: Effect of Oil Palm on the Structure of Stream. *River Res Appl* 32:2081–2094. <https://doi.org/10.1002/rra.3050>