

Table

Table 1 Oligonucleotide primer sequences used for quantitative real-time PCR.

Gene	Primer sequence (5'-3')
<i>HIF-1α</i>	F: GAACGTCGAAAAGAAAAGTCTCG R: CCTTATCAAGATGCGAACTCACA
<i>IL-1β</i>	F: ATGATGGCTTATTACAGTGGCAA R: GTCGGAGATTCGTAGCTGGA
<i>TNFα</i>	F: CCATGTTGTAGCAAACCCTCAAG R: AAGAGGACCTGGGAGTAGATGAG
<i>FADD</i>	F: GTGGCTGACCTGGTACAAGAG R: GGTAGATGCGTCTGAGTTOOAT
<i>Caspase 3</i>	F: TGAGCCATGGTGAAGAAGGAATAA R: CCCGGGTAAGAATGTGCATAAAT
<i>Caspase 8</i>	F: TCAACAAGAGCCTGCTGAAGATA R: GGAGAGTCCGAGATTGTCATTAC
<i>c-Flip</i>	F: ATGGCAGAGATTGGTGAGGATTT R: GCTCCTTGAACAGACTGCTTGTA
<i>β-Actin</i>	F: CATGTACGTTGCTATCCAGGC R: CTCCTTAATGTCACGCACGAT

Table 2 siRNA primer sequences used for siRNA interference.

Gene	Primer sequence (5'-3')
<i>FADD</i>	F: GCGAGCUGACCGAGCUCAA
	R: UUGAGCUCGGUCAGCUCGC