

Cystoscopy Results (N=106)		
Patency of prostatic Urethra (Likert Scale), n (%)	<b>Perfect</b>	<b>40 (37.7)</b>
	Good	36 (33.9)
	Discrete	20 (18.8)
	Sufficient	6 (5.6)
	Poor	4 (3.7)
Preservation of Verumontanum, n (%)	<b>100% Preserved</b>	<b>77 (72.6)</b>
	75% Preserved	22 (20.7)
	50% Preserved	4 (3.7)
	<50% Preserved	3 (2.8)
Residual Median Lobe, n (%)	12 (11.3)	
Fluffy tissue, n (%)	Abundant	10 (9.4)
	Moderate	9 (8.4)
	Few	9 (8.4)
	<b>Absence</b>	<b>80 (75.4)</b>
Mucosal flaps presence, n (%)	16 (15.0)	
Ureteral meatuses injury, n (%)	0 (0)	
Trigone appearance, n (%)	<b>Normal</b>	<b>86 (81.1)</b>
	Erythematous	20 (18.8)
	Sclerotic	0 (0)
Length of the cavity formed (mm), mean (SD)	24 (8.2)	
Distance between the apex and the sphincter (mm), mean (SD)	9.6 (5.3)	

Table 2 - morphological data of 106 patients underwent urethro-cystoscopy three months after surgery.