

		1 month	3 months	6 months	12 months
Qmax (ml/s)	N	109	109	109	109
	Baseline	8.42±2.81	8.42±2.81	8.42±2.81	8.42±2.81
	Follow up	19.73±9.77	18.30±5.57	17.98±7.83	17.10±7.83
	Change	+11.31±6.34	9.88±4.56	9.56±5.54	8.68±5.32
	% Change	+134.32	+117.34	+113.53	+103.08
	P value	<0.05	<0.05	<0.05	<0.05
PVR (ml)	Baseline	83.39±67.81	83.39±67.81	83.39±67.81	83.39±67.81
	Follow up	26.48±26.02	18.98±16.52	24.47±22.65	20.74±11.84
	Change	-56.91±33.76	-64.41±22.76	-58.92±41.27	-62.65±34.65
	% Change	-68.25	-77.24	-70.66	-75.13
	P value	<0.05	<0.05	<0.05	<0.05
IPSS (points)	Baseline	22 (17-26)	22 (17-26)	22 (17-26)	22 (17-26)
	Follow up	4 (2-6)	3 (2-5)	3 (2-6)	3 (2-5)
	Change	-18 (2-18)	-19 (2-19)	-19 (2-19)	-19 (2-19)
	% Change	-81.82	-86.36	-86.36	-86.36
	P value	<0.05	<0.05	<0.05	<0.05
IPSS-QoL (points)	Baseline	5 (4-5)	5 (4-5)	5 (4-5)	5 (4-5)
	Follow up	1 (0-2)	1 (0-1)	1 (0-1)	1 (0-1)
	Change	-4 (0-3)	-4 (0-3)	-4 (0-3)	-4 (0-3)
	% Change	-80.00	-80.00	-80.00	-80.00
	P value	<0.05	<0.05	<0.05	<0.05
SHIM (points)	Baseline	24 (20-25)	24 (20-25)	24 (20-25)	24 (20-25)
	Follow up	25 (20-25)	25 (22-25)	25 (21-25)	24 (20-24)
	Change	+1 (0-2)	+1 (0-2)	+1 (0-2)	0 (0-0)
	% Change	+4.00	+4.00	+4.00	
	P value	0.98	0.98	0.98	0.98
MSHQ-EjD (points)	Baseline	18 (17-20)	18 (17-20)	18 (17-20)	18 (17-20)
	Follow up	17 (16-20)	17 (16-20)	16 (16-20)	17 (16-20)
	Change	-1 (1-0)	-1 (1-0)	-2 (1-0)	-1 (1-0)
	% Change	-5.5	-5.5	-11.11	-5.5
	P value	0.45	0.87	0.45	0.45
ISI (points)	Baseline	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
	Follow up	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
	Change	0 (0-0)	0 (0-0)	0 (0-0)	0 (0-0)
	% Change	0	0	0	0
	P value	1	1	1	1

Table 3 - Postoperative variables at each timepoint through the 12-month follow-up (Qmax: maximum peak flow; PVR: post void residual; IPSS: international prostate symptom score; IPSS-QoL: IPSS Quality of Life; SHIM: sexual health inventory for men; MSHQ-EjD: men sexual health questionnaire-ejaculatory dysfunction; ISI: incontinence severity index)