

TABLE 4

Adjusted^a odds ratios (OR) and 95% confidence intervals (CI) of lifestyle and BMI among men and women experienced infertility in Finland 2011

	MEN					WOMEN				
	fertile n = 1677	OR	infertile n=201	OR ^b	95% CI	fertile n=1257	OR	infertile n=267	OR ^b	95% CI
	%		%			%		%		
BMI, OVERWEIGHT OR OBESE	68	1	70	1.30	0.9 - 1.89 p = 0.17	49	1	46	0.89	0.67 - 1.19 p = 0.44
SMOKING, daily or occasionally	27	1	21	0.81	0.55 - 1.21 p = 0.31	23	1	24	1.29	0.91 - 1.83 p = 0.15
SMOKING, high or average daily consumption of cigarettes^c	16	1	11	0.75	0.44 - 1.27 p = 0.28	10	1	11	0.99	0.62 - 1.59 p = 0.97
RISKY ALCOHOL USE (AUDIT-C men $\geq$6, women $\geq$5)^d	35	1	33	1.04	0.69 - 1.56 p = 0.85	26	1	20	0.83	0.56 - 1.21 p = 0.33
HIGH OR AVERAGE TOTAL ALCOHOL CONSUMPTION^d	81	1	91	2.01	1.11 - 3.67 p = 0.02	70	1	69	0.84	0.60 - 1.18 p = 0.33

^aOR adjusted for age, marital status, education and employment status
^bfertile men and women as reference groups
^c209 missing
^d729 missing