

TABLE 3

Age-adjusted prevalences and age-adjusted odds ratios (OR) and 95% confidence intervals (CI) of lifestyle and BMI among men and women experienced infertility in Finland 2011

	MEN					WOMEN				
	fertile n = 1677 %	OR	infertile n=201 %	OR ^a	95% CI	fertile n=1257 %	OR	infertile n=267 %	OR ^a	95% CI
BMI, OVERWEIGHT OR OBESE	68	1	70	1.29	0.88-1.89 p = 0.19	49	1	46	0.84	0.64 - 1.10 p = 0.21
SMOKING, daily or occasionally	27	1	21	0.71	0.48 - 1.03 p = 0.07	23	1	24	1.09	0.79 – 1.52 p = 0.59
SMOKING, high or average daily consumption of cigarettes^b	16	1	11	0.62	0.37 – 1.04 p = 0.07	10	1	11	0.86	0.54 – 1.36 p = 0.51
RISKY ALCOHOL USE (AUDIT-C men $\geq$6, women $\geq$5)^c	35	1	33	1.02	0.67 – 1.53 p = 0.94	26	1	20	0.73	0.50 – 1.06 p = 0.11
HIGH OR AVERAGE TOTAL ALCOHOL CONSUMPTION^c	81	1	91	2.49	1.39 – 4.45 p = 0.002	70	1	69	0.91	0.65 – 1.27 p = 0.58

^afertile men and women as reference groups
^b209 missing
^c729 missing