


a**Patient 1**

Case	A 73 year old male patient with a history of MPO-ANCA positive vasculitis was referred to a regional hospital in august 2020 with an episode of macro-hematuria, as well as fatigue, dizziness and progressive lower limb edema. Laboratory assessment showed acute kidney injury and elevated CRP levels. Chest imaging excluded relevant pathologies. Urinary analysis showed gross hematuria and macro-albuminuria. BVAS was elevated to 9 points. Suspecting a relapse of AAV, the patient was transferred to university hospital Hamburg-Eppendorf.		<u>organ involvement</u>																
AAV history	Initial diagnosis of AAV was made in 2017, with general symptoms, skin and kidney involvement. Kidney biopsy showed pauci-immune crescentic glomerulonephritis with small amounts of mesangial IgA deposition. Induction treatment consisted of PLEX, steroids and cyclophosphamide (cumulative CYC dose 5.0 g). Azathioprine was given for remission maintenance. After 7 months, azathioprine was discontinued due to severe side effects, and rituximab was started. Rituximab was well tolerated, and stable remission achieved, although full B cell depletion was not achieved during maintenance therapy with rituximab. Last dose RTX was given 6 months before re-admission with suspected AAV relapse.		 kidney skin																
Diagnostic workup at relapse	<table><tr><td>Hemoglobin [g/dl]</td><td>12.0</td></tr><tr><td>Leukocytes [10⁹/l]</td><td>7.8</td></tr><tr><td>Platelets [10⁹/l]</td><td>271</td></tr><tr><td>Creatinine [mg/dl]</td><td>4.25</td></tr><tr><td>eGFR [ml/min]</td><td>13</td></tr><tr><td>CRP [mg/l]</td><td>36</td></tr><tr><td>MPO-ANCA [U/ml]</td><td>22</td></tr><tr><td>B cells [%]</td><td>6 (34/μl)</td></tr></table>	Hemoglobin [g/dl]	12.0	Leukocytes [10 ⁹ /l]	7.8	Platelets [10 ⁹ /l]	271	Creatinine [mg/dl]	4.25	eGFR [ml/min]	13	CRP [mg/l]	36	MPO-ANCA [U/ml]	22	B cells [%]	6 (34/μl)	Kidney biopsy was performed, displaying a relapse of ANCA-GN with a high ARRS. Rapid biopsy immune profiling of the kidney biopsy revealed an enrichment of Th1/Tc1 and Th17/Tc17 cells in the kidney tissue.	
Hemoglobin [g/dl]	12.0																		
Leukocytes [10 ⁹ /l]	7.8																		
Platelets [10 ⁹ /l]	271																		
Creatinine [mg/dl]	4.25																		
eGFR [ml/min]	13																		
CRP [mg/l]	36																		
MPO-ANCA [U/ml]	22																		
B cells [%]	6 (34/μl)																		
Treatment considerations	Patient tolerated rituximab well, but full B cell depletion could not be achieved, despite appropriate RTX-dosage. Furthermore, the patient suffered a relapse while on RTX treatment. Urinary flow obstruction and the need of urinary diversion led the patient away from full dose CYC, while still giving consent to two doses of i.v. CYC in the inpatient setting. Rapid immune profiling showed an increase in Th1/Tc1 and Th17/Tc17 cells, suggesting suitability for an anti-IL12/IL-23 treatment.																		
Treatment	- Pulsed i.v. steroids (3 x 500 mg prednisolone), followed by an oral tapering, according to the PEXIVAS trial reduced dose regimen. By 6 months prednisolone was tapered to 5mg daily (cumulative prednisolone dose: 3900 mg). - Two i.v. pulses of cyclophosphamide (cumulative dosage 2g) were given in the inpatient setting.																		
Outcome	Combined treatment of ustekinumab, steroids and low dose CYC resulted in a rapid clinical improvement with resolution of fatigue, edema and dizziness. Creatinine rapidly improved and uACR declined, as well as ANCA-levels, CRP, and BVAS. At 6 months the patient had an elevated BVAS of 3 because of grade 1 hypertension and persistent albuminuria, mostly attributed to glomerular scarring.																		


b**Patient 2**

Case	A 52 year old male patient with known history of MPO-ANCA positive vasculitis with extensive organ manifestation was admitted to university medicine hospital Hamburg-Eppendorf, because of increasing creatinine and albuminuria after induction treatment with steroids (3 x 500 mg prednisolone and oral taper) and six pulses of i.v. cyclophosphamide (cumulative dose of 6.0 g; last dose given 4 weeks prior to admission). Initial manifestations (ENT, pulmonary, nervous and cutaneous) had improved since starting of induction treatment with steroids and CYC. BVAS was increased to 12 at admission. With suspected worsening ANCA-GN kidney biopsy was initiated.		organ involvement  ENT lung/heart kidney skin nervous system medical history - hyperlipidemia - postherpetic neuralgia													
AAV history	Initial diagnosis of AAV was made 02/2021 with general, renal, ENT, lung, nervous system, cutaneous, renal and heart manifestation. Treatment with i.v. prednisolone followed by an oral taper (according to the PEXIVAS reduced dose regimen), as well as pulsed i.v. cyclophosphamide (according to EUVAS scheme, cumulative dose 6.0 g) was initiated.															
Diagnostic workup at relapse	<table><tr><td>Hemoglobin [g/dl]</td><td>10.7</td></tr><tr><td>Leukocytes [10⁹/l]</td><td>6.5</td></tr><tr><td>Platelets [10⁹/l]</td><td>282</td></tr><tr><td>Creatinine [mg/dl]</td><td>2.27</td></tr><tr><td>eGFR [ml/min]</td><td>32</td></tr><tr><td>CRP [mg/l]</td><td>5</td></tr><tr><td>MPO-ANCA [U/ml]</td><td>29</td></tr></table>	Hemoglobin [g/dl]		10.7	Leukocytes [10 ⁹ /l]	6.5	Platelets [10 ⁹ /l]	282	Creatinine [mg/dl]	2.27	eGFR [ml/min]	32	CRP [mg/l]	5	MPO-ANCA [U/ml]	29
Hemoglobin [g/dl]	10.7															
Leukocytes [10 ⁹ /l]	6.5															
Platelets [10 ⁹ /l]	282															
Creatinine [mg/dl]	2.27															
eGFR [ml/min]	32															
CRP [mg/l]	5															
MPO-ANCA [U/ml]	29															
Treatment considerations	Cyclophosphamide treatment along with steroids could not achieve remission and renal manifestation showed a progressive disease with active glomerular lesions. COVID-19 pandemic was still on the rise, raising concerns over rituximab treatment. Rapid immune profiling of the kidney showed an enrichment of Th1/Tc1 and Th17/Tc17 cells.															
Treatment	- Pulsed steroids (3x500mg prednisolone), followed by an oral tapering. By 6 months prednisolone was tapered to 5mg daily (cumulative prednisolone dose: 3755 mg). - Two i.v. pulses of cyclophosphamide (cumulative dosage 2.0 g) were given. - Ustekinumab 90mg s.c. was administered at weeks 0, 4, 12, and 24. - Azathioprine (1 mg/kg) was started at week 22 after beginning of induction therapy as additional maintenance therapy.															
Outcome	Combined treatment of ustekinumab, steroids and low dose CYC resulted in a rapid clinical improvement with reduction in creatinine, uACR and ANCA-levels (16 U/ml). At 6 months the patient had an elevated BVAS of 2, because of persistent microalbuminuria, undulating around 200 mg/day.															

c**Patient 3**

Case	A 32 year old male with a history of PR3-ANCA positive vasculitis was re-admitted to university hospital Hamburg-Eppendorf, because of fever, night sweats, weight loss, and progressive dyspnea with hemoptysis. Four weeks before admission the patient was diagnosed with an AAV relapse- rituximab (two pulses of 1g) and pulsed steroids as well as an oral tapering regimen were started. Clinical examination at re-admission showed moist crackles in chest auscultation. BVAS was 13 at that time.	<u>organ involvement</u>  ENT lung kidney <u>medical history</u> - no chronic diseases																
AAV history	Initial diagnosis of AAV was made 2019 and treatment with i.v. prednisolone followed by an oral taper (according to the PEXIVAS reduced dose regimen) as well as rituximab was initiated. After two doses (1g each) of rituximab, the patient was lost to follow up. In 09/2022 aforementioned relapse with pulmonary and ENT manifestation occurred and pulsed prednisolone as well as re-induction with rituximab (two times 1g) was started. Initially the patient's symptoms improved and he was discharged.																	
Diagnostic workup at relapse	<table><tr><td>Hemoglobin [g/dl]</td><td>8.2</td></tr><tr><td>Leukocytes [10⁹/l]</td><td>1.7</td></tr><tr><td>Platelets [10⁹/l]</td><td>435</td></tr><tr><td>Creatinine [mg/dl]</td><td>1.88</td></tr><tr><td>eGFR [ml/min]</td><td>46</td></tr><tr><td>CRP [mg/l]</td><td>190</td></tr><tr><td>PR3-ANCA [U/ml]</td><td>101</td></tr><tr><td>B cells [%]</td><td>0 (1/μl)</td></tr></table>	Hemoglobin [g/dl]	8.2	Leukocytes [10 ⁹ /l]	1.7	Platelets [10 ⁹ /l]	435	Creatinine [mg/dl]	1.88	eGFR [ml/min]	46	CRP [mg/l]	190	PR3-ANCA [U/ml]	101	B cells [%]	0 (1/μl)	CT scans and bronchoscopy showed progressive DAH. Urinary sediment showed extensive acanthocyturia. Kidney biopsy was performed and displayed pauci-immune crescentic glomerulonephritis with active lesions, compatible with ANCA-GN, with a low ARRS. Rapid immune-profiling of the kidney biopsy revealed an enrichment of Th1/Tc1 and Th17/Tc17 cells in the kidney tissue.
Hemoglobin [g/dl]	8.2																	
Leukocytes [10 ⁹ /l]	1.7																	
Platelets [10 ⁹ /l]	435																	
Creatinine [mg/dl]	1.88																	
eGFR [ml/min]	46																	
CRP [mg/l]	190																	
PR3-ANCA [U/ml]	101																	
B cells [%]	0 (1/μl)																	
Treatment considerations	Severe leukopenia contradicted full dose cyclophosphamide re-induction treatment. Patient was also concerned of cyclophosphamide treatment-related adverse effects, and initially rejected an i.v. treatment. B cells were fully depleted, as rituximab was given four weeks prior. Rapid immune profiling of the kidney showed an enrichment of Th1/Tc1 and Th17/Tc17 cells.																	
Treatment	- Pulsed steroids (3x250mg prednisolone), followed by an oral tapering. By 6 months prednisolone was tapered to 5mg daily (cumulative prednisolone doses 3520 mg). - Oral cyclophosphamide was given as the patient was initially rejecting i.v. therapy. After repeated shared decision talks, the patient gave consent to one more i.v. pulse under close monitoring of WBC (cumulative dose: 500 mg p.o. and 1000mg i.v.). - Ustekinumab 90mg s.c. was administered at weeks 0, 4, 12, and 24. - Starting at week 16 after re-induction therapy AZA was started for additional maintenance therapy and eventually switched to low dose MMF (2x500 mg) because of AZA side effects.																	
Outcome	Combined treatment of ustekinumab, steroids and low dose CYC resulted in a rapid clinical improvement with suspended night sweats, fever and hemoptysis, a reduction in creatinine, ANCA-levels, and BVAS. At 6 months the patient had an elevated BVAS of 2, because of persisting albuminuria. Antiproteinuric treatment with ACEi was not well tolerated and could only be given at the lowest dose.																	

d**Patient 4**

Case	A 72 year old female with a history of MPO-ANCA positive vasculitis (renal limited disease) and myelodysplastic syndrome was admitted to university medicine hospital Hamburg-Eppendorf, because of acute kidney injury and reduced general condition. The patient was receiving rituximab as remission maintenance, with the last dose given 5 months prior to hospital admission. BVAS was 13 at admission. With suspected relapse of ANCA-GN diagnostic workup was initiated.		organ involvement  kidney <u>medical history</u> - myelodysplastic syndrome - osteoporosis - Hashimoto's thyroiditis															
AAV history	Initial diagnosis of AAV was made in august 2020 with a renal limited phenotype. Treatment with i.v. prednisolone followed by an oral taper (according to the PEXIVAS reduced dose regimen) as well as pulsed i.v. cyclophosphamide (cumulative dose 3600 mg) was initiated. After remission induction azathioprine was given for remission maintenance. Eventually, azathioprine had to be discontinued because of side effects and rituximab at doses of 500 mg/6 months was initiated in 05/2022.																	
Diagnostic workup at relapse	<table><tr><td>Hemoglobin [g/dl]</td><td>8.9</td></tr><tr><td>Leukocytes [10⁹/l]</td><td>3.0</td></tr><tr><td>Platelets [10⁹/l]</td><td>264</td></tr><tr><td>Creatinine [mg/dl]</td><td>3.42</td></tr><tr><td>eGFR [ml/min]</td><td>13</td></tr><tr><td>CRP [mg/l]</td><td>28</td></tr><tr><td>MPO-ANCA [U/ml]</td><td>>134</td></tr><tr><td>B cells [%]</td><td>2 (20/μl)</td></tr></table>	Hemoglobin [g/dl]		8.9	Leukocytes [10 ⁹ /l]	3.0	Platelets [10 ⁹ /l]	264	Creatinine [mg/dl]	3.42	eGFR [ml/min]	13	CRP [mg/l]	28	MPO-ANCA [U/ml]	>134	B cells [%]	2 (20/μl)
Hemoglobin [g/dl]	8.9																	
Leukocytes [10 ⁹ /l]	3.0																	
Platelets [10 ⁹ /l]	264																	
Creatinine [mg/dl]	3.42																	
eGFR [ml/min]	13																	
CRP [mg/l]	28																	
MPO-ANCA [U/ml]	>134																	
B cells [%]	2 (20/μl)																	
Treatment considerations	Leukopenia of 3.0 x10 ⁹ /l with known myelodysplastic syndrome under high dose erythropoietin therapy contradicted full dose cyclophosphamide re-induction therapy. Retrospective the patient showed a rise in MPO-ANCAs in 01/2023 while B cells were depleted (0% / 1/μl). Rapid immune profiling of the kidney showed an enrichment of Th1/Tc1 and Th17/Tc17 cells.																	
Treatment	- Pulsed steroids (3 x 500 mg prednisolone), followed by an oral tapering regimen according to the PEXIVAS study reduced dose arm. By 6 months prednisolone was tapered to 5mg daily (cumulative prednisolone dose: 4285 mg). - Three pulses of i.v. cyclophosphamide were given (cumulative dose of 1800mg). - Ustekinumab 90mg s.c. was administered at weeks 0, 4, 12, and 24. - Low dose MMF (2 x 500 mg) was started at week 16 after beginning of induction therapy as additional maintenance therapy.																	
Outcome	Combined treatment of ustekinumab, steroids and low dose CYC resulted in a rapid clinical improvement with substantial improvement of general condition, reduction in creatinine, uACR, ANCA-levels, and BVAS reduction to 5. At 6 months the patient had an elevated BVAS of 5, because of albuminuria and eumorphic hematuria.																	

Supplemental Figure 5: Detailed case vignettes for each patient of the ustekinumab treatment cohort.

Case vignettes illustrating a brief case description, history, diagnostic workup, as well as treatment considerations, treatment and outcomes for each patient of the ustekinumab treatment cohort. **a** Patient 1. **b** Patient 2. **c** Patient 3. **d** Patient 4. BVAS, Birmingham Vasculitis Activity Score. AAV, ANCA-associated vasculitis. PLEX, therapeutic plasma exchange. CYC, cyclophosphamide. ARRS, ANCA renal risk score (21). RTX, rituximab. uACR, urinary albumin to creatinine ratio. OSA, obstructive sleep apnoea. ENT, ear nose throat. DAH, diffuse alveolar hemorrhage. MMF, mycophenolate mofetil. ACEi, angiotensin-converting enzyme inhibitor.