

Two new psychrotolerant *Massilia* species inhibit plant pathogens *Clavibacter* and *Curtobacterium*

Ivo Sedláček^{1*}, Pavla Holochová¹, Karel Sedlář², Eva Staňková¹, Ondrej Šedo³, Stanislava Králová¹, Mohammad Umair², Vendula Koublová¹, Pavel Švec¹

¹Department of Experimental Biology, Czech Collection of Microorganisms, Faculty of Science, Masaryk University, Kamenice 5, 625 00 Brno, Czech Republic; ivo@sci.muni.cz (I.S.); pavlah@sci.muni.cz (P.H.); evickakroupova@seznam.cz (E.S.); kralova.s@sci.muni.cz (S.B.K.); 436839@mail.muni.cz (V.K.); mpavel@sci.muni.cz (P.Š.)

²Department of Biomedical Engineering, Faculty of Electrical Engineering and Communication, Brno University of Technology, Technická 12, 616 00 Brno, Czech Republic; sedlar@vut.cz (K.S.); 253834@vut.cz (M.U.)

³Central European Institute of Technology, Masaryk University, Kamenice 5, 625 00 Brno, Czech Republic; sedo@post.cz

***Correspondence:** ivo@sci.muni.cz; Tel.: +420-549-496-922

Supplementary material




Figure S1. Dendrograms based on cluster analysis of *Eco*RI ribotype patterns obtained using the RiboPrinter identification system from *M. scottii* sp. nov., *M. pseudviolaceinigra* sp. nov. strains, and closest *Massilia* spp. type strains. The dendrogram was calculated with Pearson's correlation coefficients by the UPGMA clustering method (r , expressed as percentage similarity values).




Figure S2 A. Polar lipid profile of strain P5043^T *M. scottii* sp. nov., after two -dimensional thin layer chromatography. Abbreviations: DPG, diphosphatidylglycerol; PG, phosphatidylglycerol; APL, aminophospholipid; AL, aminolipid; GL, glycolipid; PL, phospholipid; L lipids




Figure S2 B. Polar lipid profile of strain P3689^T *M. pseudoviolaceinigra* sp. nov. after two -dimensional thin layer chromatography. Abbreviations: DPG, diphosphatidylglycerol; PG, phosphatidylglycerol; APL, aminophospholipid; AL, aminolipid; GL, glycolipid; PL, phospholipid; L lipids .