


Fig. S1 Generalized protein folding protocol



* Solubilization buffers (S₁₋₃)

S₁: 6 M GuHCl, 50 mM Tris-HCl, 0.1 M NaCl, pH 8

S₂: 8 M Urea, 50 mM Tris-HCl, 0.1 M NaCl, pH 8

S₃: 6 M GuHCl, 50 mM Tris-HCl, 2mM EDTA, 5 mM DTT, pH 8


** Folding buffers (F₁₋₃)

F₁: 50 mM Tris-HCl, 100 mM NaCl, 7% glycerol, pH 8

F₂: 50 mM Tris-HCl, 100 mM NaCl, 2% glycerol, pH 8

F₃: 100 mM Tris-HCl, 2 mM EDTA, 1mM GSSG, 0.8 mM GSH, pH 8.5

Fig. S2



	α conotoxin 1.35 kDa	λ conotoxin 1.24 kDa	rFaxiator 12 kDa	PLA2 17 kDa	GFP 27 kDa	HRPC 34 kDa	rLuc 36 kDa	FFL 60 kDa	HSA 66 kDa	AP 52 kDa	Omp2a 39 kDa	p53 43.7 kDa
PI of protein	7.8	8	8.3	4.8	5.80	6.30	5.86	6.69	5.67	4.63	4.3	6.33
Molecular Volume	1116 Å ³	1116 Å ³	15615 Å ³	32509 Å ³	: 41198 Å ³	41198 Å ³	43585 Å ³	70007 Å ³	80426 Å ³	64545 Å ³	46001 Å ³	52819 Å ³
Charge at pH 7.0	0.8	1	3.5	-5.5	-6.2	-1.2	-13.4	-1	-15.7	-38.6	-22.6	-3.3
Disulfide (#)	2	2	3	7	0	4	0	0	17	2	0	4
Optimal tES	tES(-)	tES(-/+)	tES(-)	tES(+)	tES(+)	tES(+)	tES(+)	tES(-/+)	tES(+)	tES(+)	tES(+)	tES(+,-/+)
tES:POI encapsulation (max ratio)	N/A	N/A	1:3	N/A	N/A	1:1	1:2	N/A	N/A	N/A	N/A	N/A
tES:POI (max stabilization)	60:150	60:150	60:15	60:15	60:10	60:10	60:10	60:5	60:5	60:5	60:5	60:10
tES:POI (max fxnl yeild)	1:60	1:60	1:6	1:6	1:4	1:4	1:4	1:8	1:8	1:8	1:8	1:4
POI multimer	No	No	No	No	No	No	No	No	No	Yes	Yes	Yes
Source	Molluscs	Molluscs	Reptiles	Reptiles	Cnidarians	Plants	Cnidarians	Arthropods	Human	Arthropods	Bacteria	Human
Percentage Secondary Structure	N.A	N.A	β strand : 14.9	β strand : 5.9	β strand : 45	β strand : 1.9	β strand : 16.5	β strand : 19.3	β strand : 0	β strand : 16	β strand : 55.6	β strand : 20.6
Cysteine Density (%)	33	40	9	10	0.8	2.5	1	0.9	6	0.8	0.6	2.5
pl	8.89	7.18	8.58	5.42	5.8	5.66	5.76	6.72	5.67	4.63	4.57	6.42

Fig. S2| Comparative characteristics of monomeric and multimeric POIs




Fig. S3 | Analysis of encapsulation process with increasing POI concentration using Analytical Ultracentrifuge (AUC): Residuals plots (a,b,c) and absorbance profiles of (d) tES(+):F116H, (e) tES(+):F116H:HSA, (f) tES(+):F116H:rLuc, (g) tES(-):F116H:rFasxiator, (h) tES(+):F116H:λ conotoxin at optimal molar ratios registered at 280 nm, in time intervals of 7 min for total experimental time of approximately 7 h. AUC profile for λ conotoxin shows decreasing sedimentation of tES encapsulated conotoxin and increasing presence of free conotoxin as the run progresses.




Fig. S4 | Chromatographic profiles of α conotoxin folded in presence and absence of tES: Chromatographic profiles for α conotoxin folded in-vitro without tES or within tES(+)/F116H, tES(+/-)/F116H or tES(-)/F116H respectively with different molar ratios of tES subunits: α conotoxins from (a) 60:37.5, (b) 60:75, (c) 60:112.5, (d) 60:150, (e) 60:225 and (f) 60:300 respectively. Relative yield of globular (active) conformation when conotoxin is folded in presence and absence of tES is shown as a histogram.




Fig. S5| Chromatographic profiles of λ conotoxin folded in presence and absence of tES: Chromatographic profiles for λ conotoxin folded in-vitro without tES or within tES(+)F116H, tES(+/-)F116H or tES(-)F116H respectively with different molar ratios of tES subunits: λ conotoxins from (a) 60:37.5, (b) 60:75, (c) 60:112.5, (d) 60:150, (e) 60:225 and (f) 60:300 respectively. Relative yield of ribbon (active) conformation when conotoxin is folded in presence and absence of tES is shown as a histogram.




Fig. S6 | Optimization of molar ratios of tES subunits to POI: Size-exclusion profiles (upper panel) of (a) tES(+)-F116H:HSA (b) tES(+)-F116H:rLuc (c) tES(-)-F116H:rFasxiator with different molar ratios of tES subunits and POI. Each fraction of size-exclusion chromatography was analysed for POI activity for all molar ratios tested (lower panel). POI activity coincides with tES peak, suggesting its encapsulation inside tES assembly. All experiments were performed in triplicates, error bar represents $\pm$ standard deviation. (d) Charge matching of tES-F116H and the measured zeta potential of the POI predict optimal folding. Combinations with rank order highest functional yield are shown for nine monomeric POI's. tES-F116H(+), tES-F116H(+/-), tES-F116H(-) and POI-alone are indicated as (+), (+/-), (-) and h6, respectively.




Fig. S7 | Characterization of tES nanoparticles using Dynamic Light Scattering (DLS): DLS experiments of tES nanoparticles showed no change in the hydrodynamic diameter (12 nm approx.) of POI encapsulated tES suggesting internalization of POI within tES.




Fig. S8 | Characterization of tES nanoparticles using Transmission Electron Microscopy (TEM): TEM experiments of tES nanoparticles showed no change in the morphology and hydrodynamic diameter (12 nm approx.) of POI encapsulated tES suggesting internalization of POI within tES.




Fig. S9 | Mass spectrometry profiles of globular conformation of α conotoxin folded in presence and absence of tES: Raw mass/charge spectrum (m/z) (upper panel) and deconvoluted mass spectrum showing inferred molecular masses (lower panel) of α conotoxin folded in-vitro in presence of (a) tES(+)F116H, (b) tES(+/-)F116H and (c) tES(-)F116H and in (d) absence of tES were determined by QTOF. The protein is predominantly +3 charged, appearing with m/z of 451 amu. Protein has inferred molecular mass of 1350.48 amu which closely matches with the predicted molecular mass of 1350.92 amu. In all cases, molecular weight of the oxidized α conotoxin shows a reduction of four mass units, reflective of the formation of the two disulfide bridges. m/z at 121 and 922 were from mass calibrant. All experiments were performed in triplicates.




Fig. S10 | Mass spectrometry profiles of ribbon conformation of λ conotoxin folded in presence and absence of tES: Raw mass/charge spectrum (m/z) (upper panel) and deconvoluted mass spectrum showing inferred molecular masses (lower panel) of λ conotoxin folded in-vitro in presence of (a) tES(+)-F116H, (b) tES(+/-)-F116H and (c) tES(-)-F116H and in (d) absence of tES were determined by QTOF. The protein is predominantly +3 charged, appearing with m/z of 413 amu. Protein has inferred molecular mass of 1236.46 amu which closely matches with the predicted molecular mass of 1236.90 amu. In all cases, molecular weight of the oxidized λ conotoxin shows a reduction of four mass units, reflective of the formation of the two disulfide bridges. m/z at 121 and 922 were from mass calibrant. All experiments were performed in triplicates.




Fig. S11 | Determining yield of in-vitro folded POI in presence and absence of shells: SDS-PAGE gel of in-vitro folded POI in absence of tES, in-vitro folded POI in presence of tES and POI released and purified from tES after in-vitro folding for (a) rFasxiator, (b) PLA₂, (c) GFPuv, (d) HRPc, (e) rLuc, (f) FFL and (g) HSA.




Fig. S12 | Analysis of encapsulation process with different internal charge of tES-F116H using Analytical Ultracentrifuge (AUC): Absorbance profiles (upper panel) and residuals plots (lower panel) of (a) HSA, (b) rLuc and (c) rFasxiator encapsulated within (i) tES(+)F116H, (ii) tES(+/-)F116H and (iii) tES(-)F116H respectively at optimal molar ratios registered at 280 nm , in time intervals of 7 min for total experimental time of 7 h approximately.