


Shapiro-Wilk Test Calculator

Reporting Shapiro-Wilk test in APA Format

The Shapiro-Wilk test showed a significant departure from normality, $W(15) = .81, p = .005$

Parameter	Value
P-value	0.004639
W	0.8078
Sample size (n)	15
Average ($\bar{x}$)	90.26
Median	91.9
Sample Standard Deviation (S)	3.3481
Sum of Squares	156.936
b	11.2594
Skewness	-1.5904
Skewness Shape	 Asymmetrical , left/negative (pval=0.006)
Excess kurtosis	2.1706
Kurtosis Shape	 Potentially Mesokurtic , normal like tails (pval=0.053)
Outliers	81.5



Shapiro-Wilk-test, using tables as distribution (right-tailed)

Since $n \leq 50$ we used the Shapiro-Wilk tables to calculate the p-value.

The normal distribution chart is only for visualization.

1. H_0 hypothesis

Since $p\text{-value} < \alpha$, we reject the H_0 .

It is assumed that the data is not normally distributed.

In other words, the difference between the data sample and the normal distribution is big enough to be statistically significant.

2. P-value

The p-value equals **0.004639**, ($P(x \leq 2.6016) = 0.9954$). It means that the chance of type I error (rejecting a correct H_0) is small: 0.004639 (0.46%). The smaller the p-value the more it supports H_1 .


3. Test statistic

The test statistic **W** equals **0.8078**, which is not in the 95% region of acceptance: [0.8815, 1].

4. Effect size


The observed effect size **KS - D** is **large, 0.2541**. This indicates that the magnitude of the difference between the sample distribution and the normal distributions is large.

Distribution: Normal(mean:0, std:1)




Significance level (α)

Histogram



Q-Q Plot



-5%

-7%

Trusted US Based Suppli
MSE Supplies