

Sero-epidemiology and risk factor investigation for exposure to Japanese encephalitis virus in swine populations reared in climatically diverse regions of India

Journal name: Tropical Animal Health and Production

Akash Balasaheb Mote^a, Balbir B. Singh^b, Deepa Mehta^a, Sagar M. Patel^a, M Dhanalakshmi^c, M. Suman Kumar^a, K. N. Bhilegaonkar^a, Himani Dhanze^{a*},

^aDivision of Veterinary Public Health, ICAR – Indian Veterinary Research Institute, Uttar Pradesh, India.

^bCentre for One Health, Guru Angad Dev Veterinary & Animal Sciences University, Punjab, India

^cVeterinary College and Research Institute, Orathanadu, Tamil Nadu Veterinary and Animal Sciences University, Chennai, India.

*Corresponding author. Tel.: +919457958571

E-mail address: hdhanze@yahoo.co.in

Supplementary file S2

Table S2 Frequency distribution of samples categorized by state, district, sampling premise and the diagnostic test

Variable Name	Category	Count	Per cent
State	Assam	272	22.57
	Maharashtra	315	26.14
	Uttarakhand	239	19.83
	Uttar Pradesh	379	31.45
	Total	1205	100
District	Azamgarh	84	6.97
	Baksa	48	3.98
	Ballia	69	5.73
	Dehradun	49	4.07
	Deoria	70	5.81
	Gorakhpur	81	6.72
	Haridwar	100	8.30
	Kamrup Metropolitan	147	12.20
	Mumbai	105	8.71
	Nashik	150	12.45
	Satara	60	4.98

	Siddharth Nagar	75	6.22
	Sonitpur	77	6.39
	Udham Singh Nagar	90	7.47
	Total	1205	100
Total seropositive	Negative	846	70.21
	Positive	359	29.79
	Total	1205	100
IgM positive	Negative	959	79.59
	Positive	246	20.41
	Total	1205	100
IgG positive	Negative	993	82.41
	Positive	212	17.59
	Total	1205	100
Premise	Backyard Farm	945	78.42
	Established Farm	60	4.98
	Slaughterhouse	200	16.60
	Total	1205	100
