

Table 1. Demographics and clinical characteristics of participants (mean $\pm$ standard deviation).

	Methylphenidate		<i>p</i>	Atomoxetine		<i>p</i>
	Cognitive Training + Medication treatment	Medication treatment		Cognitive Training + Medication treatment	Medication treatment	
Male/female	24/4	29/4	1.000	29/3	29/2	1.000
Age (years)	9.9 $\pm$ 1.3	9.9 $\pm$ 1.7	0.906	8.9 $\pm$ 1.3	9.1 $\pm$ 1.6	0.627
IQ	109.6 $\pm$ 14.8	109.8 $\pm$ 16.3	0.957	106.9 $\pm$ 13.2	110.7 $\pm$ 12.3	0.240
ADHD-RS total (baseline)	32.6 $\pm$ 10.3	31.9 $\pm$ 7.1	0.753	35.5 $\pm$ 7.5	32.8 $\pm$ 6.4	0.133
Inattention (baseline)	18.8 $\pm$ 4.9	18.7 $\pm$ 3.3	0.936	20.1 $\pm$ 4.1	18.8 $\pm$ 3.4	0.189
Hyperactivity/impulsivity (baseline)	13.9 $\pm$ 6.4	13.2 $\pm$ 5.4	0.684	15.4 $\pm$ 4.9	14.0 $\pm$ 4.6	0.247
ADHD-RS total (follow-up)	23.1 $\pm$ 7.3	22.3 $\pm$ 7.5	0.662	23.0 $\pm$ 7.1	25.0 $\pm$ 6.0	0.248
Inattention (follow-up)	13.3 $\pm$ 3.5	13.1 $\pm$ 4.4	0.878	12.8 $\pm$ 3.2	14.0 $\pm$ 3.1	0.164
Hyperactivity/impulsivity (follow-up)	9.9 $\pm$ 4.3	9.2 $\pm$ 4.7	0.563	10.2 $\pm$ 5.4	11.0 $\pm$ 4.1	0.506

