

Unveiling genetic diversity in jute mallow (*Corchorus spp.*): morphological clustering reveals distinctive traits among accessions from Africa and Asia

M. Pamela Afokpe^{1,2,3}, Serge Ologou^{2,3}, S. Romaric Kouihou⁴, Simon Jan de Hoop², Sognigbé N'Danikou^{3,5,6}, Enoch G. Achigan-Dako³, M. Eric Schranz^{1*}

1 Biosystematics department, Wageningen University & Research, Wageningen, The Netherlands
 2 Hortigenetics Research (S.E. Asia) Ltd, Chiangmai, Thailand
 3 Genetics, Biotechnology and Seed Science Unit (GBioS), Laboratory of Crop Production, Physiology and Plant Breeding (PAGEV), Faculty of Agricultural Sciences (FSA), University of Abomey-Calavi, Abomey-Calavi, Republic of Benin
 4 BASF Innovation Center Gent, Belgium
 5 Eastern and Southern Africa, World Vegetable Center, P.O. Box 10, Duluti, Arusha, Tanzania
 6 Ecole d'Horticulture et d'Aménagement des Espaces Verts (EHAEV), Université Nationale d'Agriculture (UNA), BP 43, Kétou, Republic of Benin
 *Corresponding author: eric.schranz@wur.nl

Supplemental File 1: Origin of the 300 *Corchorus* accessions collected, grown and analyzed in this study

